

Isoelectric, Lewis Dot

Wednesday, January 03, 2007
9:03 AM

Reading for this week Ch. 10(10.1-10.7) , 10.8-10.9

Prof. Ken Spears

Wed 1400-1600 TA hours Nano building.

Every Thurs, quiz

Every Wed night, review sessions

1st quiz on Jan 11

Chemistry (Ken's try)

Why Lewis Dot Formulas for localized bonding?

Valence electrons of common atoms are shared to create an octet around atoms (except H). Both atoms compare

Formulas usually have central atom listed 1st

Sum all valence electrons including charges

Put a tentative single bond between all bonded atoms

Isoelectronic (CO , CN^- , N_2) Count them & add up valence e^-

Rearrange as needed to satisfy octet rule: lone pairs, multi. bonds
(1st, 2nd row use octet C, N, O, F always)

Elmt $\geq$ 3rd row can exceed octet rule.

BF_3




BeCl_2

ClF_3

$\text{B} = 3$

$\text{F} = 7 \cdot 3$

$3 + 21 = 24$




Lewis Dot structures w/ unpaired electrons
Molecular fragments have unpaired e^- 's.

CH_3 , CH_2

... stable molecules ...

very few stable molecules w/ unpaired e^- 's.
 NO , NO_2 (dimers)



This is actually bent




What improvements can be made to electron dot structures?

You have multiple structures.

Assume e^- can create an average bond. The electrons are actually delocalized. A pictorial method of showing is resonance hybrids.

Resonance Structures


Ozone, nitrogen dioxide, sulfate ion, etc.



Electron dot formulas often put e^- on an atom that are improbable. Atoms have different nuclear charges therefore diff ability to attract e^- in a bond. Electronegativity.

Invent a scheme of placing e^- that makes fewer mistakes. Counting scheme called formal charge.

Polar covalent bonds & electrostatic potential maps.



From a big distance away, it looks neutral

The closer you get, the easier it is to see imbalances

Qualitative Uses of Electronegativity

Organize the bonding trends from Ionic to Covalent

Electronegativity $\nearrow$

Size/physical
 stuff


Charge/attraction

dec ↓

inc ↓ $\xrightarrow{\text{dec}}$


dec ↓ $\xrightarrow{\text{inc}}$

Formal charge = # electrons - # lone pair - $\frac{1}{2}$ # bond pair



$$\text{FC}(\text{O}) = 6 - 4 - \frac{1}{2}(4) = 0$$

$$\text{FC}(\text{N}) = 5 - 0 - \frac{1}{2}(8) = +1$$






Lewis Dot, VSPER



Thursday, January 04, 2007
9:02 AM

Sum of FC is overall charge
FC should be as small as possible
Negative FC usually on most electronegative elmts
FC of same sign on adjacent atoms is unlikely.

what is the most likely structure, ~~FN~~ FNO or NFO?



	O(a)	O(b)	O(c)	O(d)	S
#valence	6	6			
#LP	-6	-4			
$\frac{1}{2} \cdot \# \text{ bonds}$	$-\frac{1}{2} \cdot 2$	$-\frac{1}{2} \cdot 4$			
Sum = FC	-1	0			







Used in airbags, exothermic rxn

Can we use Lewis Dot structures for 3D shape information?

Molecules are not 2D

Double bond isomers have no explanation



VSER Valence Shell Electron Pair Repulsion Model

1. Draw structure (Lewis)

2. Sum # bonds & # lone pairs around each atom (in vid.)

3. Pick appropriate e^- pair geometry around central atom & choose molecular shape

4.

Steric # lets us predict shape of molecule



$H_2O \rightarrow$ repels itself. δ^- & δ^+ forces

3 atoms
bonded to
central
atom

SN 3 makes Trigonal planar

SN 4 makes Tetrahedral / trigonal pyramidal

SN 5 makes T-shaped

See-saw is when



Properties of Crystals, Lab Procedure

Friday, January 05, 2007
8:59 AM

Bring a can to lab

What are some properties of crystals?

- Atoms arranged in a reg. pattern
- You can grow them
- Often translucent
- Can be different colors

What do you do first?


- Sand can

Then - Dissolve in KOH

Procedure for synthesis of $KAl(SO_4)_2 \cdot 12H_2O$



- Sand can
- Dissolve in KOH (long step)
- Discuss safety issues + do check-in
- Filter out solid
- Add H_2SO_4
- Put in ice bath (bc you're trying to make crystals)
- Vacuum filter crystals to get rid of excess aq. soln

When you add KOH, can will dissolve & gas will evolve ($H_2(g)$)






In lab manual


When we add H_2SO_4 , a precipitate forms, then dissolves



We use H_2SO_4 + not some other acid bc we want potassium aluminum sulfate. so we need H_2SO_4 + KOH



We need an ice bath for the product to form.
Things are less soluble at colder temps.



Mass of Alum? $466.26 \text{ g} = M_{\text{overall}}$

1 g aluminum is what we use

$$\frac{1}{26.98} = .037 \text{ moles}$$

Theoretical yield = 17.5 g

$$\left(\frac{\text{Actual yield}}{\text{Theoretical}} \right) 100 = \% \text{ yield}$$

Growing a single crystal


- Make a saturated soln using the crystals
- Filter into a small vial
- Observe next week

You can see the structure of a crystal by shining x-rays on them

Alum crystals go in white containers, leave aluminum can in lab,
filter paper, weighing paper → yellow bags, H_2SO_4 & KOH can go
in the sink if it's neutralized.

All lab reports due Thurs, Jan 18 at 6 pm

Online Safety Quiz





e- Pair Geometry

Monday, January 08, 2007
8:58 AM

VSEPR

1. Draw Lewis Structure
2. Sum # bonds + # lone pairs around each atom
- 3.


The effect of repulsion between nonbonding electrons + bond electrons
electron pair geometry



Effect of Non-bonding e⁻

Bond & decreases as # L.P. ↑ b/c e⁻ in a bond are attracted to nucleus

$LP1/LP2 > LP/Bond > Bond/Bond$



L.P. always placed in equatorial position

5 e⁻ pairs can make linear molecular geometry.



5 e⁻ pairs can give T shape.

6 e⁻ pairs where square pyramidal is distorted by L.P. < 90°.




DEMO

Eberhard's special purple plume

NI₃ explosions by touch sensitive detonation.

Ammonia + NI₃ in crystals NH₃ · NI₃

source of exothermic energy evolution?



exothermic


↑ purple plume

Dipole moment


polar covalent bonds can have displaced charges due to electronegativity differences electrostatic potential plots w/ a + test charge

$\delta^+ \text{H} - \text{Cl} \delta^-$ Partial charge δ , distance between partial charge
 $M = \delta \cdot d$ $= d.$


Field on



it would take a lot to orient them



CCl₄, nonpolar no net dipole moment




CHCl₃, polar b/c net dipole moment

Tetrahedral molecules can fit inside a cube. 109.5° of AX₄ tetrahedron.



tetrahedron.



Diff of electronegativity makes bigger dipole moment

Orbitals



Tuesday, January 09, 2007
9:00 AM

Average bond lengths

Memorize some #s, & decide if you go from single to triple bonds they get short

You can use these to come up w/ radii. Covalent radii, predict new bond lengths.

Bond energies are energies needed to break into atoms.



reactants $\rightarrow$ separate atoms $\rightarrow$ products $\rightarrow$ formation

$$\Delta H = \sum BE(\text{reactants}) - \sum BE(\text{products})$$

Break everything up & put it back together


How atomic orbitals define molecular shapes

Consider Be atoms in gas phase surrounded by H atoms
Unpaired e^- of H very reactive.





H atoms trolling for atoms w/ unpaired e^- ?

H atoms trolling for atoms that can't resist temptation?



Why doesn't carbon only make 2 bonds? Lewis Dot structure rules imply 4 e^- .



Sp orbitals form each having a huge lobe, pointing off directions




Little Angels
#1 gets her wings
#2 gets his wings
#3 - littlest doesn't



og does nice things



Littlest gets biggest wings



Be + 2 Cl atoms

sp^2 hybrids form when you move an s^1 . Taking a nice pair & moving an e^- up. Can you make a new orbital lower in energy?

Separation Δ s of 109.5°

VSER concept of trigonal pyramidal geometry. Hybrid orbital


O atoms bond w/ 2 H's. sp^3 hybrids, 2 empty e^- , 2 lone pairs. Tetrahedron, 2 lone pairs

Laughing Gas Elberhard's rxn.

N_2O - nitrous oxide, laughing gas

CS_2 - Carbon disulfide. Bonding similar to CO_2

Exothermic, obviously. Huge explosion.



Break old bonds (collision velocity, etc) & make new bonds.

Electrons moving so much faster than nuclei. Nitrogen molecules bang into us at like 1050 mph. e^- move around trying to find place where they get the lowest energy.

e^- end up in locations that match hybrid orbitals.

Reactive collision between molecules to sample new bonds
exothermic rxn encourage more bonds

orbitals are
probability density plots that show outer boundary, Computed
from wave functions.

wave functions used to compute probability density - hybrid orbitals
result.

5 e^- pairs in sp^3d hybrids

□ □ □ □ □ $3d$


□ □ □ $3p$

□
 $3s$

Hybrid bonds, Orbitals

Wednesday, January 10, 2007
9:04 AM


sp^3d^2 hybrids




sp (b d) Linear

sp^2  Trigonal planar almost asphere

sp^3 

sp^3d  Trigonal bipyramidal 

sp^3d^2  Octahedral 

Quiz

Lewis dot structures

Formal charge

VSEPR models

Bond dipoles + electronegativity trend

Resonance

Secondary

Bond length trends

Bond energies & enthalpies of rxn

Eberhard's lab demonstrations

All Ch. 10 possible


Eberhard's demo

Oxidation w/ built-in oxidants


Nitro-glycerine capsules (explode & save lives w/ heart attacks)



Nitro cellulose is like cotton



Black powder: potassium nitrate 75, charcoal 14, sulfur 11



Low velocity explosion

The spark in the hydrogen balloon didn't make the balloon blow up. Explosion occurred from outside in. Hydrogen needs oxygen to explode.

Hybrid orbitals for single, double, triple bonds.


- Overlap of hybrid orbital w/ bonded atoms gives an e^- pair w/ e^- density or bond axis. Called a σ bond (sigma)

- Single bonds are always σ bonds

- Single bonds are cylindrical & can rotate

- Double bonds have σ bond & π bond.


Ethylene



Need 3 orbitals



VSEPR doesn't explain double bond. Double bond = 1 repulsive group, so trigonal planar. Hybrid that gives trigonal planar might minimize repulsion



But we have a leftover p orbital

Lower energy region of space to hang out w/out bumping into other σ bonds


into other e^- pairs.
Hanging out is hard to do...

π bond formation in progress. p orbitals haven't overlapped yet.




This is a new way to use VSEPR. Select hybrid orbitals
CNOH repeat their bonding patterns

Overlap of 2 p orbitals is a valence bond concept that does explain π bonds.




Carbon is sp^2 for σ bond, not sp^3
If you had sp^3 , it'd be sticking out at 109.5°

Double & triple bonds easier to understand w/ orbital overlap & σ framework



$H-C \equiv C-H$ Linear
Hybridize to make linear sp



Molecular Orbitals

π bond model is a local (2 atom) model

Magnetic properties need explaining
Semi-empirical methods show π as localized over more

Spectroscopic methods show e^- can delocalize over many atoms w/ alternating double bonds (organic molecules w/ C chains)

Quantum mechanics can be done by allowing e^- to move over whole molecule

Molecular Orbitals, Energy

Tuesday, January 16, 2007
8:55 AM


Molecular Orbitals

Core $1s$ e^- too stable to share


MOs are made from math combos of wavefunctions of similar energy & diff atoms. If MOs = # of atomic orbitals combined.

MO energy order depends on atomic orbital energies

$1s$ orbitals of 2 widely separated H atoms



Create lower energy bonding MO & higher energy antibonding MO






How to fill

Distribute valence e^- for whole molecule, lowest energy 1st.


Each MO holds 2 e^- , each w/ ↑ the each w/ diff spin properties

Hand: 1st put 1 e^- in each orbital of equal energy (same spin property) before adding 2nd to make a pair.

MOs for 1-3 e^- H_2^+ H_2 H_2^- He_2^+



3-4 e^-





Stable species have more e^- in bonding order than antibonding

$$\text{Bond order} = \frac{\# e^- \text{ in bonding MOs} - \# e^- \text{ in antibonding MOs}}{2}$$

$$\frac{3-1}{2} = 1$$

$$\frac{2-1}{2} = \frac{1}{2} \text{ bond}$$


MOs can explain weaker + stronger bonds more effectively

$$\text{BO}_{\text{H}_2^+} = (1-0)/2 = \frac{1}{2} \text{ bond}$$

$$\text{BO}_{\text{H}_2} = (2-0)/2 = 1$$


$$\text{BO}_{\text{He}_2^+} = (2-1)/2 = \frac{1}{2}$$

$$\text{BO}_{\text{He}_2} = (2-2)/2 = 0$$




is too small to overlap,
→ not make MOs

Isoelectronic means same #
of e^-




B_2 has 6 valence e^- , 2 in $2s$ + 4 in p
 B_{2p} is when p overlap

xz & yz axis p orbitals overlap to make π bonds
 π bonds are MOs, & have same energy.



Mr. Sp. Japs
What is he
talking about?



Eberhard's Demo



O_2 gets stuck
No dissociation

I AGREE
No IDEA

Quiz Review, Bonding

Wednesday, January 17, 2007

8:56 AM

Quiz

Bond enthalpies, using heat of formation for rxns
Hybrid orbitals & VSEPR

Pi & Sigma bonds

Molecular orbitals of diatomic molecules (making MOs, bond order, magnetism)

Delocalized Lewis structures & resonance, simplified MO description.

end of delocalization in benzene
Eberhard's lab demonstration

2 possible diagrams.

$12 + e^-$, no matter molecule, σ lower (see chart online)
 $11 - e^-$, π



Paramagnetic is $\uparrow \uparrow$

Diamagnetic is $\uparrow\downarrow$

Smaller charge, smaller nuclei, more orbitals collide. Bigger don't collide as easily.

Bonding in diatomics

- Identify bonds in MO models
- Fundamental quantity is e^- density
- Bond order = # of bonds = $(n_b - n_a)/2$



Paramagnetism from unpaired spins

Diamagnetism from paired spins


Ferromagnetism from alignment of many atoms to make magnet

Electronegativity distorts shapes of molecular orbitals

Some rules for bonding work for heteronuclear diatomics & make molecules

Delocalisation occurs when resonance structures can be drawn
If an MO is spread out over 2+ atoms, not like C_2H_4 (isolated double bond).

Benzene is a good example, & alternating double bonds in a chain.



bonds shift but # of each type don't change.

~~Ozone~~

Delocalization could be non-bonding & pi type

π Antibonding



π Nonbonding



π Bonding



1 pi bond shared in 2 bonds
is essentially $\frac{1}{2}$ bond order contribution
so total is 1.5 bond order between
each pair of Os.

Now nonbonding e^- could have
unpaired spins

NO_3^-

4 p orbitals

delocalized bonding/antibonding

Degenerate non-bonding orbitals


HW 38, 39, 40

Simple Lewis Dot Structures, look at resonance structures.
Delocalized bonds


Can electron charge density plots help? Yes, problem is to discuss unusual bonding in terms of concepts that work for majority of cases.

If you take e^- density as $e^-/\text{unit volume}$, has 1-1 relationship w/ ground state wave function.

~~wave~~ e^- density is the bond.



For HF, it has all the motion. F barely moves.
In CO_2 , move in many ways




Formaldehyde looks like it's
when it moves



Another looks like Medieval torture



Solid state structures





refraction graphing
red refracts more

X-ray diffraction can be used to calculate solid state structures

You can take one of the crystals we made, put it under the X-ray (short wavelength) diffractometer & from refraction pattern calculate the structure.

Body
Centered
Cubic




Unit cell
has 2
atoms

$\frac{1}{8}$ corner atoms in corners

$$\frac{1}{8}(8 \text{ corner atoms}) + 1(\text{central}) = 2$$

Face
Centered
Cubic



no central
atom

$$6\left(\frac{1}{2}\right) + 1 = 4 \text{ atoms in face centered}$$

Polymorphism & allotropy

Many-shape



a material that
can change
its shape


existence of more than 1 crystal structure of a given atom/compound (Carbon [graphite & diamond])

Iron is polymorphous & allotropic

Any regular solid will pack into a repeating structure. Some of the structures are more efficient than others.



Calculating Packing efficiency of simple cubic



$$\frac{1}{8}(8) = 1$$

1.00
100%



~~OK~~
Condense so
they touch


Volume, if diameter = 1
 $\frac{4}{3} \pi \left(\frac{1}{2}\right)^3$
 $\frac{1}{6} \pi$

Volume of unit cell = 1

Packing efficiency $\left(\frac{\frac{1}{6} \pi}{1}\right) 100 = \%$


Unit cell is smallest repeating unit.

If atoms weren't touching, distance
would not be 1.



$\left(\frac{2}{\sqrt{2}}\right)^3 = \text{volume of cell}$


Body centered cubic



Calculate
that



VERTICAL FLIP



Prelab - do warm-up exercises in lab manual

Title + objective

No procedure (bring lab manual)

In end, write brief conclusion

Interaction Forces

Monday, January 22, 2007
8:58 AM

What are the interaction forces between molecules
Liquids: $10^{-9} - 10^{-12}$ s, slow changes

internal vibrations fast on this timescale

Solids: molecules don't translate or rotate, vibrations between molecules

Also fast internal vibrations

Gases: Distances very long range. Need higher P to see non ideal gas behavior

Evidence for forces? Heat of phase transitions

Vaporization is an example

1. Dipole-dipole forces, permanent dipole moments
2. Dispersion / London (transient dipoles)
3. Hydrogen Bonding

Dipole-dipole

- transient alignments, avg. net attraction. Things move.
- Larger = better interaction
- ~~less~~ very unlikely to have as much order as pics in book
- Fluctuating order makes avg energy less than expected
- Contributions about 5-20 kJ/mol. Part of H-bond forces.

Dispersion / London Forces

Fast fluctuating. can make instantaneous dipole (lasts 10^{-15} sec)

A neighbor molecule can respond w/ its own instantaneous dipole that is induced by 1st dipole. While thing is short lived BUT on avg there are forces created

Polarize neighboring molecules

Molecular properties enhance this

- Polarizability, distortability of electrons/volume. Larger atoms have more e^- & are more polarizable. Bigger w/ "mass" which ~~less~~ poor substitute for H^+ .
- Larger polarizability will have greater induced dipole
- Distance dependence: $F \propto 1/(\text{distance})^6$ so closer is better

Implies that ^{up} close alignment is good ^{more}
Long molecules in close contact have London dispersion forces
Range of energies is 0.1-20 kJ/mol

More electrons = higher boiling point (more polarizability)

Hydrogen bonds are dipoles that interact especially strongly,
Occur when H atom attached to O, N, F atoms. $F-H \cdots O-H \cdots N-H$
Interaction is between highly positive H & a lone pair. Network
of H_2O is b/c there are lone pairs & 2 O-H, 4 H bonds.
Mixture of molecules also H bond. 10-40 kJ/mol

Surface Tension - result of intermolecular forces

Exterior molecules ^{have more} symmetric forces, surface molecules have more force
pulling them to the bulk. This adds a new energy term for surface
molecules, J/m². Energy needed to increase area by 1 m²

Merise us: Balance between liquid cohesive by surface tension & surface
adhesive forces. Hg is diff

Vapor pressure T variation & pressure for equilib closed containers

Eberhard's demo - water + oil went into flasks.

Diff slopes for $\ln P$ vs $\frac{1}{T} \times 10^3$
graphs.

Diff slope means

$$\ln \frac{P_2}{P_1} = \left(\frac{\Delta H_{vap}}{R} \right) \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$


Prior graphs show slope of water larger than non-polar
toluene. (see book)



P vs T curves

Tuesday, January 23, 2007
8:53 AM

Apply fixed Joules/sec & measure T changes.
Phase changes




Slope $\propto \frac{1}{\text{heat capacity}}$

Liquids have biggest heat capacity, then solid, & a little smaller is vapor.

I Paid good money for that
↓

Mercurius!
Rocks!

Supercritical fluid is at same density as a liquid.



Vapor drives back atmosphere (boiling) Open vessel

Triple point is lowest temp for a liquid, & point where all 3 are in equilib.

X = FILES

Takes a high P to have a liquid or a triple point and it Sublimes at 1 atm

Supercritical fluid has no meniscus

At the critical point, you go from region w/ meniscus to no

At the critical point, you go from region w/ meniscus to no meniscus. The liquid is reorganizing to make a thing w/out meniscus. Chaos, light scattering over domains.

Water phase diagram



Triple pt.
4.5 Torr

I want to
Believe

↑
incomplete.
unacceptable,
despicable,
disgusting
sorry excuse for
a diagram.
There's no ice IV!

YO
YO
MAMA!

Ice has neg. slope. ΔV is neg.

$$\frac{dp}{dT} = \frac{\Delta H}{T \Delta V}$$

Most have + slopes.

$$\ln \frac{P_1}{P_2} = \left(\frac{\Delta H_{\text{vap}}}{R} \right) \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

No, yours!

Sublimation

Triple point of H_2O is 4.58 Torr, 0.01°C

$$\Delta H_{\text{subl}} = \Delta H_{\text{vap}} + \Delta H_{\text{fus}}$$

Different shape for solid-gas line
than solid-liquid line.

Types of Solids

Ionic $NaCl$, KCl

Lattice energy from Coulomb interactions

Metallic (Al , Cu , etc)

Molecular (H_2O , London dispersion, etc)

I WANT TO
GET AWAY



WHO
BELIEVES?

Network Solids (quartz, sand, diamond)


Amorphous Solids (glass, polymers)

Crystal lattice

repeating position of atoms (pattern)

Unit cell is what we use to make a lattice

X-rays have short enough wavelengths to allow spacing between atoms to act like a plane



Unit cells

Wednesday, January 24, 2007
9:03 AM

X-ray diffraction of crystals



Quiz Topics


- Intermolecular forces (types, properties, effects)
- Heats of Phase Transitions & Heat Capacities (applications)
- Vapor Pressure of Liquids & B.P.'s
- Phase diagrams
- Crystal Unit cells (sc BCC, FCC)
 - Lattice energies, NaCl CsCl, hexagonal close packing not on quiz
- Use KGS Table to organize unit cell info
- Clausius-Clayron Eqn on exam (don't need to memorize)

Unit cells



Simple cubic
net count 1
atoms/unit cell

4 touching in
same plane &
nearest neighbors



Body centered
net count 2

4 above body &
4 below, 8 nearest
neighbors.

Face-centered
net count 4

Simple cubic - Edge length = $e = 2 \times \text{atom radius} = 2R$
Volume of SC unit cell = $e^3 = 8R^3$
1 atom/unit cell
1 atom has $V = \frac{4}{3}\pi R^3$
Ratio $\frac{\text{atom } V}{\text{unit cell } V} = \frac{\pi}{6} = .5236$
Only 52.4% of lattice volume is atoms

Simple cubic has most empty space.

Body centred - $(2)^{\frac{1}{2}}e$ is diagonal of a face $(\sqrt{2})$
 Sum of squares is: $2e^2 + e^2 = \sqrt{3}e^2$ is
 body diagonal. body diagonal = $(3)^{\frac{1}{2}}e$
 Body diagonal = $4R$
 $= 3^{\frac{1}{2}}e$

$$\text{Volume unit cell} = e^3 = 64 R^3 / (3)^{\frac{3}{2}}$$

$$\text{Volume of 2 atoms in unit cell} = 2 \left(\frac{4}{3} \pi R^3 \right)$$

Ratio of volume atoms / volume unit cell = .680 or
 68% of lattice occupied by atoms

Face centred - $2^{\frac{1}{2}}e$ is diagonal of a face
 $4R = 2^{\frac{1}{2}}e$

$$\text{Volume 4 atoms in unit cell} = 4 \left(\frac{4}{3} \pi R^3 \right)$$

$$\text{Volume unit cell} = e^3$$

Ratio volume atom / unit cell volume = .740, 74%
 of lattice filled w/ atoms.

Highest fraction of space that can be filled
 by packing same size spheres.


Close Packed Structures

Cubic close packed case means face centered cubic

Coordination #s are 12

Hexagonal close packed case has no cubic unit cell
 both have smallest free volume of 26%

NaCl Lattice



octahedral





octahedral

Everything is surrounded
 by 6 atoms.

Ends up being cubic

4 formula units in unit cell

$$(2R_{Na}) + (2R_{Cl}) = 4R \text{ edge}$$



~~Face~~ Body centered cubic

NaCl requires 4 formula units ^{to describe} in the unit cell

CsCl has 1 formula unit, Cs in center. Cs, Cl touch on body diagonal, which has $2R_{Cs} + 2R_{Cl}$ edge = $\frac{\text{Body diagonal}}{\sqrt{3}}$

NaCl has FCC unit cell, density = 6.84 g/cm^3 . Calculate radius.

We have density.

Unit cell volume = $(\text{edge})^3$. We need to relate edge to R
 $4R$ is distance in units of atom radii. e = edge length.
 Always work w/ touching atoms.

$$e = \frac{4R}{\sqrt{2}}$$

Unit cell density = microscopic density. Need # atoms in unit cell, mass of atoms.

$$\text{Unit cell volume} = e^3 \quad \text{density} = 6.84 \text{ (mass)} e^3$$

$$\text{Mass} = 4 (58.69 \text{ g/Na}) = 3.898 \times 10^{-22} \text{ g}$$

Find e , solve for R.

Bar of X. Density = 10.5 g/cm^3 . FCC unit cell as $4.09 \times 10^{-10} \text{ m}$
 Atomic weight?

Titanium is BCC Density = 4.5 g/cm^3 . Edge length $\propto R$?


Experiment 11 (lab manual allowed) Optical Diffraction

OCD are Team 1!

We're using Visible Light in the lab

Bragg diffraction occurs when x-rays interact w/ crystals.

Fraunhofer diffraction occurs when light interacts w/ the slides



Crystals are constructed of atoms or molecules arranged in a reg. pattern.

<http://www.matter.org.uk/diffraction/introduction/default.htm>

Red light = 670 nm

Green light = higher energy, shorter wavelength

RIDE, Cap'tain
Ride upon your
mystery ship!

Bring a ruler, bring lab manual

WE ALL LIVE IN
A YELLOW
SUBMARINE

Diffraction caused ~~by~~ when you have a repeating, periodic array & the repeat distance is similar to wavelength of waves.

This is on the $E-10$ angstrom scale

Use X-rays


We use visible light

CH
I
DO NOT
F
A
RED
DYE!

Result

- Constructive or destructive interference

Fraunhofer Diffraction Exp.




RIDE Captain
Ride upon your
mystery ship!

$$\tan \phi \approx \sin \phi \approx \phi \quad \text{b/c infinite distance}$$

$$\tan \phi = \frac{x}{L}$$

Δ is distance between dots

Constructive interference $d \sin \phi = n \lambda = 670 \text{ nm}$




USS
ENTERPRISE
CAPTAIN
JEAN-LUC
PICARD

NX-01
ARCHER
NCC-1701
KIRK
NCC-1701-A
KIRK
NCC-1701-B
HARRISON
NCC-1701-C
GARRETT
NCC-1701-D
PICARD
NCC-1701-E
PICARD

Spontaneous change of natural phenomenon
hot $\rightarrow$ cold

Decrease entropy, put stuff together.
Phase transitions (solids melt to liquids, etc)
Dissolving (making soln)
Freezing points



Entropy, S , state function, ≥ 0 w/ units $J K^{-1} mol^{-1}$, measure of molecular randomness.

Entropy of gases $\gg$ liquids $>$ solids

Entropy of a gas increases with increased volume, increased temp (for everything solid, liquid, gas) $\circ$ at OK, perfect crystal.

Entropy of ideal mixing gases or liquids always is positive.

$$\Delta S = -R$$

$$S_A + S_B + \Delta S_{mix} = S_{tot}$$

Second Law of Thermodynamics: Entropy of universe either stays the same (equilib) or increases w/ any spontaneous changes.

We have no heat flow in forming an ideal soln, so the increase of entropy is for the universe. Etc is a spontaneous process.

Starting test tubes




are not in equilib

pure water salt + water

Spontaneous process is B has biggest entropy. Solvent moves through vapor to region w/ higher entropy.
A is empty after vacation.

Equilib point of view is Tube A + B should have same vapor pressure. But pure solvent always have greater vapor pressure than impure, and so stuff will move to soln w/ less vapor pressure.



20 mol H_2O ,
2 mol salt

20 mol H_2O ,
1 mol salt

Less salt has biggest vapor pressure.

Raoult's Law $\rightarrow$ If xlt mole fraction is not too diff then new equilb is possible by equating vapor pressure above each tube.

$$P_{soln} = X_{solvent} P^0_{solvent}$$

$$X_{solvent} + X_{junk} = 1$$

One tube gained 6.86 mol, other loses. Agains. Bloxes.

Entropy of gas has value that is proportional to partial pressure. At equilb soln entropies in A + B are equal. Gas entropy also equal.

$\approx$

Entropy also -

Gases have biggest entropy. S depends on P or V .

BP elevation, FP depression w/ entropy idea.

How can you create a vapor pressure of 1 atm above a soln? At original T of BP a soln has less v_p due to Raoult's law. Increase T find a place where $v_p = 1 \text{ atm}$.

$$\Delta T = k_b m_j \quad \text{mole fraction of solute}$$

FP Depression: New equilibrium between pure solid & junky soln. Solid is pure bc for most things crystal lattice excludes junky. Original FP is an equilibrium w/ pure solid & pure solvent. We need to lower entropy of junky by lowering T .

$$\Delta T = -k_f m_j \quad T_{\text{final}} - T_{\text{initial}} = \Delta T$$



D2

"QUACK, QUACK, QUACK"

BP elevation Create vapor pressure of 1 atm above a soln - At original T of BP a soln has $< v_p$ due to Raoult's law. Increase T & find a place where $v_p = 1 \text{ atm}$.

$$\Delta T = k_b m_j$$

FP depression new equilib is between pure solid & Jungsue soln. Solid is pure bc for most things crystals exclude Jungsue. original IFP is an equilib w/ pure solid & solvent.

$$\Delta T = -k_f m_j$$

Jungsue = solute

We don't care what Jungsue properties are, just molar of Jungsue

Ionic salts can form multiple ions

At greater dilution there are less interactions between oppositely charged ions so they can act like completely separate entities.

If a diver surfaces too quickly, N_2 dissolves in diver's blood will release quickly resulting in painful & dangerous condition called the bends. This is same as taking off cap on pop bottle.

Things have higher b.p. when they break up into ions bc concentration goes per ion.




Osmotic $>$ FP $>$ BP

for precision of determining unknown molecular weight

Osmotic Pressure effect

Membrane allows solvent to move, but not solute

Jungsue



Pressure changes the entropy of the solid. Pressure on salt side, salt has most entropy, and only takes very small diff in concentration to cause an effect.

Entropy is measurement of how well molecules can occupy diff places & be totally random.

Pressure must be lower than entropy of soln,

Volume is very dependent on pressure. $V \downarrow$, entropy reduces due to fewer locations for solvent (less random) or fewer microstates in statistical model. Higher P , $\downarrow$ entropy.

Excess pressure, π

$$\pi = CRT, \quad c = \text{solute molality}$$

c becomes ic if i is # ions.

Osmotic Pressure


Hypotonic $< .92\% \text{ m/V}$ rupture

Isotonic Saline $.92\% \text{ m/V}$


Even if solns are not very diff concentrations one can cause enough internal pressure because damage or to dehydrate cells.

Hypertonic $> .92\% \text{ m/V}$ crenation

Dialysis designed to let many things pass.



Reverse osmosis - Desalination



Colloids, Distillation

Wednesday, January 31, 2007

9:01 AM

Lewis Dot structures


Formal charge

VS PER Models

Molecular dipoles & electronegativity

Resonance (identify cases)

Bond length trends



lipstick on a pig

Bond energies, enthalpies of rxn, heats of formation.

Hybrid orbitals / VS PER relationship

Pi Sigma bonds

MO of diatomic molecules (making MOs, use to describe molecules, bond order, magnetism)

Delocalized Lewis structures & resonance, simplified MO description of MO

Intermolecular Forces (types, properties, effects)

Heats of phase transitions & heat capacities

Vapor pressure of Liquids & BB

Phase Diagrams

Crystal unit cells

NaCl, BCC unit cells

Entropy Supplement

Vapor pressure of solns, Raoult's Law

Colligative properties, FP depression, BP elevation, osmotic pressure

Henry's Law, forces in liquids, molality

Vapor composition, BP-composition diagrams, Fractional distillation, azeotropes & non-ideal solns, nBP

$$n\lambda = 2d \sin \theta$$

$$\pi = iMRT \quad \Delta T_f = -iK_f m_j \quad \Delta T_b = iK_b m_j$$


$$P_s = X_s P_s^\circ \text{ non volatile Solute (solvent)}$$

$$P_i = X_i P_i^\circ$$

Raoult's Law gives up Benzene: $5 \times 95.1 = 47.6 \text{ mm Hg}$
 $5 \times 28.4 = 14.2 \text{ mm Hg}$


So we know partial pressures Total VP Add together.

Fractional distillation: Compare BP of liquid vs vapor composition as another curve.



Start at blue arrow.
Red lines link vapor comp that is now richer in benzene.



Condense liquid in stage 1, collect vapor in stage 2 that is even richer in benzene, keep repeating & final material is pure benzene.



Lower BP has highest vapor pressure & so vapor is richer in lower BP

Intermolecular forces can be weaker than individual species or stronger. This CAN, but not always, generate azeotrope. Azeotrope is soln at particular composition where vapor & liquid have same composition (depends on external pressure). At that special comp., liquid is boiling at pressure T , which is exactly what a single pure liquid would do. Higher total VP than expected by Raoult's law. BP of azeotropes is lower b/c of higher VP.
Mixture more liquid.

chemicals pur = liquid



96% ethanol azeotrope

BUREAU
MULDER!

I ♥ U,
Kerisha!

HATE
M
JUST KIDDIN'

FBI

Colloids

Nanoparticles of gold in soln. Size controls the color. So light scattering depends on size.

Shorter wavelengths = more intense scattering

Daddy, why is the sky blue?

Well... to match your pretty eyes

Nope, not even close. See, all colors have wavelengths that are diffused by nitrogen & oxygen. Since blue is the shortest wavelength, it's diffused up to 10 times more.


Our eyes are more sensitive to blue than purple.

Interaction Forces


Thursday, February 01, 2007

9:01 AM

London Dispersion Forces




London dispersion forces based on polarizability of molecule.
Polarizability related to # electrons per mol.
Also, shapes of molecules affect it.



Dipole-Dipole

Permanent dipole. Always has δ^+ & δ^- . Water is dipole dipole.



2 molecules w/ always dipoles interact.
These are permanent, so stronger than LDF

Thiophene f.p. -38.3 b.p. 84.4°C used as solvent in place of benzene. Combustion of 2.348g sample produces 4.813g CO_2 , 1.005g H_2O & 1.788g SO_2 . When a $.867\text{g}$ sample of thiophene is dissolved in 44.56g benzene (C_6H_6) the f_p is lowered by 1.183°C .
Molecular formula of thiophene?

$$\Delta T_f = -k_{fp} m_f \rightarrow \frac{\text{mol solute}}{\text{kg solvent}}$$

$$m = \frac{\Delta T_f}{-k_{fp}} = \frac{-1.183}{-5.12} \approx 2.2 \text{ mol solute (Solute) / kg solvent (benzene)}$$

$$\frac{.2 \text{ mol}}{\text{kg benzene}} = \frac{x \text{ mol}}{44.56 \text{ g benzene}}$$

$$.0103 \text{ mol} = x \text{ mol}$$

thiophene

$$\frac{.867 \text{ g thiophene}}{.0103 \text{ mol thiophene}} = 84.2 \text{ g/mol}$$



$$\text{CO}_2: 4.913 \text{ g}$$

$$\text{H}_2\text{O}: 1.005 \text{ g}$$

$$\text{SO}_2: 1.788 \text{ g}$$


$$\text{thiophene}: 2.348 \text{ g}$$

$$4.913 \text{ g CO}_2 \left(\frac{1 \text{ mol CO}_2}{44 \text{ g CO}_2} \right) = .111 \text{ mol CO}_2$$


$$1.005 \text{ g H}_2\text{O} \left(\frac{1 \text{ mol H}_2\text{O}}{18 \text{ g H}_2\text{O}} \right) = .0558 \text{ mol H}_2\text{O}$$

$$1.788 \text{ g SO}_2 \left(\frac{1 \text{ mol SO}_2}{60 \text{ g SO}_2} \right) = .0298 \text{ mol SO}_2$$

$$2.348 \text{ g} \left(\frac{1 \text{ mol}}{84.2 \text{ g}} \right) = .0279 \text{ mol thio}$$



At 25°C , 0.5 g sample polyisobutylene in 100 mL benzene
 soln has osmotic pressure that supports 5.1 mm column of
 (d = .48 g/mL). Molar mass of polyisobutylene? For Hg, d = 13.6 g/mL



$$\pi = MRT$$

$$\pi = \frac{n}{V} RT$$

$$V = hA$$

$$V = 5.1 A$$

$$d = .48 \text{ g/cm}^3$$

$$m = Vd$$

$$m = (.51 \text{ cm})(.88 \text{ g/cm}^3) A \text{ cm}^2$$

$$m = X$$

we'll pretend this is weight of mercury

$$13.6 \text{ g/cm}^3 = X/V$$

$$A \cdot X \text{ g Hg} \left(\frac{\text{cm}^3}{13.6 \text{ g}} \right) \left(\frac{1}{A \text{ cm}^2} \right) = h(\text{Hg}) \text{ cm} \rightarrow \text{mmHg}$$

convert to
torr, = pressure


$$\pi = MRT$$

$$\text{mmHg} \left(\frac{1 \text{ torr}}{1 \text{ mmHg}} \right) = \frac{MRT}{M(R)(.5 \text{ g})}$$

$$= \frac{n}{V}$$

SO₃

$$6 + 3(6) + 2 = 26$$



$$26 - 6 = 20$$

$$20 - 12 = 8$$

$$8 - 4 = 4$$

$$4 - 4 = 0$$



3 resonance structures

$$FC = \# \text{valence} - LP - \frac{\# \text{bond } e^-}{2}$$



Prelab

Title

objective

Procedural outline

Chemical eqns

odd exercises p-64

Write eqns w/ states

Rxns

I. A1-A3


II. A1, B1

III. B1-B2, C1-C3

IV. A-D


V. A+B

We won't do red phosphorus rxn,
the sodium, or some other thing
We still fill out observations
(from here)

Combination rxns.

B. Nonmetal + Nonmetal.

Small quantity of red phosphorus in test tube heat gently
w/ Bunsen burner

RxnObservations


white smoke

White solid forms, ^{white} smoke comes out
Don't just heat bottom of tube (b/c you don't want it just concentrate in one place)

Flame inside tube, orange solid Bubbling w/ liquid & flame.


Oxidized/Reduced?

Add 5 mL ^{H₂O} to the residue, shake to dissolve, test w/ pH paper.



So P oxidized, Oxygen reduced. From 0 $\rightarrow$ -2. And 0 $\rightarrow$ +5

Rxn




Observations

Turns chocolate color, then reddish thing. Some suspended solid. P is probably in there (unreacted phosphorus) but $P_4 O_{10}$ is dissolving.

Displacement of a halide ion from a salt

Add 10 drops $Br_{(aq)}$ + 5 mL $KI_{(aq)}$ & shake. Add 3 mL hexane & shake again

Rxn



Observations

Clear soln, add Br. Br makes brown color soln. Both aqueous

Add hexane to make soln nonpolar (hexane nonpolar) & Br is nonpolar, polar KI

Hexane layer on top b/c less dense than water.

Purple color on top layer, and that means something has reacted. Br is brown. Bottom layer ~~is~~ yellow


Iodine I_2 & KBr

Bromine reduced to Bromide.

Iodine oxidized to Iodide.

Add 10 drops $I_{2(aq)}$ + 5 mL $KBr_{(aq)}$ + shake. Add

3 mL hexane & shake again.







No rxn

If bromine strong enough to oxidize iodide then ~~the~~ iodide cannot ~~not~~ oxidize bromine.

Observations

Hexane layer would be clearish purple.

Exercises



Superconductors 991-994
Colloids/Nanocrystals 557-559, 992

Self-Assembled Materials > ~~Handouts~~
 Nanomaterials


Silicon	452-456	Ch 11:45-47, 49-51
Silicates/Silicone	901-906	
LEDs	908-909	
Liquid Crystals/Polymers	1117-1121	Ch 21 45, 48 Ch 26 a1

Diff between intrinsic & extrinsic semiconductor is intrinsic is not doped, extrinsic is.

Metals have ~~a small~~^{no} gap between valence & conduction bands
 Semiconductors have a small band gap
 Insulators have a large band gap.

Network Solids

Crystal is a structure where it is one big molecule. For instance, diamond. A network of carbon




One huge network of carbon.
 Carbon is tetrahedral.

Allotrope - 3 classes of allotropes of carbon. Diamond, graphite, & things called fullerenes

Crystalline forms of a material


Graphite has a sheet-like structure.



Weak interactions between the sheets.

So graphite has diff types of bonds. Lower melting pt, easily broken. Alternating double bonds means it can...

absorb light, and so it's darker.




Soccer ball pattern for fillings

Very small ants are very highly colored. Alternating double bonds gives a very deep purple. All carbons are sp^2 hybridized instead of sp^3 hybridized

SiO_2 vs CO_2
sand, glass, quartz
($O=C=O$)

(structure is diff for these (not molecular solids))



One molecule stays in molecular form, other forms a network solid.

You have to break double bonds & form single bonds to make CO_2 a network solid, & they don't want to do that.

SiO_2 double bonds aren't natural. They're weaker than SiO_2 single bonds.

SiO_2 high melting pt
Glass, quartz, doped structures

B_2O_3 K_2O
labware (lenses)





Si Similar structure to diamond
Semiconductors

We determine if something is a semiconductor by its properties
Same for metals & insulators.

Metals (structures that have atoms that share e^-)

As you add elnts to the structure, you get more molecular


orbitals. Ones in bonding & antibonding overlap. These form a band. The bands formed are valence band or conduction band



For a material to conduct, we should be able to move e^- freely throughout the bands.

If you heat the system, the semiconductor conductivity will increase. Metals become worse conductors w/ temp. It decreases w/ a metal b/c it vibrates atoms, more collisions w/ e^- & nuclei, & impedance in free e^- .

Doping is the adding of elements. Take a material & add a small amt of another elmt. Like adding P to Si. Extra valence electrons & more will jump. Create levels:





Ga instead of P, has less e^- . Still becomes a better conductor. P-dope. Changes whether you have full shells. Take away full valence shell or bonding square, allow more to jump.

Types of Conductors, Insulators

Tuesday, February 06, 2007
9:01 AM



Eberhard's demo [↑] + how you make silicon.




Semiconductors are better conductors when heated.
Metals are better conductors at low temps.

Light - insulators have large band gap & don't absorb visible light. Semiconductors absorb some, & metals are dark b/c they have no band gap.

P is doped w/ an element to the left

N is doped w/ an element to the right

PN junction, PN diode (2 types of semiconductors brought close to one another makes current naturally flow)



Monomers coupled w/ each other make polymers


Superconductors - Mainly ceramic materials. Certain materials were discovered in the 80s that could be superconductors at high temps.

$YBa_2Cu_3O_7$ is a superconductor
↓
will conduct w/ 0 resistance.


Polymers/Liquid crystals

Oligomer is something smaller than a polymer but bigger than a monomer.


Monosilane



Disilane



Trisilane





If you keep adding, you get a polymer.

Organosilanes





↑
organic groups attached to Si

These types of structures react with water



That creates a situation where it is can happen!



← # of these guys. And N can be anything from 1 → 100 or something.

Silicone acts as oils that doesn't degrade under high temps. So it's awesome for cars.

Liquid crystals are not polymers. They're diff. Made from macromolecules

Molecules w/in liquid are not completely disordered. Align w/ themselves. Build things.

Rod-shaped molecules do this.


Ordering in Liquid-Crystal phases

1) Normal liquid

Random




2) Nematic liquid crystals



Oriented but not in planes.

3) Smectic A liquid crystals



Oriented in planes

4) Smectic C liquid crystals



Tilted & in planes & oriented

LCD display is a liquid crystal display.

Liquid crystals can also be thermometers. Change color when you touch them w/ diff temp. Fingerprint analysis, laptop touch pad

laptop touch pad
They exhibit LC behavior at diff temps.

Kevlar is made up of repeating chains (Amides NHCO group)
Molecules align a bit and that lets it stop projectiles
Chains can interact & it leads to strength

Polymers

Wednesday, February 07, 2007
9:03 AM

Hydrogen bonding between Kevlar

Addition polymers

Polypropylene 23 billion
lbs/yr
Polystyrene
Polyvinyl chloride

bottles, film, packaging, clothing,
kitchenware

> All come from double bonds,
go to single bonds.


Condensation polymers

2 types of molecules come together

Polyurethane

Polyethylene terephthalate (polyester)

Nylon




Rubber is another important polymer.

There are naturally occurring polymers.

DNA, rubber, latex, proteins, cotton, skin

Rubber:



At the ends, sometimes polymers grab whatever they can find. The ends don't matter to the whole polymer's properties. This is why grabbing whatever & not completing the octet.

Polyisoprene is the above. You have to add vulcanization (add sulfur) to it to make it cross-link. This changes properties of the polymer. This is how they make tires.

TC

If you covalently attach the polymer w/ the sulfur, it becomes stronger.

DNA is made up of sugar & phosphate. Bases hang off the sugars, and are chemicals that are there for recognition purposes.

DNA strands recognize each other through H bonding.
H bond interaction through bases.

Electrons used b/c of their charge & short wavelength.

Electron microscope works like a CRT TV.

Gold Particles, Light Interactions

Friday, February 09, 2007
8:59 AM

$$\text{Nano} = 10^{-9}$$

edge of dime is 1 mm thick
Slicing edge 1000 times is a
micron. Take that & slice it 1000
times, nanometer.

Introduction, works cited due in prelab.

Gold colloids are solid gold but are in tiny pieces suspended
in soln.

In the medieval ages, people were making nanoparticles. They
didn't know this, but diff metals, concentrations, etc they made
colored glass.

On a macroscopic scale, things don't change color b/c they
change size. But at a nano scale, they do b/c it
changes size & shape & how light interacts.

Quantum Mechanics
Schrodinger Equation
Particle in a Box

Gold soln fed to algal/plant & plants reduce Ag^{3+}
ions to solid gold, $\text{Ag}(s)$.

Use type of spectroscopy to realize what the plants were doing.

Gold particles interact w/ light waves the way a boat
interacts w/ water.




When you change the size & shape, you perceive a diff color &
you can track those changes in the visible region of the
spectrum. $\approx 320 \text{ nm} \rightarrow 670 \text{ nm}$
high energy blue low energy red

If you see red, everything but red is being absorbed & there will be an absorbance peak in the green area.

Whatever light is absorbed you'll see the complimentary color w/ your eyes.

At purple range (absorbed) you'll see clear.
At red range (absorbed) you'll see green.

Hydrogen tetrachlorate trihydrate +
Sodium citrate



NanoHVB

At gold metal in

In solvent $\rightarrow$ H_2O , refractive index of $H_2O = 1.3$ is in

Size of particles (change size until you reach λ_{max})

Scattered light

Look for scattered light

Formal reports 7 pages maximum including graphs & pics
double spaced.

Graphs should be integrated into text, should not be larger
than $\frac{1}{2}$ page.

Past tense passive voice.

Experimental section should be brief. Any modifications made
to lab summary cited.

Due week of Feb 19th on Blackboard & TA.

prelab

Title, Objective, Procedure.

Intro & references submitted to TA

References

Book, ACS journal article, Lab manual, Other.

↓


NU Library

→ electronic resources

→ ACS American Chemical Society


use ACS style guide.

Take gold & shrink it down to 13 nm, changes color.
If you take any material & shrink it down, it forms a new material w/ new properties.




Nanotubes are cousins to fullerenes.

Gold nanoparticles form a colloid



The size is much smaller than the wavelength of light



On the interior, surface atoms see below atoms like themselves & atoms on top. Surface ~~is~~ diff from bulk

Many more bulk than surface materials. Majority of atoms in bulk. Shrinking down, ratio of surface-bulk goes up

All diff color silver. Ag is neat b/c if you take a soln of silver particles (yellow in color) you can see silver nanocrystals (colored) & spherical particles break down & become silver Δ's.

By making silver in diff sizes & shape you modulate diff optical properties. Photosynthesis of Silver NPs.

Composition, size, & shape matter.

100 nm	100 nm	50 nm	90 nm
Nanoprisms	Ag spheres	Ag spheres	Spheres

Tools

Electron microscope

Scanning Tunneling Microscopes

Atomic Force Microscopes

Scanning Probe Microscopes

1990's developed these
(IBM in Switzerland Binnig & Rohrer?)

Really sharp tip.

Really sharp tip feels atoms on surface (Metal & surface conductor pops electrons from atoms at tip to surface e^-)

You can get this thing for only \$10,000.

Limitation $\rightarrow$ has to be a conducting surface. We can't look at insulators

Atomic Force Microscope allows us to look at insulators. Bring tip down & measure repulsive interactions. Deflections from tip is monitored as $f(x)$ tip $\rightarrow$ surface

You can measure the stickiness of a surface too.

Get a topological map of the surface, & figure out if the atoms are hydrophilic or hydrophobic.

Tips are not as sharp in AFM tips.

If the chemical is reactive, you put chemicals on the tip to manipulate the reactive substance.



Printing tools (electronics)

Computation is a huge industry, making computer chips.

We can make smaller & smaller features.

We can make smaller & smaller features.

We can make a mol (6.022×10^{23} !) of transistors through photolithography.



Limit of how far you can go $\rightarrow$ limit of light wavelength
(smaller & smaller features,
cost $\uparrow$. Physical limits)

So they tried etching tips but you damage the tips.

Dip Pen $\neq$ Nanolithography.

Deposit materials on a surface. Use surface like a pen to move molecules.

Water collects at point of contact b/c of thermodynamic minimum to move molecules from tip to surface.

Single layer of molecules on a surface to write out stuff w/ them

We use the water droplet as a carrier solvent instead of the ink in your pen.

Richard Feynman's *Plenty of Room At The Bottom*
Predicts that nanotechnology will become a reality.

Photolithography is a serial process. We want to print structures in parallel. Take a single pen system & go to printer.

Top $\rightarrow$ down chisels bulk materials

Bottom $\rightarrow$ up breaks down material into almost elemental state & deforms.

You get a soln of particles in a colloid, repulsive charge stabilizes it.



We can determine that the particles do this by looking at them w/ an electron microscope or:

When you add salt to the colloid, particles grow together & eventually fall out of soln, form bulk gold.

Color change from red $\rightarrow$ blue. Changes wavelength of light, as it grows. Bigger particles scatter wavelength of light.

You can come up w/ detection systems based on this principle. DNA detection w/ nanoprobles. You can tack DNA on the surface of a gold particle. You can design things coded w/ anthrax, infectious diseases, etc when the DNA is recognized, the particles come together & change color.

DNA Litmus Test (Test is almost to the point of where you can detect stuff on the spot)



If you heat the new soln, it would turn back to red b/c it breaks apart.

We use e^- or sharp tips to etch and move nanoparticles around.

Dip Pen Nanolithography.

Water condenses at point of contact (from air)

Ink based or composition (molecules that react w/ surface)



Molecules crystallize on surface. Self-assembled monolayer.

Whole process = surface modification

Octadecanethiol (ODT) - properties of surface change - surface has a lot of organic properties (terminated by an organic compound, slippery nature). Change the tail end & make it a carboxylic acid. Something that looks more like water won't be hydrophobic like SH, but hydrophilic like OH.

Pip pen lithography lets you modify it on smallest scale

Hydrophilic materials are stickier. White correlates w/ hydrophilic
Blue correlates w/ hydrophobic

4000 yrs ago	120 years ago	60 yrs ago	20 yrs ago
Quill pen	Fountain pen	Ballpoint pen	Photolithography
"Quill"	"Fountain"	"Ballpoint"	engage/disengage
3 yrs ago	1.5 yrs ago	1 yr ago	tips based on
			heating
			6 mo. ago


You can print on the nanoscopic view (you can pattern anything you want)


One chip could screen for every disease out there can do same thing w/ proteins

You can etch the Enterprise !!!





on nano-scale!






Does the amt of NO_2 + N_2O_4 change as a function of external conditions?

As we heat it, it gets darker meaning more NO_2 forms.
If you cool it down, turns clear meaning more N_2O_4 .




Calcium wants to be Ca^{2+} . wants to be oxidized.

So $\text{CaO} \rightarrow \text{Ca} + \text{O}_2$ will favor calcium oxide at equilibrium. So!



Hydrogen & Oxygen will react violently to form water.




Activation energy has to start a reaction. High activation energy means it needs a real kick to get started. Some bonds are harder to break than others.

$\langle 0 = 0 \rangle$ is hard to break.


But at equilibrium, you're going to see a lot of product.

Learn how to use concept of chemical equilib & how it works. How to get to equilib, will it stay in equilib? Thermodynamics & kinetics.





Just b/c a rxn is favored in equilib doesn't mean there's enough activation energy to get it to go.

1st have to figure out if you can get rxns to go fast enough to get product (thermodynamically possible)




Sin-gas is CO & H_2 . People used to rely a lot on these.



rate of rxn dependent on amt of materials






H_2 & N_2 do not react by themselves.

N_2 has a really strong bond.

We want to be able to predict what conditions it'll break.


Rm temp, no rxn. At equilib, we have rxn that doesn't favor products. Large activation energy.

So we heat it. What happens?



$$K = \frac{[\text{C}]^l [\text{D}]^m}{[\text{A}]^j [\text{B}]^k} = \frac{\text{products}}{\text{reactants}}$$

Balanced equation 1st.



$$K = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

[] = concentration at equilib.

associated w/ rxn as written & as balanced.

Multiplying eqn by 2 means K^2

Reversing eqn means $\frac{1}{K}$

Chemical equilibrium is dynamic. Reactants form products, products form reactants

Temperature, Pressure add stress & therefore, R & P. Remove R or P, change add R or P. Change Volume, Concentration... pH... stuff like that.




Coordination complex

Consists of a transition metal ion & ligands

Ligands are things that attach other things. Neutral molecules or anions. Water could be a ligand but that does not account for all ligands.

e^- in a ^{coordinate} bond, both e^- come from the ligand.



Distance between D orbitals is smaller, making color.

NaCl in soln is clear.

CuSO₄ in soln is blue.

Transition metals tend to make color complexes

If you change the ligands, you change energy of D orbitals & therefore color.

You can tell, that, if you have Co^{+2} vs $Co + H_2O$.



Complex 1 $\rightleftharpoons$ Complex 2
pink blue

Add acid, & soln is pink. Adding acid has changed

equilib so you have small amt complex 2 + large amt complex 1.

If you saw purple, equal amts of complex 1 & 2.



1st part of lab, just observe Cu & Co.

2nd part, disrupt equilib & observe

$[\text{Cu}(\text{H}_2\text{O})_6]\text{Cl}_2$ Hexa aquacobalt (II) chloride,

Octahedral shape.

perfect octahedron



Overall charge can be +, -, or 0 depending on oxidation state of metal.

Depending on neutral/neg. charged ligands, charge on complex can change.

Complex Ion	Color
Shift	
Cobalt	
Copper	
Not Chromium/Nickel	

All of part A

1st half of B

(Equilibria of Co complexes)

2nd week

Equilibria of Cu complexes


Diff d orbitals of a transition metal ion have diff energies when that ion is surrounded by ligands.

When light of a particular λ corresponds to the spacing b/w 2 energy levels, an e^- is raised to that energy level & light of that λ is absorbed


Dynamic equilibrium

Products making reactants & vice versa at equal rates

Forward:



Reverse:



Equilib: $\rightleftharpoons$

Part A: Color of Coordination Compds (Complex ions)

1. Cobalt (II) Complex ions
2. Copper (II) Complexions & coordination compds
3. Summary of colors

Part B: Shifting Chemical Equilibria

1. Cobalt (II) Equilibria
2. Copper (II) Equilibria.

Concentrated acids & bases used. Goggles & gloves.

Chem Rxn, Equilib State

Monday, February 19, 2007

9:00 AM

Chemical rxns, equilibrium state

1st have to know if rxn is possible.

2nd have to know if it will occur in a reasonable period of time.

We can relate products & reactants


There are strong bonds in some reactants that have to be broken, requires activation energy.

At equilib & 300°C

$$[\text{N}_2] = .25 \text{ mol/L}$$

$$[\text{H}_2] = .15 \text{ mol/L}$$

$$[\text{NH}_3] = .090 \text{ mol/L}$$



$$K_{eq} = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3} = 9.6 \text{ L}^2/\text{mol}^2$$

9.6, or 10^{45} , or 10^{-32} tells you whether it favors reactants or products

units depend on coefficients

If $K > 1$, products favored

If $K < 1$, reactants favored

If $K = 0$, NR

If $K = 1$, equal amts of reactants & products.

At 127°C ,

$$[\text{NH}_3] = 3.1 \times 10^{-2}$$


$$[\text{N}_2] = 8.5 \times 10^{-1}$$

$$[\text{H}_2] = 3.1 \times 10^{-3}$$

$$\frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3} = 3.8 \times 10^4$$

There's 1 equilib constant for a given rxn at each temp. This tells us it greatly favors products when temp is lower. This is a rxn where if you lower temp, you get more product.

In theory, you could run it at a really low temp, but the lower you go, the slower the rxn goes. You want to still get to equilib.



Initially $1 \text{ M} [\text{H}_2]$
 $1 \text{ M} [\text{N}_2]$
 $0 \text{ M} [\text{NH}_3]$ hasn't been created yet

Finally $.921 \text{ M} [\text{N}_2]$
 $.963 \text{ M} [\text{H}_2]$
 $.157 \text{ M} [\text{NH}_3]$

$$6.02 \times 10^{-2} = K_{eq}$$

Immediately, you know it was a higher temp. You actually know it's higher than 300°C , b/c it's lower K_{eq} .

Initially $1 \text{ M} [\text{NH}_3]$
 $0 \text{ M} [\text{H}_2]$
 $0 \text{ M} [\text{N}_2]$

We work under conditions where we get to equilib

Finally $.399 \text{ M} [\text{N}_2]$
 $1.197 \text{ M} [\text{H}_2]$
 $.203 \text{ M} [\text{NH}_3]$

$$K_{eq} = 6.02 \times 10^{-2}$$

So we have to be working at same temp. [] not the same.

This is equilib position. There are ∞ equilib position systems. 1 equilib constant.

Solids & liquids (pure) not included in equilib constant
 If you add more reactants, you still get same K_{eq} at one temp.

How do we describe K for gases in terms of pressure?

$$PV = nRT$$

$$P = \left(\frac{n}{V}\right)RT$$

$$P = CRT \quad \text{Partial Pressure}$$

$$C = \frac{n}{V}, \quad C = \text{molar concentration.}$$

$$K_{eq} = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$


$$K = \frac{[NH_3]^2}{[N_2][H_2]^3} = \frac{C_{NH_3}^2}{C_{N_2} C_{H_2}^3}$$

$$K_p = \frac{P_{NH_3}^2}{P_{N_2} P_{H_2}^3} \quad C = \frac{P}{RT} \quad \left(\frac{P_{NH_3}}{RT} \right)^2 \frac{1}{\left(\frac{P_{N_2}}{RT} \right) \left(\frac{P_{H_2}}{RT} \right)^3} = \frac{P_{NH_3}^2}{P_{N_2} P_{H_2}^3} (RT)^2$$

$$\text{So } K_{eq} = K_p (RT)^2$$

When # moles balanced equation total on products = reactants $3X + 3Y \rightarrow 3XY$

Coefficients in balanced eqn determine K_{eq} in terms of K_p .




This'll work b/c
moles reactants = #
moles products.

$$K_p = K_c (RT)^{\Delta n} \leftarrow \sum \text{coefficients products} - \sum \text{coefficients reactants}$$

Anytime $\Delta n \neq 0$, $K_p \neq K_c$


Synthesis of NH_3 from N_2 & H_2 at $472^\circ C$ has equilib constant K_c of .105. K_p for this rxn?



$$\Delta n = -2$$

$$2 - (3+1) = -2$$

$$K_c = K_p (RT)^{-2} \quad R = .0821, \quad T = 472^\circ C, \\ .105 (0.0821 \cdot 472)^2 = K_p \quad K_c = .105$$



$$K = \frac{[CO_2] \cancel{[CaO]}}{\cancel{[CaCO_3]}} \quad \text{Solids don't get included b/c} \\ \text{The concentration is a constant}$$

$$K = [CO_2]$$




$$K_c = \frac{[H_2]^4}{[H_2O]^4}$$

At lower temp, $H_2O = L$

$$K_c = [H_2]^4$$

Reaction quotient



$$Q = \frac{[NH_3]_0^2}{[N_2]_0 [H_2]_0^3} \quad \text{Initial concentrations!}$$

If $Q = K$, system is at equilib

If $Q > K$, products get consumed & go to reactants' side.

If $Q < K$, too many reactants & to get to eq. will form more products

Q is a great predictor


Predictive power comes from Keq.

As you increase temp, it effects Keq.

Q = initial concentration equilb constant.

Compare Q to K

1. Write balanced eqn, w/ phases
2. Write equilb expression
3. List initial concentrations
4. Calculate Q, determine which way the shift to equilb will be.
5. Define change needed to reach equilb, apply changes to initial concentrations to find equilb concentrations
6. Substitute equilb concentrations. Calculate unknown.
7. Check K vs. equilb concentration.



$$\text{expression: } K_c = \frac{[\text{CO}_2][\text{H}_2]}{[\text{CO}][\text{H}_2\text{O}]}$$

.58 =

$$\text{Concentrations: } C = \frac{\text{mol}}{V} = \frac{1 \text{ mol CO} = \text{mol H}_2\text{O}}{50 \text{ L flask}} = .02 \text{ M [CO]}$$

water-gas shift rxn

1 mol CO

1 mol H₂O in 50 L flask.

K_c = .58 at 1000°C

$$[\text{CO}_2] = [\text{H}_2] = 0 \text{ b/c nothing yet.}$$

Q:

We know we aren't at equilb & so we have to go from reactants' side to products' side.

Initial	Change	Equilibrium
$[\text{CO}] = [\text{H}_2\text{O}] = .02$	$-x$	$.02 - x$

$[\text{CO}_2] = [\text{H}_2] = 0$	$+x$	x
------------------------------------	------	-----

ICE chart

$$\text{Equilb/unknown: } K = .58 = \frac{[\text{CO}_2]_{\text{eq}} [\text{H}_2]_{\text{eq}}}{[\text{CO}]_{\text{eq}} [\text{H}_2\text{O}]_{\text{eq}}} = \frac{(x)(x)}{(0.02-x)(0.02-x)} = \frac{x^2}{(0.02-x)^2}$$

$$\pm .76 = \frac{x}{.02-x}$$

$$\int \begin{matrix} x = .0086 \\ x = \cancel{-.0086} \end{matrix} \text{ no neg concentration.}$$


Plug into Equilibrium

$$.02 - .0086 = .0114 = [\text{H}_2\text{O}]_{\text{eq}}$$

LeChatelier's Principle


Allows us to predict changes that occur in a disturbed equilib sys.

Haber Process



Very important to shift to right
Haber noted pressure favors equilib concentration of NH_3


If a change is imposed on a sys @ equilib, position of equilib will shift to reduce change.



$\uparrow$ Add $\uparrow$, get more CaCO_3
Add CaCO_3 , get more product. $Q > K$, shift to left.
 $Q < K$, shift to right

Add anything that changes Q (more CO_2), reaction will respond. But CaCO_3 & CaO don't enter into Q .

Pressure change



Add H_2 or N_2 , drive sys toward product.

Add NH_3 , drive sys toward reactants.

Add inert gas, no effect $\rightarrow$

Change overall pressure, but not equilib b/c partial pressures don't change.

Halve the volume, change $[]$ & p.p.'s $\therefore$ change Q .


Remove NH_3 , drive toward product.

Increase volume, move toward side w/ most gas molecules.

Decrease, Drive from right to left. Larger # gas molecules $\rightarrow$ smaller #.

Exothermic rxn, heat on product side. If you want product, don't heat them up.


Endothermic, heat on reactant side.



Add N_2O_4 , get more product.

Inc. volume, more product.

Half volume, more reactants.




$$\frac{[\text{NO}]^2 [\text{Cl}_2]}{[\text{NOCl}]^2} = 1.6 \times 10^{-5} \text{ M}$$

$$[\text{NOCl}]_0 = .5 \text{ M}$$

$$[\text{NO}]_0 = [\text{Cl}_2]_0 = 0$$

So we don't need to calculate Q. we know rxn $\rightarrow$.



Initial	Change	Equilib
$[\text{NOCl}] .5 \text{ M}$	$-2x$	$.5 - 2x$
$[\text{NO}] = 0$	$2x$	$2x$
$[\text{Cl}_2] = 0$	x	x

$$\frac{(2x)^2(x)}{(1.50-2x)^2} = 1.6 \times 10^{-5} \quad \text{Solve for } x$$

↑
pretty small equilib constant, so
reactants aren't favored. Means change
is small. get rid of $2x$ on
bottom.


$$\frac{4x^3}{(1.5)^2} = 1.6 \times 10^{-5}$$

$$x^3 = \frac{(1.6 \times 10^{-5})(1.5)^2}{4}$$

$$x = \#$$

Plug # back in.

If you add something the product that reacts w/ product, you get more product b/c you're using some of it up during the rxn.



Change in pressure does not change the equilib constant.

Laws of Thermodynamics.

1st In any process, spontaneous or nonspontaneous, total energy of a system & its surroundings is constant

2nd In any spontaneous process, total entropy of a system & its surroundings increases

Spontaneous process is a possible process.

3rd Entropy of a perfect crystal at 0K is 0.

$$\Delta S_{\text{sys}} = \text{change in entropy}$$

$$\Delta S_{\text{uni}} = \Delta S_{\text{sys}} + \Delta S_{\text{surr}}$$

ΔS for spontaneous process is (+)



DNA Double helix

Entropy of system goes down when it forms.

However the entropy of everything else increases.

This is a spontaneous process b/c $S_{\text{surr}} \uparrow$ while

$S_{\text{sys}} \downarrow$


$$\Delta S_{\text{uni}} = \underset{(-)}{\Delta S_{\text{sys}}} + \underset{(+)}{\Delta S_{\text{surr}}}$$

The entropy change must be positive for the universe for the process to be spontaneous.

You can't get back to an ordered state b/c of probability -
There is only 1 ordered state, and very many disordered states.
The probability of getting the ordered state is slim. Alex
is a testament to that. So nature favors disorder.


Phase Changes, Gibbs Free Energy

Thursday, February 22, 2007
8:58 AM




ΔS_{sys} has (+) sign
endothermic process. Takes heat in.

Phase changes in general



but it really looks more like



As ΔS_{sys} has (+) sign b/c sys gets more disordered,
 ΔS_{surr} has (-) sign b/c we must drive rxn to go from
 $\text{L} \rightarrow \text{G}$ or $\text{G} \rightarrow \text{L}$ under diff situations. Can't be spontaneously
going back, so ΔS_{surr} has to be (-).

Temperature plays a key role.

Heat flows out of surroundings & into sys to decrease
entropy. This is b/c it's an endothermic process.

ΔS_{sys} or ΔS_{surr} controls ΔS_{uni}

ΔS_{surr} controlled by heat flow

Higher the temp of the surroundings, less significant impact
of heat flow is.

ΔH_{sys} is (+) , ΔS_{surr} is (-).

$$\Delta S_{\text{surr}} \text{ exothermic } : \frac{+q(J)}{\text{Temp}(K)}$$

$$\text{endothermic } : \frac{-q(J)}{\text{Temp}(K)}$$

$$\Delta S_{\text{surr}} = -\frac{\Delta H}{T}, \Delta H \text{ for rxn. If } \Delta S_{\text{sys}} \text{ is } (+), \Delta S_{\text{surr}} \text{ is } (+)$$

Constant T & P

Water melting

ΔS_{sys} is (+)

ΔS_{surr} is (-) when does ΔS_{sys} win?

At high temp.

$\Delta S_{\text{sys}} \equiv$ invariant for physical state


$$\Delta S_{\text{surr}} = -\frac{\Delta H}{T} \quad \# \text{ gets smaller at higher } T.$$



O! Bessy
does her
Mudder
impression

Moo, moo, I need
help!

25°C



so (-) ΔS_{sys} . More ordered. Makes sense.

Gibb's Free Energy

$$\Delta G = \Delta H - T\Delta S_{\text{sys}}$$

$$\Delta S_{\text{uni}} = -\frac{\Delta G}{T} \quad (-)\Delta G \text{ means spontaneous process}$$

$$\Delta G = \Delta H - T\Delta S_{\text{sys}}$$

$$-\frac{\Delta G}{T} = -\frac{\Delta H}{T} + \Delta S_{\text{sys}}$$


$$-\frac{\Delta G}{T} = \Delta S_{\text{surr}} + \Delta S_{\text{sys}}$$

$$\Delta S_{\text{uni}} = \Delta S_{\text{surr}} + \Delta S_{\text{sys}}$$

$$-\frac{\Delta G}{T} = \Delta S_{\text{uni}}$$



It works



ΔS_{sys} (-).

$$\Delta G = \Delta H - (-T\Delta S) = \Delta H + T\Delta S$$

↑
must be (-) b/c it
is spontaneous.

when ΔH is (+) & ΔS is (-) then ΔG will be (+) & it

When ΔH is (+) & ΔS is (-), the ΔG will be (+) & it will always be nonspont.



When ΔH is (-) & ΔS is (+), the ΔG will be (-) & it will always be spont

When ΔH is (-) & ΔS is (-), the ΔG will be spont if T is low. Will be nonspont if T is high. Exothermic.

When ΔH is (+) & ΔS is (+), the ΔG will be spont if T is high. Will be nonspont if T is low. Endothermic
Ice melting.

Enthalpically driven vs. Entropically driven.

ΔH is (-)

ΔH is either (+) or very small.

Endothermic can never be enthalpically driven. Ice melting is entropically driven.

Level of predictability.

I ♥ CHEESE

Gibbs Free Energy, Entropy

Monday, February 26, 2007
8:59 AM

Gibb's Free energy was a restatement of the 2nd law of thermodynamics

$$\Delta G = \Delta H - T\Delta S$$

You can make predictions w/ this.
(-) is spont, ΔS is (+)
(+) is non spont

Quantitative Predictions

Calculate how much Ag_2O must be heated (to what T) to decompose Ag_2O to O_2 gas + Ag metal.



$$\Delta H = 62.1 \text{ kJ}$$
$$\Delta S = 132.75 \text{ J/K}$$


Endothermic process, so entropically driven. Must be heated to work.

$$0 = \Delta H - T\Delta S$$

$$T\Delta S = \Delta H$$

$$T = \frac{62.1 \text{ kJ} \left(\frac{1000 \text{ J}}{1 \text{ kJ}} \right)}{132.75 \text{ J/K}} = 468 \text{ K}$$

Anything $\uparrow$ 468 K will be spont.



$$\Delta H^\circ = 98.8 \text{ kJ}$$
$$\Delta S^\circ = 141.55 \text{ J/K}$$

$$\Delta G = \Delta H - T\Delta S \quad \text{given}$$

$$\Delta G = 98.8 \text{ kJ} - (298 \text{ K})(141.55 \text{ J/K})$$

$$\Delta G = +56.6 \text{ kJ}, \text{ not even close to spontaneous.}$$


Have to calculate T to make it so.

It will be spont when 298 is way higher.

$$0 = 98.8 \text{ kJ} - T(141.55 \text{ J/K})$$

$$T = \frac{98.8 \text{ kJ}}{141.55 \text{ J/K}}$$

$$T = 698 \text{ K} \text{ any temp above this makes spont process}$$



$$\text{Fe}_2\text{O}_3 \quad \Delta H = -824 \text{ kJ/mol}$$

$$S = +87.45 \text{ J/mol}\cdot\text{K}$$

$$\text{Al}_2\text{O}_3 \quad \Delta H = -1673 \text{ kJ/mol}$$



$$S = +50.925 \text{ J/mol}\cdot\text{K}$$

$$\Delta H^\circ_{298} = -849 \text{ kJ/mol}$$

$$\Delta S^\circ = -36.48 \text{ J/mol}\cdot\text{K}$$

enthalpically driven

Free energy change at 298K = -838 kJ/mol
 If you reverse it, it's pos. so it shouldn't go.
 Eberhard's demo



Entropy change

This is (+).

This is (+)

(-) ordered solid



(-) 4 molecules to 2

Can't tell w/out values

(-) ordered solid



entropy change.



This has a greater value b/c it's a more complex molecule.
 This is a smaller molecule.

However, there are more NO_2 's than N_2O_4 's, &
 $2(240) > 304.2$

$$\Delta S_{\text{rxn}}^\circ = \Delta S_{\text{prod}} - \Delta S_{\text{react}}$$

$$\Delta S_{\text{rxn}}^\circ = 480 - 304.2 = (+) 175.8 \text{ J/K}\cdot\text{mol}$$

$\text{H}_2\text{O}(\text{l})$ has a really low S° value b/c of H bonding keeps order.

2012) ...





Spontaneous?

Need to know ΔG values. $\Delta G_{\text{prod}} - \Delta G_{\text{reactants}}$.

or Find ΔH & ΔS . Find ΔG

or Find related rxns.



Flip one, mult, add, & get ΔG°

Gibbs Free Energy, Hess's Law


Tuesday, February 27, 2007
9:01 AM

1a/ 33, 37, 39, 43, 47, 53, 63, 69, 74, 79, 84, 88, 89

ΔG tells us the driving force for a rxn, and the spontaneity.
Can be calculated w/ Hess' Law, directly calculating $\Delta G_{\text{prod}} - \Delta G_{\text{reactants}}$, or $\Delta H - T\Delta S$

Hess's Law takes 2 eqs, rearranges them, & adds together to get ΔG°




$$\Delta G_{\text{rxn}} = \sum \nu_p \Delta G_f^\circ \text{prod} - \sum \nu_r \Delta G_f^\circ \text{react}$$



$$\begin{aligned} \Delta G^\circ &= (4 \text{ mol } 96.6 \text{ kJ/mol NO}) + \\ &\quad 6 \text{ mol } (-237.2 \text{ kJ/mol H}_2\text{O}) \\ &\quad - (4 \text{ mol } (-16.5 \text{ kJ/mol NH}_3) + \\ &\quad 5 \text{ mol } (0 \text{ kJ/mol O}_2)) \end{aligned}$$

elems in standard states,
 ΔG° of formation is 0

ΔG_f° is change in free energy that will occur if standard states are added



Convert reactants all the way to products $\rightarrow \Delta G^\circ$.

In the lab, we can't do this b/c you start w/ diff concentrations, temps, & pressures.

If you change the temp & pressure of rxn, you change the rxn (where it goes)


$$\Delta G = \Delta G^\circ + RT \ln Q$$

↑ any conditions ↑ standard set of conditions ↑ equilib expression w/ concentration at start of experiment

Q is rxn quotient

We can calculate ΔG° for each rxn we're interested in. $RT \ln Q$ will give us correction term.

$$\Delta G^\circ = -22.5$$




1 atm N_2 , 3 atm H_2 , .020 atm NH_3
 .01 atm N_2 , .03 atm H_2 ,

$$Q_p = \frac{(0.020)^2}{(1)(3^2)} = 1.5 \times 10^{-5}$$

$$\Delta G = (-33 \text{ kJ}) + (8.35/\text{K})(298) \ln(1.5 \times 10^{-5})$$

$$\Delta G = -60.5 \text{ kJ}$$

You use a diff R value per Q_p + Q_c



once you get to 0,
 $G = 0$

$$0 = \Delta G^\circ + RT \ln Q$$

$$-\Delta G^\circ = RT \ln Q \text{ at equilib}$$

$$\Delta G^\circ = -RT \ln K$$

Now we can calculate K.

If it goes to equilib, & its activation energy is reasonable, then the ΔG will tell what direction it will lean.



catalysts lower activation energy w/o changing ps of react. & prod.

Heating breaks up bonds & might help. It will change K. Change relative positions, though.

A catalyst interacts w/ reactants to stretch & weaken bonds.

Perm - H_2O_2 goes to $\text{H}_2 + \text{O}_2$. Thermodynamically unstable, spontaneous. Meaning large activation barrier keeps it from going right away. Add catalyst & see what happens

Free Energy, Nuclear Rxns

Wednesday, February 28, 2007
9:04 AM

You can calculate ΔG for any conditions as long as you had ΔG° . ΔG° for a rxn is taking all reactants & converting them all to products. We don't really do this.

At equilib, $\Delta G = 0$
 $0 = \Delta G^\circ + RT \ln Q$ ← at equilib, $Q = K$
 $-\Delta G^\circ = RT \ln K$



Now we can calculate K w/ ΔG° & temp & R .
Enormous equilib K or really small K have been tabulated w/ calculation, not rxn. This is why $-\Delta G^\circ = RT \ln K$ is really important

$\Delta G^\circ < 0$, then it is spont. Will go to prod, & so $K > 1$

$\Delta G^\circ > 0$, then it is nonspont. Will favor reactants, so $K < 1$

If you know RT & K , you can find ΔG° .

At equilib, ΔG is 0. So it's uphill both ways & forward & reverse are nonspont.



$\Delta G^\circ = -RT \ln K$
At 25° $\ln K = \frac{-(-33.0 \text{ kJ})}{(8.3)(298 \text{ K})}$

$K = 6.5$ This is K_p

$K_p = K_c RT^{\Delta n}$ ($R = 0.0821$)
b/c $K_c = K_p RT^{-\Delta n}$

ΔG°_f Mercury at 25°C is 31.85 kJ/mol. Vapor pressure of mercury at 25°C?

$K_p = P_{H_2O}$ b/c pure liquid not in eqn.

$$\ln K_p = \frac{\Delta G^\circ}{RT} = \frac{-31.85 \text{ kJ}}{(8.314)(298 \text{ K})} = , K_p = 2.6 \times 10^{-6}$$

this is pretty high,
and that's why it's
volatile.

As $T \uparrow$, $K \uparrow$ endo. As $T \uparrow$, $K \downarrow$ exo.


$$\Delta G^\circ = -RT \ln K = \Delta H^\circ - T\Delta S^\circ$$

$$\ln K = \frac{-\Delta H^\circ}{RT} + \frac{T\Delta S^\circ}{RT}$$

$$\ln K = \frac{-\Delta H^\circ}{R} \left(\frac{1}{T} \right) + \frac{\Delta S^\circ}{R}$$

2 variables here - K & T ,
so you get a linear graph

I ♥ Kenisha!



How does W relate to ΔG ?

Max theoretical work is ΔG .

$$W_{\max} = \Delta G$$

We can never get to $W_{\max}$, though.

As you do work, you lose energy to heat. Heat increases entropy of the environment. And entropy is not a useful form of energy - there's no further component.

Thermal energy increases entropy of the surroundings.

1st law - the best one can do w/ this process is break even.

Calculate ΔG° for



$$\Delta G = \Delta G^\circ + RT \ln Q$$

At equilib $\Delta G = 0$

$$\Delta G^\circ = -RT \ln K$$

$$\Delta G^\circ = (-8.314)(373 \text{ K}) \ln(5.3 \times 10^9)$$

$$\Delta G^\circ = -69,400 \text{ J} \quad \underbrace{-69.4 \text{ kJ}}_{\text{max work}}$$




It's spontaneous process. If $\Delta G^\circ_{\text{rxn}} > 0$, most work you could get out.

Is 0 b/c it's a nonspont process.

Nuclear rxns, nuclear chemistry

Radioactive materials. radioactive dating. power & other operation of

Radioactive materials, radioactive dating, power & other generation of stuff. Homer Simpson works at a Nuc power plant.



lead blocks radioactive particles.
goes through glass

The Los Alamos sample is no longer radioactive. It's all decayed now. Nuclear radius is unstable.

There's radioactive radon in the air in dust particles.


Nuclear Rxns

Monday, March 05, 2007
8:58 AM

Nuclear Rxn

Fundamentally, rxn occurs in nucleus. No molecules reacting. You can have nuc. rxns occurring in molecules, but actual rxn occurs in nucleus. Radioactive sugar leads to labelling.


Radioactive material is constantly decaying, so we're constantly losing radioactive material.



21 neutrons
19 protons

22 neutrons
18 protons

proton converted to neutron.



β particle emission

Positron

e^- capture

α

γ

spont. fission

Decrease in N-P ratio


Inc.

Inc

Inc

None

None



Half-life is time required for half of material to go through half of decay

half-life is time required for half of material to go through half of decay



Uranium has a half-life of 4 billion years.
Some materials have half-lives of seconds.

If we know how old something is by half-lives, then it's important to know this.

You need to see the types of particles liberated (like you don't want γ b/c too high energy) and the half-lives for storage purposes.

$$t_{1/2} = \frac{0.693}{k} \quad \text{rate constant associated w/ decay rxn}$$

Nuclear reactors are not great things to have in the backyard for safety concerns and terrorism concerns.

70% of Illinois is powered by nuc. reactors.

Radioactive hydrogen is tritium. Mass 3


Deuterium is H w/ mass 2. Heavy water.

^{131}I used for thyroid

^{60}Co used for cancer treatment

Thallium 201 used as tracer after heart attacks

Tc-99m for bone abnormalities, heart damage in heart attack



displacement reaction.

calculate amt radioactive material, calculate amt insulin.

Radiation pill is how you protect yourself from radiation $\rightarrow$ saturate thyroid w/ I before attack so radioactive I doesn't come in.

Radioactive Dating, Fusion

Tuesday, March 06, 2007
9:05 AM

Radioactive Dating $t_{1/2} = \frac{0.693}{k}$

Derivation $\ln \frac{N_t}{N_0} = -kt$ Integrated 1st order rate law.
 N_0 is amt radioactive material at start of process.
 N_t is amt left


$$\ln \frac{50}{100} = -kt_{1/2}$$

$$-0.693 = -kt_{1/2}$$

^{14}C $t_{1/2} = 5730 \text{ y}$

^{14}C exists in atm due to cosmic ray bombardment of earth

React w/ C & N.



Then ^{14}C reacts w/ O, converts to $^{14}\text{CO}_2$
 plants consume it (CO_2)

Libby recognized that once animals & plants die,
 $^{14}_6\text{C} \rightarrow ^{14}_7\text{N} + ^0_{-1}\text{e}$ they no longer consume
 ^{14}C & they liberate it.

If we know amt ^{14}C in living things & something dies, we
 can back calculate how old object is.

Biggest assumption is that ^{14}C - ^{12}C ratio hasn't changed
 in thousands of years

Tree died, gave 7 disintegrations of ^{14}C per min per gram carbon.
 Present day gives 15.3 disintegrations/min. when did it die?

We use ^{14}C b/c it's accurate but long.

$$^{14}\text{C} \ t_{1/2} = 5730 \text{ y}$$

$$\ln \frac{N_t}{N_0} = -kt$$

$$t_{1/2} = \frac{0.693}{k}$$

$$k = \frac{0.693}{t_{1/2}}$$

$$\ln \frac{7}{15.3} = -\frac{0.693}{5730} t$$

$$t = 6.5 \times 10^3 \text{ yrs old}$$

Fission splits

Fusion puts together

It's possible to do fusion w/ certain things below
 60 b/c of the relative stabilities of elmts.
 60 is mass #



Any time you go toward a more stable nucleus, it is a spont. process.


The probs w/ having 2 nuclei or making them fuse is spont. but the rate is really slow unless we know something about activation barrier. w/ fission, activation barrier is huge.



Fission is splitting of nuclei. Release a lot of energy when they do that.

ATLANTA

Peace up,
A-Town
Yeah Yeah




As this happens, it rises in energy. Basis for bomb is the chain reaction that cascades out of control & explodes.

To control process, control amt of neutrons.

Nuclear reactors don't have chain rxn or we'd die.

Neutrons are controlled by control rods - they absorb neutrons.

Fusion brings them together.



deuterium + tritium $\rightarrow$ helium / α particle + neutron

This generates an enormous amt of energy

Another benefit is just helium and a neutron vs all that radioactive waste.

No chain rxn. Very controllable. No cascading out of control.

Activation energy, however, is enormous & we don't have a catalyst.

Other option is really high temps. 40 million° on sun.

So have to generate that kind of heat w/out melting everything around the reactor.

Hydrogen bomb is uncontrolled fusion. You have those starting materials & you initiate the rxn by slamming them into each other. You use fission to produce enough heat to initiate fusion.



