

Classifying Diff Eqns

Monday, March 26, 2007

8:59 AM

Comp lab Th. 10-1100

What's a diff eqn?

Eqn involving an unknown fn & its derivatives

e.g. $\frac{dx}{dt} + x = 0$ $x = e^{-t}$, $x = 0$, $x = 3e^{-t}$
 $x = Ce^{-t}$

$m \frac{d^2 y}{dt^2} + b \frac{dy}{dt} + ky = 0$ $m, b, k \geq 0$ damped spring
 damping coeff. = b

$\frac{d^2 \theta}{dt^2} + \frac{g}{l} \sin \theta = 0$ $g = \text{grav acc.}$ Spring constant = k
 Pendulum eqn $l = \text{length moment arm}$

Terminology for classifying diff eqns

Order - power / degree of highest derivative

linear or nonlinear "power of unknown in eqn". If only 1st power appears of the unknown fn, then eqn is linear; otherwise nonlinear.

$\frac{d^2 \theta}{dt^2} = -\frac{g}{l} \sin \theta$ is nonlinear b/c of the $\sin \theta$

Systems of diff eqns

ex $\vec{F} = m\vec{a}$, 3D space
 $F_1 = m \frac{d^2 x}{dt^2}$


$\vec{a} = \frac{d^2}{dt^2} \langle x, y, z \rangle$
 $\vec{F} = \langle F_1, F_2, F_3 \rangle$

$F_2 = m \frac{d^2 y}{dt^2}$


$F_3 = m \frac{d^2 z}{dt^2}$

$\vec{F} = -\frac{GMm}{r^2} \hat{r}$

$= \left\langle -\frac{GMm x}{(x^2 + y^2 + z^2)^{\frac{3}{2}}}, -\frac{GMm y}{(x^2 + y^2 + z^2)^{\frac{3}{2}}}, -\frac{GMm z}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \right\rangle$



$\vec{r} = \langle x, y, z \rangle$
 $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$



$$|\vec{r}| = r = \sqrt{x^2 + y^2 + z^2}$$

$$\hat{r} = \frac{\vec{r}}{|\vec{r}|} = \frac{\langle x, y, z \rangle}{\sqrt{x^2 + y^2 + z^2}}$$

System of 3 second-order differential nonlinear eqns
 Ordinary vs. Partial differential eqns (ODEs) (PDEs)
 Has to do w/ 1 variable or 2 variables that fn depends on
 If unknown depends on >1 variable, PDE.

Heat eqn

$$\frac{\partial T}{\partial t} = K \frac{\partial^2 T}{\partial x^2}$$

$T(x, t) = \text{temp}$

First example

Spherical particle falling through highly viscous fluid



Forces - Gravity, $\frac{\text{drag}}{\uparrow \text{viscous}}$, buoyancy

Viscous drag depends on viscosity, temp, velocity

$$\text{Grav} = -mg\hat{j}$$

$$F = ma = m \frac{d\vec{v}}{dt} = -m \frac{dv}{dt} \hat{j} = -mg\hat{j}$$

$$\vec{v} = -v\hat{j}$$

↑ speed

For small enough particles $F_{\text{drag}} \approx 6\pi\eta a v$

Result due to Stokes

$a = \text{rad of sphere}$


$\eta = \text{viscosity}$

$v = \text{speed}$

good approx
for small Reynolds

$$R = \frac{\rho a v}{\eta}$$

only true if $R \ll 1$



ρ = density of fluid
 a = radius of spherical particle
 μ = viscosity of fluid

Assume: $\vec{F} = \vec{F}_{\text{grav}} + \vec{F}_{\text{drag}} \hat{=} m\vec{a}$

$$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d}{dt}(-v\hat{j}) = (-\hat{j}) \frac{dv}{dt}$$

$\hat{j}$ | unit vector pointing in vertical direction.

$$\vec{F}_{\text{grav}} = mg(-\hat{j})$$

Assume Stokes' Approx to drag

$$F_{\text{drag}} \approx 6\pi\mu a v$$

Good approx if Reynold's # = $R = \frac{\rho a v}{\mu} \ll 1$

$$\vec{F}_{\text{drag}} = 6\pi\mu a v \hat{j}$$

$$-m \frac{dv}{dt} \hat{j} = -mg \hat{j} + 6\pi\mu a v \hat{j}$$

$$\boxed{\frac{dv}{dt} = g - \left(\frac{6\pi\mu a}{m}\right)v}$$

= diff eq. for the speed of the particle v as a function of time

Solution depends on initial speed

~ "initial condition" at $t = t_0$ $v(t_0) = v_0$

if v is pos it's going down

if v is neg it's going up

Euler's Approx Numerical soln

Based on the fact that $\frac{dv}{dt} = \lim_{\Delta t \rightarrow 0} \frac{v(t+\Delta t) - v(t)}{\Delta t}$

$$\frac{dv}{dt} \approx \frac{v(t+\Delta t) - v(t)}{\Delta t}$$

if t is small

$$\frac{dv}{dt} = g - \left(\frac{6\pi\eta a}{m} \right) v$$

$$v(t_0) = v_0$$

$\left(\frac{1}{\tau} \right)$ just b/c

t	v
t_0	v_0
$t_0 + \Delta t$	$v_1 = v_0 + \left[g - \frac{v_0}{\tau} \right] \Delta t$
$t_0 + 2\Delta t$	v_2
$\vdots$	$\vdots$
$t_0 + n\Delta t$	v_n
$t_0 + (n+1)\Delta t$	v_{n+1}

$$v_1 = v(t_0 + \Delta t) \approx \underbrace{v(t_0)}_{v_0} + \underbrace{\frac{dv}{dt}(t_0)}_{g - \frac{v_0}{\tau}} \Delta t$$

Taylor series
but not higher order
terms
Linear Approx

$$v_1 \approx v_0 \left[g - \frac{v_0}{\tau} \right] \Delta t$$

Assume we have $v_n \approx v(t_0 + n\Delta t)$

$$\text{What is } v_{n+1} = v(t_0 + (n+1)\Delta t) = v(t_0 + n\Delta t + \Delta t)$$

Linearize v at time $t_n \approx t_0 + n\Delta t$

$$v_{n+1} \approx \underbrace{v(t_0 + n\Delta t)}_{v_n} + \underbrace{\frac{dv}{dt}(t_0 + n\Delta t)}_{g - \frac{v_n}{\tau}} \Delta t$$


$$v_{n+1} = v_n + \left[g - \frac{v_n}{\tau} \right] \Delta t$$

$$v_2 = v_1 + \left[g - \frac{v_1}{\tau} \right] \Delta t$$


$$\frac{dy}{dx} = f(y) \quad \text{or} \quad \frac{dy}{dx} = f(y, x), \quad y(x_0) = y_0$$

Qualitative Analysis of Solns

Based on direction field



$$\boxed{\frac{dv}{dt} = g - \frac{v}{\tau}}$$




$$\boxed{\frac{dv}{dt} = g - \frac{v}{\tau}}$$

$$v=0 \Rightarrow \frac{dv}{dt} = g$$

if $v=g\tau$, no slope

$$\frac{dv}{dt} = g - \frac{v}{\tau} = \text{slope of tangent line determined by direction field}$$



Analytic soln $v(t) = \text{---}$

$$\frac{dv}{dt} = g - \frac{v}{\tau} \quad \leftarrow \text{separable } 1^{\text{st}} \text{ order differential eqn}$$

$$v(t) = v_0$$

$$\tau = \frac{m}{6\pi\eta r a} = \text{constant}$$

Separable eqn soln method

$$\frac{dy}{dx} = f(y)g(x)$$

if $f(y)$ is 0, y isn't changing

$$\int \frac{dy}{f(y)} = \int g(x) dx \quad \text{indefinite}$$

$$\frac{1}{g - \frac{v}{\tau}} \frac{dv}{dt} = 1$$

$$\frac{\tau}{g\tau - v} \frac{dv}{dt} = 1$$

$$\int \frac{1}{g\tau - v} \frac{dv}{dt} = \int \frac{1}{\tau} dt$$

$$\int \frac{1}{g\tau - v} dv = \int \frac{1}{\tau} dt$$

$$-\ln|g\tau - v| = \frac{t}{\tau} + C$$

$$-\ln|T-r| = \frac{t}{T} + C$$

$$\ln|T-r| = -\frac{t}{T} - C$$

$$e^{\downarrow} = e^{\downarrow}$$


$$|T-r| = e^{-\frac{t}{T}}$$

$$T-r = K e^{-\frac{t}{T}} \quad e^{-C} \text{ constant } K$$

Evaporating Snowball

Wednesday, March 28, 2007

9:01 AM



$$\frac{dv}{dt} = g - \frac{v}{\tau} \quad \tau = \frac{m}{6\pi\eta a}$$

$v(0) = v_0$ initial speed downwards

$$\int g - \frac{1}{\tau} \frac{dv}{dt} dt = \int 1 dt \quad \text{provided } v_0 \neq g\tau$$

Special soln $v(t) = g\tau$ (singular soln)

$$\left(-\frac{1}{\tau}\right) \int g - \frac{1}{\tau} dv = \left(-\frac{1}{\tau}\right) \int dt$$

$$\int \frac{1}{v - g\tau} dv = -\frac{t}{\tau} + C$$

$$\ln|v - g\tau| = C - \frac{t}{\tau}$$

$$v - g\tau = e^C e^{-\frac{t}{\tau}}$$

$$v = g\tau + K e^{-\frac{t}{\tau}}$$

$$\text{Check: } \frac{dv}{dt} = -\frac{1}{\tau} K e^{-\frac{t}{\tau}}$$


$$g - \frac{v}{\tau} = g - \left[g + \frac{K}{\tau} e^{-\frac{t}{\tau}} \right]$$

$$= -\frac{1}{\tau} K e^{-\frac{t}{\tau}}$$

$$t=0, v=v_0 = g\tau + K$$

$$K = v_0 - g\tau$$

$$v(t) = g\tau + (v_0 - g\tau) e^{-\frac{t}{\tau}} \quad \text{when } t=0, v(t)=v_0$$



as $t \rightarrow \infty$, $v \rightarrow g\tau$
terminal

all solns decay exponentially toward terminal velocity.

$$\tau_{\text{form}} = \eta \tau = \frac{9\eta}{6\pi\mu a}$$

We assumed Stokes' eqn for viscous drag, which is valid if Reynolds # $R \ll 1$

$$R = \frac{\rho a v_{\text{typical}}}{\mu}$$

plug into this

$$= \rho a \frac{9\eta}{6\pi\mu^2 a} = \frac{\rho \eta}{6\pi\mu^2} \xrightarrow{\text{density of the fluid}} \frac{\rho \eta}{6\pi\mu^2} \xrightarrow{\text{density of the sphere}} \left(\frac{4}{3}\pi a^3\right)$$

$$\frac{\rho_{\text{fluid}}}{\mu^2} \left(\frac{4}{3}\pi\right) (\rho_{\text{sphere}} a^3) \ll 1$$

↑
fluid properties

$$\frac{\rho_{\text{fluid}}}{\mu^2} \left(\frac{2}{9}g\right) (\rho_{\text{sphere}} a^3) \ll 1$$

$$a \ll \left(\frac{\mu^2}{\rho_{\text{fluid}} \rho_{\text{sphere}}}\right)^{\frac{1}{3}} \left(\frac{9}{2g}\right)^{\frac{1}{3}}$$

Evaporating snowball (spherical)

Assume evaporation rate is $\propto$ surface area S



Surface area S , Volume V $\frac{dV}{dt} = \text{evap rate}$

$$\frac{dV}{dt} \propto S \quad \frac{dV}{dt} = -KS \quad K \text{ is a pos. constant}$$

$$V = \frac{4}{3}\pi a^3 \quad S = 4\pi a^2$$

$$a = \left(\frac{3V}{4\pi}\right)^{\frac{1}{3}} \quad S = 4\pi \left(\frac{3V}{4\pi}\right)^{\frac{2}{3}}$$

$$S = (4\pi)^{\frac{1}{3}} (3V)^{\frac{2}{3}}$$

initial condition,
 $V(0) = V_0$

$$-KS = -K(4\pi)^{\frac{1}{3}} (3V)^{\frac{2}{3}}$$

$$\frac{dV}{dt} = -\beta V^{\frac{2}{3}} \quad \beta \text{ is combo of } K(4\pi)^{\frac{1}{3}} (3^{\frac{2}{3}})$$

$$V(0) = V_0$$

$$\int \frac{1}{V^{\frac{2}{3}}} \frac{dV}{dt} dt = \int -\beta dt \quad \text{provided } V_0 \neq 0$$

$$\int V^{-\frac{2}{3}} dV = \int -\beta dt$$

indefinite integral

$$3V^{\frac{1}{3}} = -\beta t + C$$

$$\text{at } t=0, V=V_0$$

$$3V_0^{\frac{1}{3}} = C$$

$$3V^{\frac{1}{3}} = -\beta t + 3V_0^{\frac{1}{3}}$$

$$V = \left(V_0^{\frac{1}{3}} - \frac{\beta}{3} t \right)^3$$

definite integral

$$\int_{V_0}^{V(t_f)^{\frac{2}{3}}} V^{-\frac{2}{3}} dV = -\beta \int_{t_0=0}^{t_f} dt$$

$$3V^{\frac{1}{3}} \Big|_{V_0}^{V_f} = -\beta t_f$$

$$3V_f^{\frac{1}{3}} - 3V_0^{\frac{1}{3}} = -\beta t_f$$



same thing

Volumes must be pos. So this soln valid for $t \leq \frac{3V_0^{\frac{1}{3}}}{\beta}$

$$t = t_{\text{nothing}} = \frac{3V_0^{\frac{1}{3}}}{\beta} \quad \text{soln} = 0 \quad \text{for } t > \frac{3V_0^{\frac{1}{3}}}{\beta}$$

$$V_0 = 1 \text{ cm}^3$$

$$V_0 = 10 \text{ cm}^3$$



$$V(t) = \left(V_0^{\frac{1}{3}} \left(1 - \frac{\beta t}{3 V_0^{\frac{1}{3}}} \right) \right)^3$$

$$= V_0 \left(1 - \frac{\beta t}{3 V_0^{\frac{1}{3}}} \right)^3$$

ratio of final to initial $\frac{V(t)}{V_0} = \left(1 - \frac{\beta t}{3 V_0^{\frac{1}{3}}} \right)^3$

$$\frac{V}{V_0}$$



all curves
collapse to single curve

Existence & Uniqueness Theorem

Friday, March 30, 2007
9:01 AM



p. 23 in book

Existence & Uniqueness Theorem for First Order ODEs (Ordinary Differential Eqns)

$$* \frac{dy}{dx} = f(x, y) ; y(a) = b$$

conditions on this fn so you get existence & uniqueness

Thm Suppose $f(x, y)$ is continuous on some rectangle in the xy plane. Containing (a, b) in its interior



(1) Then (*) has at least 1 solution defined on some open interval J containing the pt $x=a$ [Statement says there exists a soln]

(2) If, in addition, the partial derivative $\frac{\partial f}{\partial y}$ is continuous on the rectangle then the soln is unique on some (perhaps smaller) open interval J_0 containing $x=a$ [Uniqueness]

Revisit Snowball Problem

$$\frac{dV}{dt} = -\beta V^{\frac{2}{3}} , V(t_0) = V_0$$

(1) $f(t, V) = -\beta V^{\frac{2}{3}}$ is a continuous function on a neighborhood (t_0, V_0)
exists a soln to initial value prob

(2) $\frac{\partial f}{\partial V} = -\frac{2}{3}\beta V^{-\frac{1}{3}}$ There's a prob at $V=0$. Continuous everywhere except $V=0$ so initial value problem (*) has a unique soln provided $V_0 \neq 0$

If $V_0 = 0$, there was no snowball. CSM would choose this soln. "blows puff of smoke" There was no snowball!

Section 1.5

First order linear eqns & their solns

$$\boxed{\frac{dy}{dx} + p(x)y = q(x)} \quad y(a) = b$$

depending on p & q you can determine if soln exists & is unique.

$$\frac{dy}{dx} = q(x) - p(x)y = f(x, y)$$

$$- p(x)y = \frac{\partial f}{\partial y}$$

Ex. 1 Soln of falling head problem using integrating factor

((⊗))) $\frac{dv}{dt} = g - \frac{v}{\tau} \quad \tau = \frac{m}{6\pi\eta a}$

$$\frac{dV}{dT} = 1 - V$$

Nondimensionalizing

$$v = (g\tau) V(T)$$

$\uparrow$ units here $\uparrow$ dimensionless
 $t = \frac{T}{\tau} \cdot \tau$
 $\uparrow$ units here

$$\frac{dv}{dt} = \frac{d}{dt}(g\tau V) = g\tau \frac{dV}{dt}$$

$$= g\tau \frac{dV}{dT} \frac{dT}{dt} = g \frac{dV}{dT}$$

$\underbrace{\frac{dT}{dt}}_{\frac{1}{\tau}}$

$$g - \frac{v}{\tau} = g - \frac{g\tau V}{\tau}$$

$$g \frac{dV}{dT} = g(1 - V)$$

$$\frac{dV}{dT} = (1 - V)$$

$$\frac{dV}{dT} + V = 1$$

$$e^T \frac{dV}{dT} + e^T V = e^T \quad e^T \text{ is integrating factor}$$

$$\int \frac{d}{dT}(e^T V) dT = \int e^T dT$$

$$e^T V = e^T + C$$

$$V = 1 + Ce^{-T}$$

Ex. 2 $\frac{dy}{dx} + 2xy = 2x \leftarrow$ separable $\frac{dy}{dx} = 2x(1-y)$
 but we'll solve w/ an integrating factor. "integrating factor" $m(x)$

$$m(x) \frac{dy}{dx} + m(x) 2xy = 2xm(x)$$

$$\frac{d}{dx} (?) = 2xm(x)$$

try: $\frac{d}{dx}(my) = m \frac{dy}{dx} + \frac{dm}{dx} y$

Thus will work if $\frac{dm}{dx} = 2xm$

$$m = e^{x^2}$$

$$\mu \frac{dy}{dx} + 2x\mu y = 2x\mu \quad \mu = e^{x^2}$$

$$e^{x^2} \frac{dy}{dx} + 2xe^{x^2}y = 2xe^{x^2}$$

$$\int \frac{d}{dx}(e^{x^2}y) = \int 2xe^{x^2}$$

$$e^{x^2}y = e^{x^2} + C$$

$$y = 1 + Ce^{-x^2}$$

$$\int \frac{d\mu}{dx} \frac{1}{\mu} \int 2x dx$$

$$\ln |\mu| = x^2 + C$$

$$\mu = Ae^{x^2}$$

Soln of First Order ODE

Monday, April 02, 2007
8:56 AM

Soln of the general first order ODE using an integrating factor

$$(*) \frac{dy}{dx} + p(x)y = g(x)$$

① Multiply eqn (*) by integrating factor $M(x)$, a function to be determined

$$M(x) \frac{dy}{dx} + M(x)p(x)y = g(x)M(x)$$

② Choose $M(x)$ so that the left-hand side is a derivative
Specifically $\frac{d}{dx}(My) = M \frac{dy}{dx} + \frac{dM}{dx} y$

$$L = M \frac{dy}{dx} + M p y$$

Hence $\boxed{\frac{dM}{dx} = M(x)p(x)}$ $\leftarrow$ eqn for the integrating factor M

③ Solve for $M(x)$

$$\int \frac{dM}{M} = \int p(x) dx$$

$$\ln |M| = \int p(x) dx$$

$$M = e^{\int p(x) dx}$$

No arbitrary constant of integration b/c it drops out in the end.

④ Solve

$$e^{\int p(x) dx} \frac{dy}{dx} + e^{\int p(x) dx} p(x)y = e^{\int p(x) dx} g(x) \quad \text{Multiply by integrating factor}$$

$$\Rightarrow \frac{d}{dx} [e^{\int p(x) dx} y] = e^{\int p(x) dx} g(x)$$

$$\int \frac{d}{dx} [e^{\int p(x) dx} y] dx = \int e^{\int p(x) dx} g(x) dx$$

$$e^{\int p(x) dx} y = \int e^{\int p(x) dx} g(x) dx + C$$

$$y = e^{-\int p(x) dx} \left[\int e^{\int p(x) dx} g(x) dx \right] + C e^{-\int p(x) dx}$$

Check this:


$$\frac{dy}{dx} = -p(x)y + g(x)$$

$$\frac{dy}{dx} = \frac{d}{dx} \left(e^{-\int p(x) dx} \left[\int e^{\int p(x) dx} g(x) dx \right] + C e^{-\int p(x) dx} \right)$$

$$= e^{-\int p(x) dx} (-p(x)) \left[\int e^{\int p(x) dx} g(x) dx \right] + e^{-\int p(x) dx} \left[e^{\int p(x) dx} g(x) \right]$$

$$+ \dots - \int p(x) dx \dots$$

$$+ Ce^{-\int p(x) dx} (-p(x)) \quad \overbrace{g(x)} \\ = -p(x) \underbrace{\left[e^{-\int p(x) dx} \int e^{\int p(x) dx} g(x) dx + Ce^{-\int p(x) dx} \right]}_{y \text{ (what we started with)}}$$



Initially: 60 gallons pure H_2O in the tank

Find amt of saltwater in tank at time t

Find amt of salt in tank at time t

What is max amt salt in tank?

Concentration of salt in the liquid = $\frac{s}{v}$

We want diff eqns for $v(t)$ & $s(t)$ $\frac{dv}{dt} = ?$ $\frac{ds}{dt} = ?$ $v(0) = 60 \text{ gal}$
 $s(0) = 0 \text{ salt lbs}$

What is Δv in time Δt ?

$$\Delta v = (2 \text{ gal/min}) \Delta t - (3 \text{ gal/min}) \Delta t = (-1 \text{ gal/min}) \Delta t$$

$$t > 60 \Rightarrow v(t) = 0 \\ s(t) = 0$$

$$\frac{dv}{dt} = -1 \Rightarrow \boxed{\frac{dv}{dt} = -1, v(0) = 60}$$

What is Δs at Δt ?

$$\Delta s = (1 \text{ lb/gal}) (2 \text{ gal/min}) \Delta t - \left(\frac{s}{v} \right) \left(\frac{3 \text{ gal}}{\text{min}} \right) \Delta t$$

$$\frac{ds}{dt} = 2 - \frac{3s}{v} \Rightarrow \boxed{\frac{ds}{dt} = 2 - \frac{3s}{v}, s(0) = 0}$$

$$\int \frac{dv}{dt} dt = \int -1 dt$$

$$v(t) = -t + C$$

$$v(0) = 0 + C$$

$$60 = C$$

$$v(t) = 60 - t \quad 0 \leq t \leq 60$$

$$\frac{ds}{dt} = 2 - \frac{3s}{v} = 2 - \frac{3s}{60-t}$$

linear 1st order, but not separable

$$\frac{ds}{dt} + \frac{3s}{60-t} = 2$$

$$p(x) = \frac{3}{60-t} \quad g(x) = 2$$

$$\frac{dy}{dx} + p(x)y = q(x)$$

Solve by integrating factors


$$\mu = e^{\int \frac{3}{60-t} dt} = -3 \ln|60-t| = e^{-3 \ln|60-t|}$$

$$= (e^{\ln|60-t|})^{-3}$$

$$= (60-t)^{-3}$$

$$\mu = \frac{1}{(60-t)^3}$$

Salt water tank Prob

2 gal/min
1 lb/gal
salt

$V(t)$ = salt water
in gal
in tank
after
 t min

$s(t)$ = salt in
lb in
tank after
 t min

$$V(0) = 60$$

$$s(0) = 0$$

$$\frac{dV}{dt} = -1 ; V(0) = 60 \quad 0 \leq t \leq 60$$

$$\frac{ds}{dt} = 2 - \frac{3s}{V} ; s(0) = 0$$

$$V(t) = 60 - t \quad \frac{ds}{dt} = 2 - \frac{3}{60-t} s, \quad s(0) = 0$$

$$\frac{ds}{dt} + \left(\frac{3}{60-t}\right)s = 2 \quad \text{Solve w/ an integrating factor}$$

$$\mu = e^{\int \frac{3}{60-t} dt} = e^{-3 \ln|60-t|} = \left(\frac{1}{60-t}\right)^3$$

$$\left(\frac{1}{60-t}\right)^3 \frac{ds}{dt} + \left(\frac{3}{60-t}\right)^4 s = \frac{2}{(60-t)^3}$$

$$\int \frac{d}{dt} \left(\frac{1}{(60-t)^3} s \right) dt = \int \frac{2}{(60-t)^3} dt$$

$$\frac{1}{(60-t)^3} s = \int \frac{2}{(60-t)^3} dt \quad \begin{matrix} v = 60-t \\ dv = -dt \end{matrix}$$

$$\frac{s}{(60-t)^3} = \frac{1}{(60-t)^2} + C$$

$$t=0, s=0 \quad 0 = \frac{1}{60^2} + C$$

$$\Rightarrow \quad -\frac{1}{3600}$$

$$s(t) = (60-t)^3 \left[\frac{1}{(60-t)^2} - \frac{1}{60^2} \right]$$

$$s(t) = (60-t) - \frac{(60-t)^3}{60^2}$$

Max amt of salt, $s_{\max}$ occurs at time $t = t_{\max}$
At $t = t_{\max}$, $\frac{ds}{dt} = 0$

$$\frac{ds}{dt} = -1 + \frac{3(60-t)^2}{60^2}$$

$$0 = -1 + \frac{3(60-t)^2}{60^2}$$

$$\sqrt{1200} = \sqrt{(60-t)^2}$$

$$\frac{60}{\sqrt{3}} = 60 - t$$

$$60 - \frac{60}{\sqrt{3}} = t_{\max}$$

$$2 S_{\min} \approx t_{\max}$$

$$\begin{aligned} S_{\max} &= S(t_{\max}) = 60 - 60 + \frac{60}{\sqrt{3}} - \left(\left(\frac{60}{\sqrt{3}} \right)^3 \right) \\ &= \frac{60}{\sqrt{3}} \left(1 - \frac{1}{3} \right) \\ &= \frac{40}{\sqrt{3}} \approx 23.16 \end{aligned}$$

Population growth models

Experimental Growth (too simplistic)

Let $P(t)$ = size of pop at time t . Continuous variable
(treat^{as} continuous real variable)

Let ΔP = change in pop size in amt time Δt
= # births in time Δt - # deaths in time Δt

Assume birth rate $\propto P + \Delta t$ $\Delta P_{\text{birth}} = a P \Delta t$
death rate $\propto P + \Delta t$ $\Delta P_{\text{death}} = -b P \Delta t$

$$\Delta P = \underbrace{(a-b)}_K P \Delta t \Rightarrow \frac{\Delta P}{\Delta t} = K P \rightarrow \lim_{\Delta t \rightarrow 0} \frac{\Delta P}{\Delta t} = \frac{dP}{dt} = K P$$

$$P(0) = P_0$$

$$P(t) = C e^{kt}$$

$$P(0) = P_0 = C$$


$P(t) = P_0 e^{kt}$ if k is pos, exponential divergence in population

99% of the ppl you meet in college

Logistic Model Population growth

Assume birth and death rates (a & b) depend linearly on pop size


$$\begin{aligned} a &\rightarrow a_0 - a_1 P & a_0, a_1 > 0 \\ b &\rightarrow b_0 + b_1 P \end{aligned}$$



$$\begin{aligned} \frac{dP}{dt} &= aP - bP = (a_0 - a_1 P)P - (b_0 + b_1 P)P \\ &= (a_0 - b_0)P - (a_1 + b_1)P^2 \\ &= \underbrace{(a_1 + b_1)}_{K > 0} P \left[\underbrace{\frac{a_0 - b_0}{a_1 + b_1}}_{M > 0} - P \right] \end{aligned}$$

$$\frac{dP}{dt} = K P (M - P)$$

$$\frac{dP}{dt} \begin{cases} > 0 & \text{if } P < M \\ < 0 & \text{if } P > M \end{cases} \quad M = \text{carrying}$$



Logistic eqn for pop

$$\frac{dP}{dt} = K(M-P)P$$

$$K, M > 0 \quad (\text{pos constants}) \quad P(0) = P_0$$

$M = \text{Carrying capacity}$

Nondimensionalize $\hat{P} = \frac{P}{M}$

$$\tau = MKt \quad \text{dimensionless time}$$

$$\frac{dP}{dt} = \frac{d}{dt}(M\hat{P}) = M \frac{d\hat{P}}{dt}$$

$$M \frac{d\hat{P}}{d\tau} \frac{d\tau}{dt} = M^2 K \frac{d\hat{P}}{d\tau}$$

$$K(M-P)P = M^2 K \frac{d\hat{P}}{d\tau} \quad P = M\hat{P}$$

$$K(M-M\hat{P})M\hat{P} = M^2 K \frac{d\hat{P}}{d\tau} \quad t = \frac{\tau}{MK}$$


$$K M^2 (1-\hat{P})\hat{P} = M^2 K \frac{d\hat{P}}{d\tau}$$

$$\boxed{\hat{P} - \hat{P}^2 = \frac{d\hat{P}}{d\tau}}$$

$$P(0) = P_0 = M\hat{P}_0$$

$$P(t) = P_0 e^{Kt}$$

$$\frac{d\hat{P}}{d\tau} = (1-\hat{P})\hat{P}$$



Population can't be neg,
 so only pos. $\hat{P}$.

$$\frac{d\hat{P}}{d\tau} > 0 \quad \frac{d\hat{P}}{d\tau} < 0$$

$\hat{P} = 0$ & 1 are equilib solns (singular) regions

expect $0 < \hat{P}_0 < 1 \Rightarrow$ soln increases toward $\hat{P} = 1$
 $\hat{P}_0 > 1 \Rightarrow$ soln decreases towards $\hat{P} = 1$

Solve $\frac{d\hat{P}}{d\tau} = (1-\hat{P})\hat{P}$; $\hat{P}(0) = P_0 \neq 0, 1$

$$\int \frac{d\hat{P}}{(1-\hat{P})\hat{P}} = \int d\tau = \tau + C$$

$$\int \frac{d\hat{P}}{(1-\hat{P})\hat{P}}$$

$$\int \frac{1}{1-\hat{p}} + \frac{1}{\hat{p}} d\hat{p} = \frac{1}{(1-\hat{p})\hat{p}} = \frac{A}{(1-\hat{p})} + \frac{B}{\hat{p}}$$

$$= \ln|1-\hat{p}| + \ln|\hat{p}| = \ln\left|\frac{\hat{p}}{1-\hat{p}}\right|$$

$$= \frac{A\hat{p} + B(1-\hat{p})}{(1-\hat{p})\hat{p}}$$

$$A\hat{p} + B(1-\hat{p}) = 1$$

$$\ln\left|\frac{\hat{p}}{1-\hat{p}}\right| = \tau + C \Rightarrow e^{\ln\left|\frac{\hat{p}}{1-\hat{p}}\right|} = e^{\tau+C}$$

$$= e^{\tau} e^C$$

$$\hat{p} = 0, B=1$$

$$\hat{p}=1, A=1$$

$$\left|\frac{\hat{p}}{1-\hat{p}}\right| = e^C e^{\tau}$$

$$\hat{p} = 0, B=1$$

$$\hat{p}=1, A=1$$

$$B + \hat{p}(A-B) = 1$$

$$\frac{\hat{p}}{1-\hat{p}} = K e^{\tau}; \quad \tau=0 \Rightarrow \hat{p} = \hat{p}_0 \Rightarrow \frac{\hat{p}_0}{1-\hat{p}_0} = K$$

$$\frac{\hat{p}}{1-\hat{p}} = \frac{\hat{p}_0}{1-\hat{p}_0} e^{\tau} \Rightarrow \hat{p} = \frac{\hat{p}_0}{1-\hat{p}_0} e^{\tau} (1-\hat{p})$$

$$\hat{p} \left[1 + \frac{\hat{p}_0 e^{\tau}}{1-\hat{p}_0} \right] = \frac{\hat{p}_0}{1-\hat{p}_0} e^{\tau}$$

$$\hat{p}(\tau) = \frac{\hat{p}_0}{(1-\hat{p}_0) \left[1 + \frac{\hat{p}_0 e^{\tau}}{1-\hat{p}_0} \right]} = \frac{\hat{p}_0 e^{\tau}}{1-\hat{p}_0 + \hat{p}_0 e^{\tau}} = \frac{\hat{p}_0}{(1-\hat{p}_0) e^{-\tau} + \hat{p}_0}$$

$$\hat{p}(\tau) = \frac{\hat{p}_0}{(1-\hat{p}_0) e^{-\tau} + \hat{p}_0}$$

Back to original dimensionless variable

$$p(t) = M \hat{p}(MKt)$$

$$\uparrow$$

$$\tau$$

$$p_0 = M \hat{p}_0$$

$$p(t) = \frac{M \hat{p}_0}{(1-\hat{p}_0) e^{-MKt} + \hat{p}_0} = \frac{p_0}{(1-\frac{p_0}{M}) e^{-MKt} + \frac{p_0}{M}}$$

$$p(t) = \frac{M p_0}{(M-p_0) e^{-MKt} + p_0}$$

Check behavior of soln as $t \rightarrow \infty$

$$\lim_{t \rightarrow \infty} m p = \frac{M p_0}{p_0} = M$$

$$\ddot{x} = 0 \quad \overline{f_0} = 11$$


Qualitative Analysis of the following

$$\frac{dx}{dt} = f(x)$$

Autonomous 1st order DEQ

Autonomous means $\frac{dx}{dt}$ depends on x but not on t

Non Autonomous $\frac{dx}{dt} = f(x, t)$



$$x = x_j; f(x_j) = 0 \Leftarrow \text{equilib}$$

x_1, x_3, x_5 are unstable equilibria

x_2, x_4, x_6 are stable

$$x = x_j \text{ is stable eq if } f(x_j) = 0 \text{ \& } \frac{df}{dx}(x_j) < 0$$


$$\text{unstable if } \frac{df}{dx}(x_j) > 0$$

Linear stability of equilib solns

Autonomous First order Ordinary Diff eqns

$$\frac{dx}{dt} = f(x)$$

Qualitative Analysis




c_1, c_2 are equilib solns,
or critical pts

So when $f(x) = y = \frac{dx}{dt} = 0$,

Slope is 0.

$$f(c_1) = f(c_2) = 0$$



Phase diagram



$x = c_1$ is an unstable equilib

$x = c_2$ is a stable equilib

Wants to return to starting pt



To infinity &
beyond!

An equilib $x=c$ is stable provided that initial condition $x=x_0$ is sufficiently close to $x=c$, then $x(t)$ remains close to c for all $t \geq 0$

If condition is not met, it's unstable.

Sufficient conditions for stability/instability

Assume $x=c$ is an equilib soln. $f(c)=0$

Now consider initial condition $x_0 = c + \hat{x}$ $0 < |\hat{x}| < 1$

$$\frac{dx}{dt} = f(x) \quad f(c)=0 \quad x_0 = c + \hat{x}$$

Approximate soln

If $\hat{x}(t)$ remains small then


$$x(t) = c + \tilde{x}(t)$$

c is stable

$$\text{LHS } \frac{dx}{dt} = \frac{d}{dt}(c + \tilde{x}) = \frac{d\tilde{x}}{dt}$$

$$\text{RHS } f(x) = f(c + \tilde{x})$$

Approximate $f(x)$ by
a straight line slope
 $\frac{df}{dx}(c)$





$$y = f(x) \approx f'(c)(x-c) \quad \frac{y-0}{x-c} = f'(c)$$

$$= f'(c) \tilde{x}$$

$$\frac{d\tilde{x}}{dt} \approx f'(c) \tilde{x}, \quad \tilde{x}_0 = \tilde{x}(0) \Rightarrow \boxed{\tilde{x} e^{f'(c)t}}$$

Sufficient condition for stability of $x=c$
 $f'(c) < 0$

Sufficient condition for instability of $x=c$
 $f'(c) > 0$



doesn't work
for $f'(c)=0$

Taylor's Thm

Suppose the $(n+1)^{\text{th}}$ derivative of $f(x)$ exists on an interval containing a, b . Then $f(b) = f(a)$

$$f(b) = f(a) + f'(a)(b-a) + \frac{f''(a)}{2}(b-a)^2 + \frac{f'''(a)}{3!}(b-a)^3$$

$$+ \dots + \frac{f^{(n)}(a)}{n!}(b-a)^n + \frac{f^{(n+1)}(z)}{(n+1)!}(b-a)^{n+1}$$

z is between a & b

ex Expand $f(x) = \frac{1}{x}$ about $x=1$

$$f(x) = \frac{1}{x} \approx f(1) + f'(1)(x-1) + \frac{f''(1)}{2}(x-1)^2 + \frac{f'''(1)}{3!}(x-1)^3$$

$$+ \dots + \frac{f^{(n)}(1)}{n!}(x-1)^n$$

$$f(x) = \frac{1}{x} \quad f'(x) = -\frac{1}{x^2} \quad f''(x) = \frac{2}{x^3} \quad f'''(x) = -\frac{6}{x^4}$$

$$f^{(4)}(x) = \frac{24}{x^5} \quad f^{(n)}(x) = (-1)^n \left(\frac{n!}{x^{n+1}} \right)$$

$$f(1) = 1$$

$$f(x) = \frac{1}{x} \approx 1 - (x-1) + (x-1)^2 - (x-1)^3 +$$

$$\begin{aligned}
 f(1) &= 1 \\
 f'(1) &= -1 \\
 f''(1) &= 2 \\
 f'''(1) &= -3! \\
 f^{(4)}(1) &= 4!
 \end{aligned}$$

$$f(x) = \frac{1}{x} \approx 1 - (x-1) + (x-1)^2 - (x-1)^3 + \underbrace{(x-1)^4 + \dots + (-1)^n (x-1)^n}_{p_n(x)}$$

$$\left| \frac{1}{x} - p_n(x) \right| = \frac{|f^{(n+1)}(z)|}{(n+1)!} |x-1|^{n+1}$$

$$|x-1|^{n+1} \frac{|f^{(n+1)}(z)|}{(n+1)!} \leq \frac{M}{(n+1)!} |x-1|^{n+1}$$


$$f^{(n+1)}(z) = (-1)^{n+1} \frac{n!}{z^{n+1}}$$

Solving 1st Order Diff Eqns

Monday, April 09, 2007
9:03 AM

Numerical Methods for solving 1st order diff eqns

$$\begin{cases} \frac{dy}{dx} = f(x, y) \\ y(x_0) = y_0 \end{cases}$$



Estimate $y = y(x)$
at discrete values
of x

$$h = \frac{x_n - x_0}{n}$$

x	y
x_0	y_0
$x_0 + h = x_1$	$y_1 = y_0 + h f(x_0, y_0)$
$x_0 + 2h = x_2$	$y_2 = y_1 + h f(x_1, y_1)$
$x_0 + nh = x_n$	$y_n = y_{n-1} + h f(x_{n-1}, y_{n-1})$

$$\frac{dy}{dx} = \frac{y_1 - y_0}{x_1 - x_0} = f(x_0, y_0)$$

$$y_1 = y_0 + h f(x_0, y_0)$$

$$y_{n+1} = y_n + h f(x_n, y_n)$$

Estimate how the error in Euler's method scales w/ h

Consider error in one step of integration

Let $y = \Phi(x)$ be the exact soln to *

$$y_{n+1} = \Phi(x_{n+1}) = \Phi(x_n + h)$$

$$\begin{aligned} \Phi(x_n) &= \frac{dy}{dx} \bigg|_{x_n} = f(x_n, y_n) \\ &= \Phi(x_n) + \Phi'(x_n)h + \frac{1}{2}\Phi''(x_n)h^2 + \dots \\ &= y_n + h f(x_n, y_n) + E \end{aligned}$$

Soln by Euler's Method + Error

Taylor's Thm allows us to estimate Error

$$E = \frac{1}{2}h^2\Phi''(\xi)$$

$$\Phi(x_{n+1}) = \Phi(x_n + h) \approx \Phi(x_n) + h\Phi'(x_n) + \frac{1}{2}h^2\Phi''(\xi)$$

ξ between x_n, x_{n+1}

$$|E_n| = Ch^2 \text{ constant}$$

So the error is some constant of h^2

$$\text{Total error} \propto Nh^2$$

integration steps

As $h \downarrow$, $N \uparrow$

$$N = \frac{x_n - x_0}{h}$$

$$TE \propto \frac{x_n - x_0}{h} h^2 \propto (x_n - x_0) h$$

$$E_n \propto h$$

error $\propto h$
if $2N$, then $\frac{1}{2}h$

EITHER/OR
IF/THEN

Demonstrate this error scaling w/ prob. we can BEFORE/AFTER

Solve.

$$\frac{dy}{dx} = y$$

$$y(0) = 1$$

$$y(1) = ?$$

Exact soln
 $\Rightarrow y(x) = Ce^x$
 $1 = C$

$$y(x) = e^x$$

$$y(1) = e = 2.7182818...$$

$$x_0 = 0$$

$$y_0 = 1$$

$$x_1 = h$$

$$y_1 = y_0 + y_0 h = (1+h)y_0 = (1+h)$$

$$x_2 = 2h$$

$$y_2 = y_1 + y_1 h = (1+h)y_1 = (1+h)^2$$

$$x_3 = 3h$$

$$y_3 = y_2 + y_2 h = (1+h)y_2 = (1+h)^3$$

$$x_n = nh \quad y_n = y_{n-1} + y_{n-1} h = (1+h)y_{n-1} = (1+h)^n$$

$$h = \frac{x_n - x_0}{n} = \frac{1-0}{n} = \frac{1}{n}$$

$$y(1) \approx (1+h)^n, \quad n = \frac{1}{h}$$

h	n	approx	Error
.1	10	$y_1 \approx 1.1^{10} = 2.59$	$2.59 - e = .124$
.01	100	$y \approx 1.01^{100} = 2.705$	$2.705 - e = .0135$
.001	1000	2.717	.00136

Improve on Euler's Method

$$\begin{cases} \frac{dy}{dx} = f(x, y) \\ y(x_0) = y_0 \end{cases} \quad \text{1 subinterval from } x_n \text{ to } x_{n+1}$$

$$\int_{x_n}^{x_n+h} \frac{dy}{dx} = \int_{x_n}^{x_n+h} f(x, y) dx$$

$$y \Big|_{x_n}^{x_n+h} = y(x_{n+1}) - y(x_n) = y_{n+1} - y_n$$

$$y_{n+1} = y_n + \int_{x_n}^{x_n+h} f(x, y) dx \quad \text{exact}$$

$$= \int_{x_n}^{x_{n+1}} f(x, y) dx$$


$$y_{n+1} = y_n + \int_{x_n}^{x_{n+1}} f(x, y) dx$$

exact

$$= \int_{x_n}^{x_{n+1}} f(x, y) dx$$

Left endpt approx

$$\int_{x_n}^{x_{n+1}} f(x, y) dx$$



$$\int_{x_n}^{x_{n+1}} f(x, y) dx \approx \int_{x_n}^{x_{n+1}} f(x_n, y_n) dx = f(x_n, y_n) h$$


Right end approx

Backward
Euler
Method


$$y_{n+1} = y_n + h f(x_{n+1}, y_{n+1})$$

implicit eqn for y_{n+1}

First Order ODE's in particular
improvements to Euler's Method



$$\frac{dy}{dx} = f(x, y) ; y(x_0) = y_0$$



$$x_{n+1} - x_n = h$$

$$N = \frac{x_N - x_0}{h}$$


$$\int_{x_n}^{x_{n+1}} \frac{dy}{dx} dx = \int_{x_n}^{x_{n+1}} f(x, y) dx$$

$$\underbrace{y(x_{n+1})}_{y_{n+1}} - \underbrace{y(x_n)}_{y_n} = \int_{x_n}^{x_{n+1}} f(x, y) dx$$

Problem / this is
we don't know
 $y(x)$

$$y_{n+1} = y_n + \int_{x_n}^{x_{n+1}} f(x, y) dx \quad \text{exact}$$


Euler's Method is a left-endpt. approximation to integral



$$y_{n+1} \approx y_n + f(x_n, y_n) h$$

Explicit formula for y_{n+1} in terms of x_n & y_n


Backward Euler Method : Right endpt. approx




$$y_{n+1} \approx y_n + f(x_{n+1}, y_{n+1}) h$$

Implicit formula for y_{n+1} in
terms of x_n, y_n

Trapezoidal Approx to integral idea behind improved Euler
method





$$y_{n+1} \approx y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_{n+1})]$$

Improved Euler $y_{n+1} \approx y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_n + hf(x_n, y_n))]$

Predictor $\rightarrow y_{n+1} = y_n + hf(x_n, y_n)$

Corrector $\rightarrow y_{n+1} = y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_{n+1})]$

Improved Euler

Error scales as h^2 , meaning that if you double N the error goes down by $\frac{1}{4}$

Example

$$\frac{dy}{dx} = y$$

$$y(0) = 1$$

$$\Rightarrow y(x) = e^x$$

$$y(1) = e$$

h	N	$(1+h)^N$	$y(1)$ Euler	error for Euler	$(1+h\frac{h^2}{2})^N$	$y(1)$ imp. Euler	error imp. Euler
$\frac{1}{5}$	2	2.25	2.25	.468	2.64	2.64	.078
$\frac{1}{4}$	4	2.441	2.441	.277	2.645	2.645	.023
$\frac{1}{8}$	8	2.566	2.566	.153	2.712	2.712	.0064

$$y_{n+1} = y_n + hf(x_n, y_n) \\ = y_n + hy_n = (1+h)y_n$$

$$y_{n+1} = y_n + \frac{h}{2} [y_n + (1+h)y_n] \\ = [1+h+\frac{h^2}{2}]y_n$$

Euler is 1st order accurate for $E \sim h$

Improved Euler is 2nd order accurate $E \sim h^2$

4th order Runge Kutta $E \sim h^4$

$$k_1 = f(x_n, y_n)$$

$$k_2 = f(x_n + \frac{h}{2}, y_n + \frac{h}{2}k_1)$$

$$k_3 = f(x_n + \frac{h}{2}, y_n + \frac{h}{2}k_2)$$

$$k_4 = f(x_n + h, y_n + hk_3)$$

$$y_{n+1} = y_n + \frac{h}{6} [k_1 + 2k_2 + 2k_3 + k_4]$$

If $f(x, y)$ depends only on x , then this is just Simpson's

approx. $\frac{dy}{dx} = f(x)$

$$y_{n+1} - y_n = \int_{x_n}^{x_{n+1}} \frac{dy}{dx} dx = \int_{x_n}^{x_{n+1}} f(x) dx = \frac{h}{6} [f(x_n) + 4f(x_n + \frac{h}{2}) + f(x_{n+1})]$$

Simpson's

$$RK4 \quad y_{n+1} - y_n = \frac{h}{6} [k_1 + 2k_2 + 2k_3 + k_4]; \begin{matrix} k_1 = f(x_n) \\ k_2 = f(x_n + \frac{h}{2}) \\ k_3 = f(x_n + \frac{h}{2}) \\ k_4 = f(x_n + h) \end{matrix}$$

$$k_3 = f(x_n + \frac{h}{2})$$

$$k_4 = f(x_{n+1})$$

4th order Runge Kutta ($E \sim h^4$)

$$\frac{dy}{dx} = f(x, y)$$

$$\begin{cases} k_1 = f(x_n, y_n) \\ k_2 = f(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hk_1) \\ k_3 = f(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hk_2) \\ k_4 = f(x_n + h, y_n + hk_3) \end{cases}$$

$$y_{n+1} = y_n + \frac{h}{6}[k_1 + 2k_2 + 2k_3 + k_4]$$

2 Special Cases

$$\frac{dy}{dx} = f(x) \Rightarrow y_{n+1} = y_n + \int_{x_n}^{x_{n+1}} f(x) dx$$

R-K is just Simpson's Approx of the integral in this case

$$\frac{dy}{dx} = f(y)$$

Show that $E \sim h^4$; We do this by showing that the error in 1 step (from y_n to y_{n+1}) is $E_n < C_n h^5$

Then the total error is $E_T < K N h^5 = K(x_n - x_0) \frac{h^4}{h}$

Idea behind proof that $E_n \sim h^5$

Approx soln: $y_{n+1} = y_n + \frac{h}{6}[k_1 + 2k_2 + 2k_3 + k_4]$

"True soln": $y(x) = \Phi(x)$

$$y_{n+1} = \Phi(x_n + h)$$

Compare Taylor Series through h^4 termsApproximate soln: Taylor Series Approx about (x_n, y_n)

$$k_1 = f(y_n)$$

$$k_2 = f(y_n + \frac{1}{2}hk_1) = f(y_n) + f'(y_n)(\frac{1}{2}hk_1) + \frac{1}{2}f''(y_n)(\frac{1}{2}hk_1)^2 + \frac{1}{3!}f'''(y_n)(\frac{1}{2}hk_1)^3 + \frac{1}{4!}f^{(4)}(y_n)(\frac{1}{2}hk_1)^4 + \dots$$

$$k_3 = f(y_n + \frac{1}{2}hk_2) = f(y_n) + f'(y_n)(\frac{1}{2}hk_2) + \frac{1}{2}f''(y_n)(\frac{1}{2}hk_2)^2 + \frac{1}{3!}f'''(y_n)(\frac{1}{2}hk_2)^3 + \frac{1}{4!}f^{(4)}(y_n)(\frac{1}{2}hk_2)^4 + \dots$$

$$k_4 = f(y_n + hk_3) = f(y_n) + f'(y_n)(hk_3) + \frac{1}{2}f''(y_n)(hk_3)^2 + \frac{1}{3!}f'''(y_n)(hk_3)^3 + \frac{1}{4!}f^{(4)}(y_n)(hk_3)^4 + \dots$$

Taylor Series expansion of approx soln $f = f(y_n)$

$$y_{n+1} = y_n + fh + \frac{1}{2}f'f h^2 + \frac{1}{6}(f''f^2 + (f')^2f)h^3 + \frac{1}{24}(4f''f'f^2 + (f')^3f + f'''f^3)h^4 + \Theta(h^5)$$

"of order h^5 "

$f, f', f'', \dots$ are evaluated at y_n

Now Taylor expand the true soln $y = \Phi(x)$ $\left[\frac{dy}{dx} = \Phi'(x) = f(y) \right]$

$$y_{n+1} = \Phi(x_{n+1}) = \Phi(x_n + h) = \Phi(x_n) + \Phi'(x_n)h + \frac{1}{2}\Phi''(x_n)h^2 + \frac{1}{3!}\Phi'''(x_n)h^3 + \frac{1}{4!}\Phi^{(4)}(x_n)h^4 + \dots$$

$$= y_n + f(y_n)h +$$

$$\begin{aligned} \Phi''(x_n) &= \frac{d}{dx}(\Phi'(x)) \Big|_{x=x_n} \\ &= \frac{d}{dx}(f(y(x))) \Big|_{x=x_n} \\ &= \underbrace{f'(y(x))}_{f'(y_n)} \underbrace{y'(x)}_{f(y_n)} \Big|_{x=x_n} = f'(y_n)f(y_n) \quad \checkmark \end{aligned}$$

$$\Phi'''(x_n) = \frac{d}{dx}(\Phi'') = \frac{d}{dx}(f'(y_n)f(y_n)) = \frac{d}{dx} \left[\underbrace{f'(y_n)}_{f'(y_n)} \underbrace{f(y_n)}_{f(y_n)} \right]$$

$$= f''(y_n) \underbrace{\frac{dy}{dx} \Big|_{x=x_n}}_{f(y_n)} f(y_n) + f'(y_n) f'(y_n) \frac{dy}{dx}$$

$$\Phi'''(x_n) = f''f^2 + (f')^2f \quad \checkmark$$

$$\Phi^{(4)}(x_n) = \frac{d}{dx}(\Phi''') \Big|_{x=x_n} = \frac{d}{dx} [f''f^2 + (f')^2f]$$

$$\frac{dy}{dx} = f(x, y)$$

$$y(x_0) = y_0$$

$$x_n = x_0 + nh$$

$$y_n = y(x_n)$$

$$h = x_{n+1} - x_n$$

Euler $y_{n+1} = y_n + hf(x_n, y_n)$

Improved Euler:

$$u_{n+1} = y_n + hf(x_n, y_n)$$

$$y_{n+1} = y_n + \frac{h}{2} (f(x_n, y_n) + f(x_{n+1}, u_{n+1}))$$

Runge Kutta

$$k_1 = f(x_n, y_n)$$

$$k_2 = f(x_n + \frac{h}{2}, y_n + \frac{h}{2} k_1)$$

$$k_3 = f(x_n + \frac{h}{2}, y_n + \frac{h}{2} k_2)$$

$$k_4 = f(x_n + h, y_n + k_3)$$

$$y_{n+1} = y_n + \frac{h}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$

$$\frac{dy}{dx} = y \quad y(0) = 1 \Rightarrow y = e^x$$

$$y_0 = 1$$

$$y_1 = y(x_0 + h) = y(h) = e^h \quad \text{exact}$$

$$= \sum_{n=0}^{\infty} \frac{h^n}{n!} = 1 + h + \frac{1}{2}h^2 + \frac{1}{3!}h^3$$

Euler $y_1 = y_0 + hf(x_0, y_0)$
 $f(x, y) = y$ for this example

$$x_0 = 0$$

$$y_0 = 1$$

$$y_1 = 1 + h(y_0) = 1 + h$$

Error in Euler is $\Theta(h^2)$ "order h^2 " means
 leading term is Ch^2



Improved Euler

$$y_1 = y_0 + h \underbrace{f(x_0, y_0)}_{y_0} = (1+h) \underbrace{y_0}_1 = 1+h$$


$$y_1 = y_0 + \frac{h}{2} (\underbrace{f(x_0, y_0)}_{y_0=1} + \underbrace{f(x_{n+1}, y_{n+1})}_{y_1=1+h})$$

$$y_1 = 1 + \frac{h}{2} (1+h)$$

$$= 1 + h + \frac{h^2}{2}$$

Error is $\mathcal{O}(h^3)$

ow! Improved Euler!



Runge Kutta

$$k_1 = f(x_0, y_0) = y_0 = 1$$

$$k_2 = f(x_0 + \frac{h}{2}, y_0 + \frac{h}{2} k_1) = y_0 + \frac{h}{2} k_1 = 1 + \frac{h}{2}$$

$$k_3 = f(x_0 + \frac{h}{2}, y_0 + \frac{h}{2} k_2) = y_1 + \frac{h}{2} k_2 = 1 + \frac{h}{2} (1 + \frac{h}{2})$$

$$= 1 + \frac{h}{2} + \frac{h^2}{4}$$

$$k_4 = f(x_0 + h, y_0 + h k_3) = y_0 + h k_3$$

$$= 1 + h (1 + \frac{h}{2} + \frac{h^2}{4})$$

$$= 1 + h + \frac{h^2}{2} + \frac{h^3}{4}$$

$$y_1 = y_0 + \frac{h}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$

$$y_1 = 1 + \frac{h}{6} (1 + 2 + h + 2 + h + \frac{h^2}{2} + 1 + h + \frac{h^2}{2} + \frac{h^3}{4})$$

$$y_1 = 1 + h + \frac{h^2}{2} + \frac{h^3}{6} + \frac{h^4}{24}$$

Error is $\mathcal{O}(h^5)$



ow! Runge Kutta!

Ode45 is an adaptive Runge Kutta

Chapter 3 Second order diff eqns

$$\frac{d^2 y}{dx^2} = f(x, y, \frac{dy}{dx})$$

These eqns show up all the time in mechanics

These eqns show up all the time in mechanics

$$F = ma$$

$$F = m \frac{d^2x}{dt^2}$$

$$F(x, t, v) = F(x, t, \frac{dx}{dt})$$

$$\frac{d^2y}{dx^2} = F(x, y, \frac{dy}{dx})$$

A necessary condition for a unique soln is to have 2 initial conditions eg $y(x_0) = y_0$ $\frac{dy}{dx}(x_0) = y_1$

First examples


$$m \frac{d^2x}{dt^2} = F(x)$$

These conserve energy
so energy conservation
can be used as an initial
condition

example mass on a spring obeying Hooke's law



x_0 equilib position



$k = \text{spring constant}, > 0$

$$\frac{d}{dt} \left[ma = m \frac{d^2x}{dt^2} = -kx \right]$$

$$m \frac{d^2x}{dt^2} \frac{dx}{dt} = -kx \frac{dx}{dt}$$

$$m \frac{d}{dt} \left(\frac{1}{2} \left(\frac{dx}{dt} \right)^2 \right) = -k \frac{d}{dt} \left(\frac{x^2}{2} \right) = -k \left(\frac{dx}{dt} \right) \frac{x}{2}$$

$$\frac{d}{dt} \left[\frac{m}{2} \left(\frac{dx}{dt} \right)^2 + \frac{k}{2} x^2 \right] = 0$$

energy

Conservation of energy eqn

$$\frac{m}{2} \left(\frac{dx}{dt} \right)^2 + \frac{k}{2} x^2 = E_0$$

$$\left(\frac{dx}{dt} \right)^2 = \frac{2}{m} \left(E_0 - \frac{k}{2} x^2 \right)$$

$$\frac{dx}{dt} = \pm \sqrt{\frac{2}{m} \left(E_0 - \frac{k}{2} x^2 \right)}$$


$$\int \frac{dx}{\sqrt{\frac{2}{m} \left(E_0 - \frac{k}{2} x^2 \right)}} = \pm t + C$$

$$\int \sqrt{\frac{2E_0}{m} - \frac{k}{m}x^2} = \pm t + C$$

$$m \frac{d^2x}{dt^2} = F(x)$$

express $F(x) = -\frac{dV}{dx}$

RETICULAN
IRON DEPLETION



$V(x)$ = potential energy = $-\int F(x) dx$ = Work done by force

$$\int_{t_0}^{t_f} m \frac{d^2x}{dt^2} \frac{dx}{dt} dt = \int_{t_0}^{t_f} -\frac{dV}{dx} \frac{dx}{dt} dt$$

$$\int_{t_0}^{t_f} \frac{m}{2} \frac{d}{dt} \left(\left(\frac{dx}{dt} \right)^2 \right) dt = - \int_{x_0}^{x_f} \frac{dV}{dx} dx$$

$$\left. \frac{m}{2} \left(\frac{dx}{dt} \right)^2 \right|_{t_0}^{t_f} = - \left. V(x) \right|_{x_0}^{x_f}$$

$$\left. \frac{m}{2} \left(\frac{dx}{dt} \right)^2 \right|_{t=t_f} - \frac{m}{2} v_0^2 = V(x_f) - V(x_0)$$

$$m \frac{d^2x}{dt^2} = F(x) = -\frac{dV}{dx}; V(x) = -\int F(x) dx$$


$$x(t_0) = x_0; \left. \frac{dx}{dt} \right|_{t=t_0} = v_0$$

$$\frac{m}{2} \left(\frac{dx}{dt} \right)^2 + V(x) = E_0 \leftarrow \text{Initial energy}$$

$$\frac{m}{2} v_0^2 + V(x_0) =$$

$$\frac{dx}{dt} = \pm \sqrt{\frac{2}{m}(E_0 - V(x))} \quad \text{1st order DEQ, can solve}$$

Ex Mass on a spring obeying Hooke's Law



$x=0$ at equilib

$$m \frac{d^2x}{dt^2} = -kx$$

$$x(0) = x_0$$

$$v(0) = v_0 = \left. \frac{dx}{dt} \right|_{t=0}$$

$$V(x) = -\int F(x) dx = k \int x dx = \frac{k}{2} x^2$$

$$\frac{1}{2} m \left(\frac{dx}{dt} \right)^2 + \frac{k}{2} x^2 = E_0$$

$$\frac{1}{2} m v_0^2 + \frac{1}{2} k x_0^2$$

$$\frac{dx}{dt} = \pm \sqrt{\frac{2}{m} (E_0 - k \frac{x^2}{2})}$$

$$\int \frac{1}{\sqrt{\frac{2E_0}{m} - \frac{k}{m} x^2}} dx = \int dt = \pm t + C$$

$$\sqrt{\frac{2E_0}{m} - \frac{k}{m} x^2} = \sqrt{\frac{2E_0}{m}} \sqrt{1 - \frac{k}{2E_0} x^2}$$

$$x = \sqrt{\frac{2E_0}{k}} \sin \phi$$

$$dx = \sqrt{\frac{2E_0}{k}} \cos \phi d\phi$$

$$\sqrt{\frac{m}{2E_0}} \int \frac{1}{\sqrt{1 - \frac{k}{2E_0} x^2}} dx = \pm t + C$$

$$1 - \frac{k}{2E_0} x^2 = \sqrt{1 - \frac{k}{2E_0} \frac{2E_0}{k} \sin^2 \phi} = \sqrt{\cos^2 \phi} = \pm \cos \phi$$

$$\sqrt{\frac{m}{2E_0}} \int \frac{\pm 1}{\cos \phi} \sqrt{\frac{2E_0}{k}} \cos \phi d\phi$$

$$\sqrt{\frac{m}{k}} \int \pm d\phi = \pm t + C$$

$$\int d\phi = \pm \sqrt{\frac{k}{m}} t + \tilde{K}$$


$$\phi = \pm \sqrt{\frac{k}{m}} t + \tilde{K}$$

$$x(t) = \sqrt{\frac{2E_0}{k}} \sin\left(\pm \sqrt{\frac{k}{m}} t + \tilde{K}\right)$$

$$x(t) = \sqrt{\frac{2E_0}{k}} \sin\left(\pm \sqrt{\frac{k}{m}} t + \tilde{K}\right)$$

$$= \pm \sqrt{\frac{2E_0}{k}} \sin\left(\sqrt{\frac{k}{m}} t \pm \tilde{K}\right)$$

$$= \sqrt{\frac{2E_0}{k}} \sin\left(\sqrt{\frac{k}{m}} t + \theta_0\right)$$




$$\text{if } (+), \tilde{K} = \theta_0, \text{ if } (-) \tilde{K} = -\theta_0$$

$$E_0 = \frac{1}{2} m v_0^2 + \frac{1}{2} k x_0^2$$

$$v(0) = x_0 = \sqrt{\frac{2E_0}{k}} \sin(\theta_0)$$

$$\theta_0 = \sin^{-1}\left(x_0 \sqrt{\frac{m}{2E_0}}\right)$$



Amplitude of oscillation
↑ w/ Energy.
larger oscillations
if E is ↑

$$T = \frac{2\pi}{\omega} = \frac{2\pi}{\sqrt{\frac{k}{m}}} = \sqrt{\frac{m}{k}} 2\pi$$

period ↑ w/ mass ↑
 ↓ w/ k

Linear 2nd order Eqns

$$y'' + p(x)y' + q(x)y = f(x)$$

Special case where $f(x)=0$, then eqn is homogeneous & linear

If $f(x) \neq 0$, then eqn is linear, inhomogeneous

Linear, homogeneous eqns have a superposition principle

If $y_1(x)$ & $y_2(x)$ solve the eqn (*), then $y(x) = c_1 y_1(x) + c_2 y_2(x)$
 (*) $y'' + p(x)y' + q(x)y = 0$ also solves the eqn for any ^{real} constants c_1 & c_2

$$\begin{aligned} y' &= c_1 y_1' + c_2 y_2' \\ y'' &= c_1 y_1'' + c_2 y_2'' \\ 0 &= c_1 y_1'' + c_2 y_2'' + p(x)[c_1 y_1' + c_2 y_2'] + q(x)(c_1 y_1 + c_2 y_2) \end{aligned}$$

$$c_1 \underbrace{(y_1'' + p(x)y_1' + q(x)y_1)}_0 + c_2 \underbrace{(y_2'' + p(x)y_2' + q(x)y_2)}_0$$

b/c y_1, y_2 are solns of the eqn.

$$0 = 0$$

The general soln will involve 2 constants. In this case, the constants are c_1 & c_2 .

$$y(x) = c_1 y_1(x) + c_2 y_2(x)$$

If you have initial conditions that $y(x_0) = y_0$ & $y'(x_0) = y_1$

$$\Rightarrow c_1 y_1(x_0) + c_2 y_2(x_0) = y_0$$

$$c_1 y_1'(x_0) + c_2 y_2'(x_0) = y_1$$

Examples

i) $y'' + y = 0$

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

General soln - $y(x) = c_1 \cos(x) + c_2 \sin(x)$

$$y_3 = 0 \quad \times$$

$$y_1 = \sin(x - \frac{\pi}{4}) = -\cos(x)\sin(\frac{\pi}{4}) + \cos(\frac{\pi}{4})\sin(x)$$

$$y_2 = \cos(x + \frac{\pi}{4}) = \cos(x)\cos(\frac{\pi}{4}) - \sin(x)\sin(\frac{\pi}{4})$$

$$y_2 = -y_1 \quad \times$$

Easily check if solns are independent or not

We find 2 solns. $y = y_1(x)$ & $y = y_2(x)$

Want to check if they are independent or not

If they are independent then we've found the general soln

$$y(x) = c_1 y_1(x) + c_2 y_2(x)$$

We obtain a criterion for independence by considering an initial value problem

$$y(x_0) = A \quad y'(x_0) = B$$

$$\begin{aligned} y \text{ at } x_0 &= c_1 y_1(x_0) + c_2 y_2(x_0) = A \\ y'(x_0) &= c_1 y_1'(x_0) + c_2 y_2'(x_0) = B \end{aligned} \quad \begin{array}{l} \text{Solve for} \\ \text{unknowns} \end{array}$$

$$\begin{bmatrix} y_1(x_0) & y_2(x_0) \\ y_1'(x_0) & y_2'(x_0) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} A \\ B \end{bmatrix} \quad \text{2 eqns, 2 unknowns } c_1, c_2$$

This can be solved, no matter what A & B are if

$$\text{Det} \begin{bmatrix} y_1(x_0) & y_2(x_0) \\ y_1'(x_0) & y_2'(x_0) \end{bmatrix} \neq 0$$

$$\begin{bmatrix} m_1 & m_2 \\ m_3 & m_4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$m_3 (m_1 x + m_2 y = a)$$

$$-m_1 (m_3 x + m_4 y = b)$$

$$m_3 m_2 y - m_1 m_4 y = m_3 a - m_1 b$$

$$y = \frac{m_3 a - m_1 b}{m_3 m_2 - m_1 m_4}$$

$$\text{So } m_3 m_2 - m_1 m_4 \neq 0$$

Wronskian

$$W(x) = \text{Det} \begin{bmatrix} y_1(x) & y_2(x) \\ y_1'(x) & y_2'(x) \end{bmatrix}$$

Det.

$$= y_1(x)y_2'(x) - y_1'(x)y_2(x)$$

$W=0 \Rightarrow$ solns not independent

$$y'' + y = 0$$

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

$$y_3 = 0$$

$$y_4 = \sin(x - \frac{\pi}{4})$$

$$y_5 = \cos(x + \frac{\pi}{4})$$

$$\{\cos(x), 0\}$$

$$W = \det \begin{bmatrix} \cos(x) & 0 \\ \sin(x) & 0 \end{bmatrix}$$

$$W=0$$

So that doesn't work

$$\{\cos(x), \sin(x)\}$$

$$W = \det \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix} = \cos^2(x) + \sin^2(x) = 1$$

$$\{\sin(x - \frac{\pi}{4}), \cos(x + \frac{\pi}{4})\}$$

$$W = \det \begin{bmatrix} \sin(x - \frac{\pi}{4}) & \cos(x + \frac{\pi}{4}) \\ \cos(x - \frac{\pi}{4}) & -\sin(x + \frac{\pi}{4}) \end{bmatrix}$$

$$= -\sin(x - \frac{\pi}{4})\sin(x + \frac{\pi}{4}) - \cos(x - \frac{\pi}{4})\cos(x + \frac{\pi}{4})$$

X

$$= -\cos(x - \frac{\pi}{4} - x - \frac{\pi}{4})$$

$$\cos(\frac{\pi}{2}) = 0$$

If the Wronskian $= 0$ at any pt it $= 0$ everywhere

The following statements are equivalent

- 1) y_1 & y_2 form a fundamental set of solns (ie the gen soln)
- 2) y_1 & y_2 are linearly independent. $y_2 \neq c y_1$ is $c_1 y_1 + c_2 y_2$
- 3) Wronskian $W(y_1, y_2) = y_1 y_2' - y_1' y_2 \neq 0$ at one pt $x = x_0$.

$$4) W(y_1, y_2) \neq 0 \text{ for all } x$$

Linear 2nd order Homogenous ODE's

Wednesday, April 18, 2007
8:58 AM

Recap

Linear, 2nd order homogeneous ODE's

$$(*) y'' + p(x)y' + q(x)y = 0$$

There is a superposition principle. If $y_1(x)$ & $y_2(x)$ solve $(*)$ then so does $c_1 y_1(x) + c_2 y_2(x)$ for any constants c_1, c_2

Need 2 linearly ind. solns y_1 & y_2 to get the general soln $y = c_1 y_1 + c_2 y_2$;

- linearly independent means $y_2 \neq C y_1$ for any constant C

- y_1 & y_2 are a "fundamental set"

Easy check of linear independence

$$\text{Compute Wronskian } W(y_1, y_2) = \det \begin{bmatrix} y_1(x) & y_2(x) \\ y_1'(x) & y_2'(x) \end{bmatrix}$$

If $W(y_1, y_2) = 0$ then y_1 & y_2 are lin. dependent

If $W(y_1, y_2) \neq 0$ then y_1 & y_2 are lin. independent

It's enough to check W at one pt. If $W(y_1(x_0), y_2(x_0)) \neq 0$ then solns are lin. independent

If $W(y_1(x_0), y_2(x_0)) = 0$ then $W(y_1(x), y_2(x)) = 0$

Special case where p, q are constants

$$a y'' + b y' + c y = 0$$

$$a \neq 0$$

a, b, c are constants

linear 2nd order homogeneous constant coefficient eqn

Method: try $y = y_0 e^{rx}$

$$y' = r y_0 e^{rx} = r y$$

$$y'' = r^2 y_0 e^{rx} = r^2 y$$

$$a y'' + b y' + c y = 0$$

$$a r^2 y + b r y + c y = 0$$

$\Rightarrow$ Need r to satisfy $\boxed{a r^2 + b r + c = 0}$

"




"Characteristic eqn"

$$y = y_0 e^{ry}$$

$$ar^2 + br + c = 0$$

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



① $b^2 - 4ac > 0$ r_1, r_2 are real
 $y = C_1 e^{r_1 x} + C_2 e^{r_2 x}$ 2 solns, distinct + real $\Rightarrow r_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, r_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$

② $b^2 - 4ac = 0$

one soln, $r_1 = -\frac{b}{2a}$ $\} y_1 = e^{r_1 x}; y_2 = ?$

③ $b^2 - 4ac < 0$

2 complex solns $r_1 = -\frac{b}{2a} + i \frac{\sqrt{4ac - b^2}}{2a}$

$$r_2 = -\frac{b}{2a} - i \frac{\sqrt{4ac - b^2}}{2a}$$

Case 1 2 solns

Ex 1

$$2y'' - 3y' + y = 0$$

$$2r^2 - 3r + 1 = 0$$

$$(2r-1)(r-1) = 0$$

$$r = \frac{1}{2}, r = 1$$

$$y(x) = C_1 e^{\frac{x}{2}} + C_2 e^x$$

$$y = e^{rx}$$

Check that $y_1 = e^{r_1 x}$ $y_2 = e^{r_2 x}$ are lin ind

$$W(y_1, y_2) = \det \begin{bmatrix} e^{r_1 x} & e^{r_2 x} \\ r_1 e^{r_1 x} & r_2 e^{r_2 x} \end{bmatrix}_{x=0}$$

$$\det \begin{vmatrix} 1 & 1 \\ r_1 & r_2 \end{vmatrix} = r_2 - r_1 \neq 0$$

Ex 2 $y = e^{rx}$

$$y'' + 8y' - 9y = 0$$

$$y(0) = 1$$

$$y'(0) = 0$$

$$r^2 + 8r - 9 = 0$$

$$(r+9)(r-1) = 0$$

$$r = 1, -9$$

$$y = C_1 e^{-9x} + C_2 e^x$$

$\therefore 2.90 \quad 0.9x \quad \cdot \quad \cdot$

$$y = c_1 e^{-x} + c_2 e^x$$

$$y(0) = c_1 + c_2 = 1$$

$$y'(0) = -c_1 + c_2 = 0 \Rightarrow c_2 = c_1$$

$$c_1 + c_1 = 1$$

$$c_1 = \frac{1}{2}$$

$$c_2 = \frac{1}{2}$$

$$y = \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

Case 2 170h

$$b^2 - 4ac = 0$$

$$r_1 = -\frac{b}{2a}$$

$$y_1 = e^{-\frac{b}{2a}x} \quad y_2 = v(x) e^{-\frac{b}{2a}x}$$

$$y = v(x) e^{-\frac{b}{2a}x}$$

$$y' = v'(x) e^{-\frac{b}{2a}x} - v(x) \frac{b}{2a} e^{-\frac{b}{2a}x}$$

$$= (v' - \frac{b}{2a}v) e^{-\frac{b}{2a}x}$$

$$y'' = (v'' - \frac{b}{2a}v') e^{-\frac{b}{2a}x} + \text{same method}$$

$$(v' - \frac{b}{2a}v) (-\frac{b}{2a} e^{-\frac{b}{2a}x})$$

$$= (v'' - \frac{b}{2a}v' + \frac{b^2}{4a}v) e^{-\frac{b}{2a}x}$$

$$ay'' + by' + cy = e^{-\frac{b}{2a}x} [av'' - bv' + \frac{b^2}{4a}v + bv' - \frac{b^2}{2a}v + cv]$$

$$= 0$$

$$= e^{-\frac{b}{2a}x} [av'' + \underset{\substack{\uparrow \\ \text{since} \\ \frac{b^2}{4a} = c}}{c}v - 2cv + cv] = 0$$

$$= e^{-\frac{b}{2a}x} [av''] = 0$$

$$\text{so } v'' = 0$$

$$\Rightarrow v' = 0$$

$$\Rightarrow v = \alpha + \beta x$$

$$y_2 = (\alpha + \beta x) e^{-\frac{b}{2a}x}$$

$$y_2 = \alpha y_1 + \beta x y_1$$

if $b^2 - 4ac = 0$, then 1 soln is

$$y_1 = e^{-\frac{b}{2a}x}$$

$$y_2 = x e^{-\frac{b}{2a}x}$$

$$\text{set } k = 0$$

Ex 1

$$y'' - 7y' + 12y = 0$$

$$\begin{aligned}
 r^2 - 2r + 1 &= 0 \\
 (r-1)^2 &= 0 \\
 r=1 &\rightarrow y_1 = e^x \\
 &\quad y_2 = x e^x
 \end{aligned}$$

gen soln $y = C_1 e^x + C_2 x e^x$

2nd Order linear Homogenous Const Coefficient Diff Eqns

Friday, April 20, 2007
9:04 AM

$$ay'' + by' + cy = 0 \quad ; \quad a \neq 0$$

2nd order linear homogeneous constant coefficient diff eqn

$$y = e^{rx} \quad ; \quad y' = r e^{rx} \quad ; \quad y'' = r^2 e^{rx}$$

$$e^{rx} [ar^2 + br + c] = 0$$

$\Rightarrow r$ solves $ar^2 + br + c = 0 \Leftrightarrow$ characteristic eqn

$$r = \frac{-b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

① $b^2 - 4ac > 0$

② $b^2 - 4ac = 0$

③ $b^2 - 4ac < 0$

① if $b^2 - 4ac > 0$

2 real roots that are distinct

$$r_1 = \frac{-b}{2a} + \frac{\sqrt{b^2 - 4ac}}{2a}$$

$$r_2 = \frac{-b}{2a} - \frac{\sqrt{b^2 - 4ac}}{2a}$$

$$y = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

② $b^2 - 4ac = 0$

$$r = -\frac{b}{2a}$$

$$y = c_1 e^{rx} + c_2 x e^{rx}$$

Ex. $y'' + 4y' + 4y = 0$

$$y(0) = 0$$

$$y'(0) = 1$$

$$r^2 + 4r + 4 = 0$$

$$(r+2)(r+2) = 0$$

$$r = -2$$

$$y = c_1 e^{-2x} + c_2 x e^{-2x}$$

$$y' = c_1 e^{-2x} - 2c_2 x e^{-2x}$$

$$y(0) = c_1 = 0$$

$$y'(0) = c_2 = 1$$

=

$$y'' + 4y' + 4y = \begin{cases} y = x e^{-2x} \\ y' = e^{-2x} - 2x e^{-2x} \\ y'' = -2e^{-2x} - 2e^{-2x} + 4x e^{-2x} = -4e^{-2x} + 4x e^{-2x} \end{cases}$$

$$= -4e^{-2x} + 4x e^{-2x} + 4e^{-2x} - 8x e^{-2x} + 4x e^{-2x} - 8x e^{-2x} = 0$$

Beher
Professional Patent

$$+ 4xe^{-2x} = 0$$

$$\text{if } b^2 - 4ac < 0$$

$$\begin{array}{l} 2 \text{ complex roots} \quad r = \alpha \pm i\beta \\ \left. \begin{array}{l} r_1 = \alpha + i\beta \\ r_2 = \alpha - i\beta \end{array} \right\} \text{"complex conjugate pairs"} \end{array}$$

$$r = \frac{-b}{2a} \pm \sqrt{\frac{b^2 - 4ac}{2a}}$$

$$= \frac{-b}{2a} \pm \frac{\sqrt{(-1)(4ac - b^2)}}{2a}$$

$$= \frac{-b}{2a} \pm \frac{i\sqrt{4ac - b^2}}{2a}$$

$$\alpha = -\frac{b}{2a} \quad \beta = \frac{\sqrt{4ac - b^2}}{2a}$$

$$y_1 = e^{(\alpha + i\beta)x}$$

$$= e^{\alpha x} e^{i\beta x}$$

$$= e^{\alpha x} [\cos(\beta x) + i\sin(\beta x)]$$

Euler's Formula

$$y_2 = e^{(\alpha - i\beta)x}$$

$$= e^{\alpha x} e^{-i\beta x}$$

$$= e^{\alpha x} [\cos(\beta x) - i\sin(\beta x)]$$

$$e^{i\theta} = \cos \theta + i\sin \theta$$

$$e^{\pm i\theta} = \cos \theta \pm i\sin \theta$$

$$u_1 = \frac{1}{2}(y_1 + y_2)$$

$$= e^{\alpha x} \cos(\beta x)$$

$$u_2 = \frac{1}{2i}(y_1 - y_2)$$

$$= e^{\alpha x} \sin(\beta x)$$

$$\text{Det} \begin{bmatrix} c_1 e^{\alpha x} \cos \beta x & c_2 e^{\alpha x} \sin \beta x \\ c_1 \alpha x e^{\alpha x} \cos \beta x - c_1 e^{\alpha x} \sin \beta x & c_2 \alpha x e^{\alpha x} \sin \beta x + c_2 e^{\alpha x} \cos \beta x \end{bmatrix}$$

$$\textcircled{1} b^2 - 4ac > 0 \quad y = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

$$\textcircled{2} b^2 - 4ac = 0 \quad y = c_1 e^{rx} + x c_2 e^{rx}$$

$$\textcircled{3} b^2 - 4ac < 0 \quad y = c_1 e^{\alpha x} \cos \beta x + c_2 e^{\alpha x} \sin \beta x$$

$$\alpha = \text{Re}(r) \quad \text{real}$$

$$\beta = \text{Im}(\beta) \quad \text{imaginary}$$

Use Taylor's series to motivate Euler's Formula.

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} \dots$$

$$e^{i\theta} = 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} \dots$$

$$i^2 = -1 \quad i^3 = -i \quad i^4 = 1 \quad i^5 = i \quad i^6 = -1$$

$$e^{i\theta} = 1 + i\theta + \frac{-\theta^2}{2} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \frac{\theta^6}{6!} \dots$$

$$= \underbrace{\left(1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \frac{\theta^8}{8!} \dots\right)}_{\cos \theta} + i \underbrace{\left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} \dots\right)}_{\sin \theta}$$

$$z = \cos \theta + i \sin \theta$$

$$\frac{dz}{d\theta} = -\sin \theta + i \cos \theta$$

$$\frac{dz}{d\theta} = i(\sin \theta + i \cos \theta) = iz$$

$$\frac{dz}{d\theta} = iz \Rightarrow z = ce^{i\theta}$$

Ex

$$u_1 = e^{\alpha x} \cos(\beta x)$$

$$u_2 = e^{\alpha x} \sin(\beta x)$$

$$W(u_1, u_2) = \det \begin{bmatrix} u_1 & u_2 \\ u_1' & u_2' \end{bmatrix}$$

$$\det \begin{bmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) & \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x) \end{bmatrix}$$

$$W(u_1, u_2) \Big|_{x=0} = \det \begin{bmatrix} 1 & 0 \\ \alpha & \beta \end{bmatrix} = \beta$$

β cannot be 0 b/c that would be a diff case.

Ex $y'' + 2y' + 5y = 0$

$$y(0) = 0$$

$$y'(0) = 1$$

$$r^2 + 2r + 5 = 0$$

$$r = \frac{-2 \pm \sqrt{4 - 4(5)}}{2a} = \frac{-2 \pm 4i}{2a} = -1 \pm 2i$$

$$y = C_1 e^{-x} \cos(2x) + C_2 e^{-x} \sin(2x)$$

$$y(0) = C_1 = 0 \Rightarrow y = C_2 e^{-x} \sin(2x)$$

$$y' = C_2 e^{-x} \sin(2x) + 2C_2 e^{-x} \cos(2x)$$

$$y'(0) = 2C_2 = 1$$

Soln

Soln

$$y = \frac{1}{2} e^{-x} \sin(2x)$$

$$y = a_1 e^{(-1+2i)x} + a_2 e^{(-1-2i)x}$$

Linear n^{th} order homogeneous differential eqns

$$y^{(n)} = \frac{d^{(n)}y}{dx^n} = n^{\text{th}} \text{ derivative}$$

$$(*) y^{(n)} + p_1(x)y^{(n-1)} + p_2(x)y^{(n-2)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$

There is a superposition principle

If $y_1(x)$ & $y_2(x)$ solve $(*)$ then so does

$$c_1 y_1(x) + c_2 y_2(x) = y \text{ for any real constants } c_1, c_2$$

We can expect n constants of integration for n^{th} order eqn.

Unique solⁿ requires n initial conditions

General soln of $(*)$ is $y(x) = c_1 y_1 + c_2 y_2 + \dots + c_n y_n$
where $y_1, y_2, y_3, \dots, y_n$ are n linearly ind. solns of $(*)$

Solns $y_1, \dots, y_n$ are linearly independent if the only solⁿ to $c_1 y_1 + c_2 y_2 + c_3 y_3 + \dots + c_n y_n = 0$ is to choose $c_1 = c_2 = c_3 = \dots = c_n = 0$

Otherwise one of the solns could be expressed as a linear combination of the others.

$n \times n$ matrix

We test for linear independence of a set of functions $\{y_1, y_2, \dots, y_n\}$ by computing the Wronskian

$$W(y_1, \dots, y_n) = \det \begin{bmatrix} y_1 & y_2 & y_3 & \dots & y_n \\ y_1' & y_2' & y_3' & \dots & y_n' \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & y_3^{(n-1)} & \dots & y_n^{(n-1)} \end{bmatrix}$$

Must get a square matrix

If $W \neq 0 \Rightarrow y_1, \dots, y_n$ are linearly dependent
If $W \neq 0 \Rightarrow y_1, \dots, y_n$ are linearly independent

Can check w/ just 1 pt.

Problem: Use the Wronskian to show that $y'' + p(x)y' + q(x)y = 0$ cannot have 3 l.n. ind. solns.

y_1, y_2, y_3

$$\begin{aligned}
 W(y_1, y_2, y_3) &= \det \begin{bmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{bmatrix} \\
 &= y_1(y_2' y_3'' - y_3' y_2'') - y_2(y_1' y_3'' - y_3' y_1'') \\
 &\quad + y_3(y_1' y_2'' - y_2' y_1'') \\
 &= y_1(y_2'(-p y_3' - q y_3) - y_2'(-p y_2' - q y_2)) \\
 &\quad - y_2(y_1'(-p y_3' - q y_3) - y_3'(-p y_1' - q y_1)) \\
 &\quad + y_3(y_1'(-p y_2' - q y_2) - y_2'(-p y_1' - q y_1)) \\
 &= 0
 \end{aligned}$$

Diff eqns w/ const. coefficients

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + a_{n-2} y^{(n-2)} + \dots + a_1 y' + a_0 y = 0$$

Assume soln in the form of $e^{rx} = y$

$$e^{rx} [r^n a_n + a_{n-1} r^{n-1} + \dots + a_1 r + a_0] = 0$$

characteristic eqn that r must satisfy

Ex If the characteristic eqn is $(r-1)(r-2)^3(r^2+4)^2 = 0$
What is the general soln?

Roots 8th order so 8 roots.
 $\{1, 2, 2, 2, 2i, -2i, 2i, -2i\}$

Prescription for going from roots of the char eqn to the gen soln

(1) n real, distinct roots, then we have n linearly ind. solns

$$e^{r_1 x}, e^{r_2 x}, \dots, e^{r_n x} \quad \{r_1, \dots, r_n\}$$

(2) Any ^{real} root r , that is repeated k times ($k=2$ or 3 or n)
then k solns associated w/ that root, $e^{rx}, x e^{rx}, x^2 e^{rx}, \dots, x^{k-1} e^{rx}$
 r are

(3) Distinct complex pair $r = \alpha + i\beta$ & $r = \alpha - i\beta \Rightarrow 2$ (in ind solns).
 $e^{\alpha x} \cos(\beta x), e^{\alpha x} \sin(\beta x)$

(4) Repeated complex roots $r = \alpha + i\beta$, $r = \alpha - i\beta$ repeated k

times. Same thing.

$$e^{\alpha x} \cos(\beta x), e^{\alpha x} \sin(\beta x), \\ x e^{\alpha x} \cos(\beta x), x e^{\alpha x} \sin(\beta x), \\ \dots, x^{k-1} e^{\alpha x} \cos(\beta x), x^{k-1} e^{\alpha x} \sin(\beta x)$$

$$y(x) = c_1 e^x + c_2 e^{2x} + c_3 x e^{2x} + c_4 x^2 e^{2x} + c_5 \cos(2x) + c_6 \sin(2x) \\ + c_7 x \cos(2x) + c_8 x \sin(2x)$$

$$a_n r^n + a_{n-1} r^{n-1} + \dots + a_1 r + a_0 = 0$$

$$a_n \neq 0$$

(*) $a_n, a_{n-1}, \dots, a_0$ are real

claim if $r = \alpha + i\beta$ ($\beta \neq 0$)


is a soln of (*) then so is $\bar{r} = \alpha - i\beta$

$$a_n (r - r_1)(r - r_2) \dots (r - r_n) = 0$$



Spring Constant

Tuesday, April 24, 2007

8:58 AM



$k = \text{spring constant}$



$$\vec{F} = m\vec{a} \quad \vec{a} = \frac{d^2x}{dt^2} \hat{i} = \ddot{x} \hat{i}$$

$$\vec{F} = \vec{F}_{\text{grav}} + \vec{F}_{\text{spring}} + \vec{F}_{\text{dumper}} \quad \text{def used when indep. var. is } t$$

$$= \hat{i} [mg - kx - b\dot{x}]$$

$$m\ddot{x} = mg - kx - b\dot{x}$$

2nd order lin eqn. Not homogeneous b/c of mg

$$m\ddot{x} + kx + b\dot{x} = mg$$


Introduce a new coordinate system where $y = x - x_s$

$$x = x_s + y$$

$$x_s : \dot{x} = \ddot{x} = 0$$

$$kx_s = mg$$

$$y = x - \frac{mg}{k}$$



$$\dot{y} = \dot{x}$$

$$\ddot{y} = \ddot{x}$$

$$m\ddot{y} + b\dot{y} + k(y + \frac{mg}{k}) = mg$$

$$m\ddot{y} + b\dot{y} + ky = 0$$

$$\Gamma y e^{rt} = y$$

$$mr^2 + br + k = 0$$

$$r = \frac{-b \pm \sqrt{b^2 - 4(m)(k)}}{2m}$$

Ⓘ Undamped Case $b=0$

$$r = \pm \frac{\sqrt{-4mk}}{2m} = \pm \frac{2i\sqrt{mk}}{2m} = \pm i \frac{\sqrt{mk}}{m} = \pm i \sqrt{\frac{k}{m}} = \pm i\omega_0$$

$$y = c_1 \cos(\omega_0 t) + c_2 \sin(\omega_0 t)$$

or equivalently $y = A \cos(\omega_0 t - \phi)$
 $\{c_1, c_2\}$ or $\{A, \phi\}$

$$\cos(A-B) = \text{Re}[e^{i(A-B)}] = \text{Re}[e^{iA} e^{-iB}]$$

↑ "real part"

$$\text{Re}[(\cos A + i \sin A)(\cos B - i \sin B)]$$

$$= \cos A \cos B + \sin A \sin B$$


$$A \cos(\omega_0 t) \cos(\phi) + A \sin(\omega_0 t) \sin(\phi)$$

$$A \cos(\phi) = c_1, \quad A \sin(\phi) = c_2$$

$$A = \sqrt{c_1^2 + c_2^2}$$

$$\phi = \tan^{-1}\left(\frac{c_2}{c_1}\right)$$

$$\frac{A \sin \phi}{A \cos \phi} = \frac{c_2}{c_1}$$



$$T = \frac{2\pi}{\omega_0}$$

II Damped Case

- (A) $b^2 - 4mk < 0$ b/c b is very small damped vibration
- (B) $b^2 = 4mk$ "critically damped"
- (C) $b^2 - 4mk > 0$ overdamped case

$$(A) r = -\frac{b}{2m} \pm i \sqrt{\frac{4mk - b^2}{4m^2}}$$

$$= -\frac{b}{2m} \pm i \sqrt{\frac{k}{m} - \frac{b^2}{4m^2}}$$

$$\gamma = \frac{b}{2m} \quad \omega_0 = \sqrt{\frac{k}{m}}$$


$$= -\gamma \pm i \sqrt{\omega_0^2 - \gamma^2}$$

$$= c_1 e^{-\gamma t} \cos(\omega t) + c_2 e^{-\gamma t} \sin(\omega t)$$

$$= A e^{-\gamma t} \cos(\omega t - \phi)$$


$$\omega^2 = \omega_0^2 - \gamma^2$$

$$\omega^2 < \omega_0^2$$



$$y = A \cos(\omega t - \phi)$$

$$0 < b < \sqrt{4mk}$$



$$y = A e^{-\gamma t} \cos(\omega t - \phi)$$

amp. of cos function

$$-Ae^{-\delta t}$$

② $b = \sqrt{4mk}$ $y = c_1 e^{-\delta t} + c_2 t e^{-\delta t}$
 $= e^{-\delta t}(c_1 + c_2 t)$


if c_1, c_2 are both pos. then $y(t)$ never goes through $y=0$

if $c_1 > 0$ & $c_2 < 0$ then $y(t) = 0$ when $t = -\frac{c_1}{c_2}$

$$y = e^{-\delta t}(c_1 + c_2 t) \quad y' = -\delta e^{-\delta t}(c_1 + c_2 t) + c_2 e^{-\delta t} = \dot{y}$$

$$y(t=0) = c_1 = \text{initial displacement}$$

$$\dot{y}(0) = -\delta c_1 + c_2 = \text{initial velocity}$$



Linear Non-Homogeneous eqns

eg $y'' + p(x)y' + q(x)y = f(x) \neq 0 \quad *$

$[f(x) = 0 \Rightarrow \text{homogeneous}]$

Introduce as shorthand linear operator L

$$L = \frac{d^2}{dx^2} + p(x)\frac{d}{dx} + q(x)$$

$$Ly = \frac{d^2 y}{dx^2} + p(x)\frac{dy}{dx} + q(x)y$$

Eq $*$ is then written $Ly = f(x)$

Observation 1

$y = 0$ is not a soln of $*$

Observation 2

Eqn $*$ does not have a superposition principle

If y_1 & y_2 solve $*$ then $y = c_1 y_1 + c_2 y_2$ does not necessarily solve it.

$$Ly_1 = f(x)$$

$$Ly_2 = f(x)$$

$$L(c_1 y_1 + c_2 y_2) = c_1 Ly_1 + c_2 Ly_2 \\ = (c_1 + c_2)f \neq f$$

If $c_1 + c_2 = 1$, then it works - otherwise no.

Observation 3

If y_1 & y_2 solve $*$ then it must be that

$y_2 = y_1 + y_h$, where y_h is a soln of the homogeneous problem associated w/ $*$.

$Ly_h = 0$

$Ly_h = 0$

$$Ly_1 = f(x)$$

$$Ly_2 = f(x)$$

$$L(y_2 - y_1) = Ly_2 - Ly_1 = f - f = 0$$

$$L(y_2 - y_1) = 0$$

∴ solves the homogeneous problem,

Strategy for finding the general soln of *

① Find a particular soln $y = y_p$ ($Ly_p = f(x)$)

② Solve the associated homogeneous problem ($Ly_h = 0$)
(gen soln)

③ Gen soln is $y = y_p + y_h$

Ex 1

$$y'' - y = e^{2x}$$

Look for particular soln $y_p = ce^{2x}$
 $y_p'' = 4ce^{2x}$

$$y_p'' - y_p = 4ce^{2x} - ce^{2x} = 3ce^{2x} = e^{2x} \quad \boxed{c = \frac{1}{3}} \quad \checkmark$$

$$① y = \frac{1}{3} e^{2x}$$

$$y_p'' - y_h = 0$$

$$r^2 - 1 = 0$$

$$r = \pm 1$$

$$② y_h = c_1 e^x + c_2 e^{-x}$$

$$③ \boxed{\frac{1}{3} e^{2x} + c_1 e^x + c_2 e^{-x} = y}$$

Ex 2

$$y'' + y = x^2$$

Look for particular soln

$$y_p = x^2 + Ax + B$$

$$y_p'' = 2$$

$$y_p'' + y_p = x^2 + \underbrace{A x}_{A=0} + \underbrace{(B x^2)}_{B=-2} = x^2$$

$$\textcircled{1} y_p = x^2 - 2$$

$$y_h'' + y_h = 0$$

$$r^2 + 1 = 0$$

$$r = \pm i$$

$$y_h = C_1 \cos(x) + C_2 \sin(x)$$

$$y = x^2 - 2 + C_1 \cos(x) + C_2 \sin(x)$$

Ex3

$$y'' - 2y' - y = \sin(x)$$

$$y_p = A \cos(x) + B \sin(x)$$

$$y_p' = -A \sin(x) + B \cos(x)$$

$$y_p'' = -A \cos(x) - B \sin(x)$$

$$\sin(x) = -A \cos(x) - B \sin(x) + 2A \sin(x) - 2B \cos(x) - A \cos(x) - B \sin(x)$$

$$\sin(x) = \sin(x) [-B + 2A - B] + \cos(x) [-A - 2B - A]$$

$$2(A - B) = 1$$

$$-2(A + B) = 0$$

$$B = -A$$

$$A = \frac{1}{4}$$

$$B = -\frac{1}{4}$$

$$\textcircled{1} \text{ Particular soln } \frac{1}{4}(\cos(x) - \sin(x))$$

$$y_h'' - 2y_h' - y_h = 0$$

$$r^2 - 2r - 1 = 0$$

$$r = \frac{2 \pm \sqrt{4+4}}{2} = 1 \pm \sqrt{2}$$

② Homogeneous soln

$$y_h = C_1 e^{(1+\sqrt{2})x} + C_2 e^{(1-\sqrt{2})x}$$

$$y = y_p + y_h$$

Method of Undetermined Coefficients

Works when $f(x)$ involves functions like $e^{\alpha x}$, $\cos(\beta x)$, $\sin(\beta x)$, polynomials, & combinations of these.

Solving * w/ const. coefficients

e.g. $e^{\alpha x} \cos(\beta x)$ or $x^3 e^x$

Ex 4

$$y'' + ay' + by = e^x$$

I, what instance does $y_p \neq ce^x$?

$$y'' + 2y' - y = x^3$$

$$y_p = Ax^3 + Bx^2 + Cx + D$$

$$\begin{matrix} \uparrow \\ (-1) \end{matrix}$$

y

Examples

Tuesday, May 01, 2007
9:07 AM

Ex 1 $y'' + 2y' + y = xe^{2x}$ Ex 2 $y'' - y' = e^x$ Ex 3 $y'' - y' = x^2 e^x$
Ex 4 $y'' + 2y' + y = e^{-x}$ Ex 5 $y'' + y = \sin x$

Ex 1 $y'' + 2y' + y = xe^{2x}$

$$y_p = Axe^{2x} + Be^{2x}$$

$$y_p' = 2Axe^{2x} + Ae^{2x} + 2Be^{2x} = (A+2B)e^{2x} + 2Axe^{2x}$$

$$y_p'' = 2(A+2B)e^{2x} + 2Ae^{2x} + 4Axe^{2x} = (4A+4B)e^{2x} + 4Axe^{2x}$$

$$y_p'' + 2y_p' + y_p = e^{2x} [4A+4B + 2A+4B+B] + xe^{2x} [4A+4A+A] = xe^{2x}$$

$4A+4B=0$ if eqn =
 $9A=1$ if eqn =

$$A = \frac{1}{9}$$

$$B = -\frac{2}{27}$$

$$r^2 + 2r + 1 = 0 = (r+1)^2$$

homogeneous soln = $y = c_1 e^{-x} + c_2 x e^{-x}$

$$y = y_p + y_h = \frac{1}{9} x e^{2x} - \frac{2}{27} e^{2x} + c_1 e^{-x} + c_2 x e^{-x}$$

Ex 2 $y'' - y' = e^x$

$$y_p = Ae^x$$

$$y_p' = Ae^x$$

$$y_p'' = Ae^x$$

$$Ae^x - Ae^x = e^x$$

$$0 = e^x$$

so there is no way to choose A to get e^x out.

$$y'' - y' = 0$$

$$r^2 - r = 0$$

$$r^2 = r$$

$$r = 1, 0$$

$$y_h = c_1 + c_2 e^x$$

this is a soln, & we used a guess like this so it doesn't work.

$$u' - u = e^x$$

$$\mu = e^{\int -1 dx} = e^{-x}$$

$$e^{-x} u' - e^{-x} u = 1$$

$$\left(\frac{d}{dx} (e^{-x} u) \right) = 1$$

$$e^x u = x + C$$

$$u = xe^x + Ce^x$$

$$y' = xe^x + Ce^x$$

$$y = \int (xe^x + Ce^x) dx$$

$$y = xe^x - e^x + B + Ce^x$$

$$= xe^x + C_1 + C_2 e^x$$

$$\int u dv = uv - \int v du$$

$$\int xe^x dx$$

$$u = x \quad du = dx$$

$$dv = e^x dx \quad v = e^x$$

$$\int xe^x dx = xe^x - \int e^x dx$$

$$= xe^x - e^x + B$$

Advice

First determine y_h , so you will know what to guess for y_p

Exception #1 if the forcing term $f(x)$ (right-hand side) contains an exponential or a trig function or a combo of the 2 that comes from a simple root (not repeated) of the characteristic eqn then one must multiply the trial function (your guess) for y_p we'd normally use by x .

Ex. 2

① Find homogeneous soln

$$y_h'' - y_h' = 0$$

$$r^2 - r = 0$$

$$r(r-1) = 0$$

$$y_h = C_1 + C_2 e^x$$

② See e^x change to xe^x

$$y_p = Axe^x$$

$$y_p' = Ae^x + Axe^x$$

$$y_p'' = Ae^x + Ae^x + Axe^x$$

$$= 2Ae^x + Axe^x$$

$$y_p'' - y_p' = 2Ae^x + Axe^x - Ae^x - Axe^x = Axe^x \Rightarrow A=1$$

$$y = y_h + y_p = C_1 + C_2 e^x + xe^x$$

Ex 3. $y'' - y' = x^2 e^x$

① $y_h'' - y_h' = 0$

$$y_h = C_1 + C_2 e^x$$

② $y_p = x(Ax^2 e^x + Bx e^x + \underbrace{Ce^x}_{\text{prob term}})$

$$y_p = Ax^3 e^x + Bx^2 e^x + Cx e^x$$

$$y_p' = 3Ax^2 e^x + Ax^3 e^x + 2Bx e^x + Bx^2 e^x + Ce^x + Cx e^x$$

$$y_p'' = y_p' + 3Ax^2 e^x + 2Bx e^x + Ce^x + 6Ax e^x + 2B e^x$$

$$y_p'' - y_p' = 3Ax^2 e^x + (2B+6A)x e^x + (C+2B)e^x$$

$$\begin{aligned}
 &3A \text{ must be } 1 \\
 &2B + 6A \text{ must be } 0 \\
 &C + 2B = 0 \\
 &A = \frac{1}{3} \\
 &B = -1 \\
 &C = 2
 \end{aligned}$$

$$y = C_1 + e^x + \frac{1}{3}x^3e^x - x^2e^x + 2xe^x$$

$$y'' + ay' + by = f(x)$$

Lab due Mon May 14

$$y = y_h + y_p$$

y_p = "particular soln"

y_h = gen soln of homogeneous prob

$$y_h'' + ay_h' + by_h = 0$$

Step 1 Find y_h

Step 2 Check to see relationship btwn y_h & guess y_p
If not, then "guess" y_p w/ undetermined
coeffs & solve for these

Exception 1 If yes, then modify guess by multiplying it
by x if the root is simple

Exception 2 If the forcing $f(x)$ contains an exponential
that comes from a repeated root of the
characteristic eqn, then multiply the trial function
for y_p by x^2

Ex

$$y'' + 2y' + y = e^{-x}$$

$y_c = y_h$ in the
book

$$\textcircled{1} \quad y_h'' + 2y_h' + y_h = 0$$

$$r^2 + 2r + 1 = 0$$

$$(r+1)^2 = 0$$

$r = -1 \Rightarrow$ repeated root

$$y_h = C_1 e^{-x} + C_2 x e^{-x}$$

$$\textcircled{2} \quad y_p = (A e^{-x}) x^2 = A x^2 e^{-x}$$

$$y_p' = 2A x e^{-x} - A x^2 e^{-x}$$

$$y_p'' = 2A e^{-x} - 2A x e^{-x} - 2A x e^{-x} + A x^2 e^{-x}$$

$$y_p'' = 2A e^{-x} - 4A x e^{-x} + A x^2 e^{-x}$$

$$y_p'' + 2y_p' + y_p = e^{-x}$$

$$= e^{-x} [2A]$$

$$+ x e^{-x} [-4A + 4A]$$

$$+ x^2 e^{-x} [A - 2A + A] = 2A e^{-x} = e^{-x}$$

$$A = \frac{1}{2}$$

$$y = \frac{1}{2} x^2 e^{-x} + C_1 e^{-x} + C_2 x e^{-x}$$

" - $x^2 x$

$$\ddot{x} = \frac{d^2x}{dt^2}$$

$$m\ddot{x} + b\dot{x} + kx = F_0 \cos(\omega t)$$

Forced Mechanical Vibration
- resonance

Homogeneous soln

$$m\ddot{x}_h + b\dot{x}_h + kx_h = 0$$

$$mr^2 + br + k = 0$$

$$-b \pm \frac{\sqrt{b^2 - 4(m)k}}{2m} = r$$

$$-\frac{b}{2m} \pm \sqrt{\left(\frac{b}{2m}\right)^2 - \frac{k}{m}}$$

Focus on
decaying oscillations
when not much damping

$$\left(\frac{b}{2m} < \sqrt{\frac{k}{m}}\right)$$


$$\omega_0 = \sqrt{\frac{k}{m}} = \text{frequency of oscillations in undamped}$$

$$\gamma = \frac{b}{2m} = \text{decay rate associated w/ damping}$$

$$-\gamma \pm \sqrt{\omega_0^2 - \gamma^2}$$

$$x_h = c_1 e^{-\gamma t} \cos(\sqrt{\omega_0^2 - \gamma^2} t) + c_2 e^{-\gamma t} \sin(\sqrt{\omega_0^2 - \gamma^2} t)$$

damping causes shift to lower freq.
 $\sqrt{\omega_0^2 - \gamma^2} < \omega_0$



$$x = R e^{-\gamma t} \cos(\sqrt{\omega_0^2 - \gamma^2} t + \phi)$$

$$m\ddot{x}_p + b\dot{x}_p + kx = F_0 \cos(\omega t)$$

unless both $\gamma = 0$ & $\omega = \omega_0$ the forcing term is not
one of the homogeneous solns

$$x_p = A \cos(\omega t) + B \sin(\omega t)$$

$$\dot{x}_p = -\omega A \sin(\omega t) + \omega B \cos(\omega t)$$

$$\ddot{x}_p = -\omega^2 A \cos(\omega t) - \omega^2 B \sin(\omega t)$$

$$m\ddot{x}_p + b\dot{x}_p + kx_p = \cos(\omega t) \underbrace{[-m\omega^2 A + b\omega B + kA]}_{F_0} + \sin(\omega t) \underbrace{[-m\omega^2 B - b\omega A + kA]}_{0} = F_0 \cos(\omega t)$$

$$\left. \begin{aligned} \frac{1}{m} [(-m\omega^2 + k)A + (b\omega)B] &= F_0 \\ \frac{1}{m} [(-m\omega^2 + k)B - (b\omega)A] &= 0 \end{aligned} \right\} \begin{aligned} (\omega_0^2 - \omega^2)A + (2\gamma\omega)B &= \frac{F_0}{m} \\ (\omega_0^2 - \omega^2)B - (2\gamma\omega)A &= 0 \end{aligned}$$

$$\left[\frac{(\omega_0^2 - \omega^2)^2}{2\gamma^2\omega} + \frac{(2\gamma\omega)^2}{2\gamma\omega} \right] B = \frac{F_0}{m}$$

$$B = \frac{2\gamma\omega F_0}{m[(\omega_0^2 - \omega^2)^2 + (2\gamma\omega)^2]}$$

$$x_p = \frac{F_0}{m[(\omega_0^2 - \omega^2)^2 + (2\gamma\omega)^2]} \left[(\omega_0^2 - \omega^2) \cos(\omega t) + 2\gamma\omega \sin(\omega t) \right] A = \frac{(\omega_0^2 - \omega^2) F_0}{m[(\omega_0^2 - \omega^2)^2 + (2\gamma\omega)^2]}$$

$$= R \cos(\omega t + \phi)$$

$$C = \sqrt{C_1^2 + C_2^2}$$

$$R^2 = \frac{C_1^2 + C_2^2}{\frac{F_0^2 \sqrt{(\omega_0^2 - \omega^2)^2 + (2\gamma\omega)^2}}{m \text{ (all that stuff)}}}$$

$$R = \frac{F_0}{m \left(\sqrt{(\omega_0^2 - \omega^2)^2 + (2\gamma\omega)^2} \right)^{-1}}$$

Denominator is small when $\omega_0 = \omega$
Biggest response.

To solve a 2nd order DEQ $y'' = f(x, y, y')$ (*)

Using Euler or Runge-Kutta or... first rewrite (*) as 2 1st order eqns

$$y' = v$$

$$m\ddot{x} + b\dot{x} + kx = F_0 \cos(\omega t)$$

$$\dot{x} = v$$

$$v' = y'' = f(x, y, v)$$

$$\dot{v} = \frac{1}{m} (F_0 \cos \omega t - b v - k x)$$

Introduce a vector variable $\vec{u} = \begin{pmatrix} x \\ v \end{pmatrix}$

$$\vec{u}' = \begin{pmatrix} v \\ f(x, v) \end{pmatrix} = \vec{F}(x, \vec{u})$$

$$\vec{u} = \begin{pmatrix} x \\ v \end{pmatrix} \Rightarrow \dot{\vec{u}} = \begin{pmatrix} v \\ \frac{1}{m} (F_0 \cos \omega t - b v - k x) \end{pmatrix}$$

Euler method

$$\vec{u}' = \vec{F}(x, \vec{u})$$

$$\vec{u}' = \lim_{h \rightarrow 0} \frac{\vec{u}(x+h) - \vec{u}(x)}{h} = \vec{F}(x, \vec{u})$$

$$\vec{u}(x_{n+1}) = \vec{u}_{n+1} = \vec{u}_n + h \vec{F}(x_n, \vec{u}_n)$$

RK ~~for~~ $y' = f(x, y)$

$$y_{n+1} = y_n + \frac{h}{2} (k_1 + 2k_2 + 2k_3 + k_4)$$

$$k_1 = f(x_n, y_n)$$

RK ~~3~~ $\vec{u}' = \vec{F}(x, \vec{u})$

$$\vec{u}_{n+1} = \vec{u}_n + \frac{h}{6} (\vec{k}_1 + 2\vec{k}_2 + 2\vec{k}_3 + \vec{k}_4)$$

$$\vec{k}_1 = \vec{F}(x_n, \vec{u}_n)$$

Forced vibration

$$m\ddot{x} + b\dot{x} + kx = F_0 \cos \omega t$$

$$\left. \begin{aligned} r &= -\gamma + i\sqrt{\omega_0^2 - \gamma^2} \\ \omega_0 &= \sqrt{\frac{k}{m}} \\ \gamma &= \frac{b}{2m} \end{aligned} \right\} x_h = e^{-\gamma t} \left[c_1 \cos(\sqrt{\omega_0^2 - \gamma^2} t) + c_2 \sin(\sqrt{\omega_0^2 - \gamma^2} t) \right]$$

transient
pt. of soln $\left\{ \begin{array}{l} \text{homogeneous soln} \rightarrow 0 \text{ as} \\ t \rightarrow \infty \end{array} \right.$

$$x_p(t) = A \cos \omega t + B \sin \omega t$$

$$A = \frac{F_0 (\omega_0^2 - \omega^2)}{m[(\omega_0^2 - \omega^2)^2 + 4\gamma^2 \omega^2]}$$

$$B = \frac{2\gamma \omega F_0}{m[(\omega_0^2 - \omega^2)^2 + 4\gamma^2 \omega^2]}$$

$$x(t) = x_h(t) + x_p(t) \quad \text{as } t \rightarrow \infty \quad x(t) \rightarrow x_p(t)$$

$$\text{As } t \rightarrow \infty$$

$$x(t) \rightarrow x_p(t)$$

$$x_p(t) = R \cos(\omega t + \phi)$$

$$\cos(A+B) = \underset{\substack{\uparrow \\ \text{"real part"}}}{\text{Re}(e^{i(A+B)})} = \text{Re}(e^{iA} e^{iB})$$

$$= \text{Re}(\cos A + i \sin A)(\cos B + i \sin B)$$

$$= \cos A \cos B - \sin A \sin B$$

$$x_p(t) = R \cos \omega t \cos \phi + R \sin \omega t \sin \phi$$

$$A = R \cos \phi \quad B = R \sin \phi$$

$$\tan \phi = \frac{B}{A}$$

$$A^2 + B^2 = R^2$$

$$\phi = \arctan \frac{B}{A}$$


$$R = \sqrt{A^2 + B^2}$$

$$R = \sqrt{A^2 + B^2}$$


$$R^2 = \frac{F_0^2 [(\omega_0^2 - \omega^2)^2 + 4\delta^2 \omega^2]}{m^2 [(\omega_0^2 - \omega^2)^2 + 4\delta^2 \omega^2]^2}$$

$$R = \frac{F_0}{m \sqrt{(\omega_0^2 - \omega^2)^2 + 4\delta^2 \omega^2}}$$


$$R \propto F_0$$



examine the dependence of R on ω



$$M = \frac{1}{\sqrt{(\omega_0^2 - \omega^2)^2 + 4\delta^2 \omega^2}}$$



Resonant Response at $\omega = \omega_0$

δ small: large narrow peak in $R(\omega)$
 δ larger: smaller, broader peak in $R(\omega)$

Assume $\frac{\delta}{\omega_0} \ll 1$ & $\omega \approx \omega_0$

$$(\omega_0^2 - \omega^2)^2 = (\omega_0 - \omega)^2 \underbrace{(\omega_0 + \omega)^2}_{\approx 2\omega_0^2} \approx (\omega_0 - \omega)^2 4\omega_0^2$$

$$M = \frac{1}{\sqrt{(\omega_0^2 - \omega^2)^2 + 4\gamma^2 \omega^2}} \approx \frac{1}{\sqrt{4\omega_0^2 [(\omega_0 - \omega)^2 + \gamma^2]}}$$


$\uparrow$ $\uparrow$
 $(\omega_0 - \omega)^2 4\omega_0^2$ $\approx \omega_0$

$$\frac{1}{2\omega_0} \frac{1}{\sqrt{(\omega_0 - \omega)^2 + \gamma^2}} \leftarrow \text{"Lorentzian" for probability peak is at } \omega_0 = \omega$$

$$R_{\text{peak}}^L = \frac{F_0^2}{n^2 4\omega_0^2 \gamma^2}$$

Frequency

Monday, May 07, 2007
8:59 AM



$R_{max} \approx R(\omega_0)$ for small γ

$$R = \frac{F_0}{m} \frac{1}{\sqrt{(\omega_0^2 - \omega^2)^2 + 4\gamma^2 \omega^2}}$$

$$R(\omega_0) = \frac{F_0}{2m\gamma\omega_0} \Rightarrow R \uparrow \text{ as } \frac{1}{\gamma}$$

Determine $\Delta\omega$ by estimating the values of ω for which

$$[R(\omega)]^2 \approx \left(\frac{R_{max}}{\sqrt{2}}\right)^2 \approx \frac{R_{max}^2}{2}$$

b/c the peak does not occur exactly at ω_0

$$R^2 = \frac{F_0^2}{m^2((\omega_0^2 - \omega^2)^2 + 4\gamma^2 \omega^2)} \quad \frac{1}{2} R_{max}^2 \approx \frac{F_0^2}{8m^2\gamma^2 \omega_0^2}$$

ω_0 by approx

$$(\omega_0^2 - \omega^2)^2 = (\omega_0 - \omega)^2 (\omega_0 + \omega)^2 \approx 4\omega_0^2 (\omega_0 - \omega)^2$$


ω_0 by approx

$$\frac{1}{2} R_{max}^2 = R^2 \approx \frac{F_0^2}{m^2 4 \omega_0^2 [(\omega_0 - \omega)^2 + \gamma^2]} \approx \frac{F_0^2}{8m^2 \gamma^2 \omega_0^2}$$

$$(\omega_0 - \omega)^2 + \gamma^2 \approx 2\gamma^2$$

$$(\omega_0 - \omega)^2 = \gamma^2$$

Q: Mon Captain!



$$\Delta\omega \approx 2\gamma \text{ b/c } \omega_0 + \gamma - (\omega_0 - \gamma)$$

$$Q = \text{Quality factor} = \frac{\omega_0}{\Delta\omega} \approx \frac{\omega_0}{2\gamma}$$

as $\gamma \downarrow$, $Q \uparrow$

$$Q \text{ or } R_{max} \propto \frac{1}{\gamma}$$

2nd order linear eqns

Variation of parameters of a 2nd order eqn

$$(*) \quad y'' + p(x)y' + q(x)y = f(x)$$

Assumes we have the gen. soln of $y_h'' + p(x)y_h' + q(x)y_h = 0$

Assumes we have the gen. soln of $y_h'' + p(x)y_h' + q(x)y_h = 0$

$$y_h = C_1 y_1(x) + C_2 y_2(x)$$

Seek a soln of (*) of the form $y_p = v_1(x)y_1(x) + v_2(x)y_2(x)$

It involves 2 unknown functions
 $v_1(x) + v_2(x)$

We'll pick a convenient constraint between v_1 & v_2 to find a particular soln

$$y_p' = \underline{v_1'} y_1 + v_1 y_1' + \underline{v_2'} y_2 + v_2 y_2'$$

Apply constraint to eliminate underlined terms
 $v_1' y_1 + v_2' y_2 = 0$

$$y_p' = v_1 y_1' + v_2 y_2'$$

$$y_p'' = v_1' y_1' + v_1 y_1'' + v_2' y_2' + v_2 y_2''$$

Plug into (*)

$$y_p'' + p(x)y_p' + q(x)y_p = f(x) = v_1 [y_1'' + p(x)y_1' + q(x)y_1] + v_2 [y_2'' + p(x)y_2' + q(x)y_2] + v_1' y_1' + v_2' y_2'$$

$\begin{matrix} \nearrow \\ \text{0 b/c} \\ \text{y}_1 \text{ solves} \\ \text{homogeneous} \\ \text{eqn} \end{matrix}$

$$\left. \begin{aligned} v_1' y_1 + v_2' y_2 &= 0 \\ v_1' y_1' + v_2' y_2' &= f(x) \end{aligned} \right\} \begin{array}{l} \text{solve these 2} \\ \text{eqns for } v_1' \text{ \& } v_2' \end{array}$$

Then integrate

$$y'' + y = \tan(x)$$

1) Solve $y_h'' + y_h = 0$

$$A \cos x + B \sin x = y_h$$

$$v_1(x) \cos x + v_2(x) \sin x = y_p$$

$$v_1'(x) \cos x + v_2'(x) \sin x - v_1(x) \sin x + v_2(x) \cos(x) = y_p'$$

Assume

$$v_1'(\cos(x)) + v_2'(\sin(x)) = 0$$

$$y_p' = -v_1 \sin(x) + v_2 \cos(x)$$

$$y_p'' = -v_1' \sin(x) + v_2' \cos(x) - v_1 \cos(x) - v_2 \sin(x)$$

$$y_p'' + y_p = v_1 [-\cos(x) + \cos(x)]$$

$$\begin{aligned}
 & +v_2' [m(x) - \sin(x)] \\
 & +v_1' [-\sin(x)] \\
 & +v_2' [-\sin(x)] = \tan(x)
 \end{aligned}$$

$$\begin{aligned}
 v_1' \cos(x) + v_2' \sin(x) &= 0 \\
 v_1' \sin(x) + v_2' \cos(x) &= \tan(x)
 \end{aligned}$$

$$\begin{aligned}
 \sin(x) [v_1' \cos(x) + v_2' \sin(x)] &= 0 \\
 + \cos(x) [-v_1' \sin(x) + v_2' \cos(x)] &= \sin(x) \\
 \hline
 v_2' \sin(x) &\Rightarrow v_2 = -\cos(x)
 \end{aligned}$$

$$v_1' \cos(x) + v_2' \cos(x) = 0 \Rightarrow v_1' = -\frac{\sin^2 x}{\cos x}$$

$$v_1 = -\int \frac{\sin^2 x}{\cos x} dx$$

$$= -\int \frac{1 - \cos^2 x}{\cos x} dx$$

$$= -\int \frac{1}{\cos x} dx + \int \cos x dx$$

$$= -(\ln|\sec(x) + \tan(x)|) + \sin(x)$$

$$\begin{aligned}
 y = y_h + y_p &= [A + v_1] \cos x + [B + v_2] \sin x \\
 &= A \cos(x) + B \sin(x) - \ln|\sec(x) + \tan(x)| \cos(x)
 \end{aligned}$$

$$+ \sin(x) \cos(x)$$

$$- \cos(x) \sin(x)$$

$$y = A \cos(x) + B \sin(x) - \ln|\sec x + \tan x| \cos x$$

Variation of Parameters, Kirchoff's Laws

Tuesday, May 08, 2007
9:04 AM

Variation of Parameters

$$\text{let } y_p = v_1(x)y_1(x) + v_2(x)y_2(x)$$

we know y_1, y_2 solve eqn

$$\begin{aligned} y_1'' + p(x)y_1' + q(x)y_1 &= 0 \\ y_2'' + p(x)y_2' + q(x)y_2 &= 0 \end{aligned}$$

$$y_p = v_1 y_1 + v_2 y_2$$

$$y_p' = v_1' y_1 + v_2' y_2 + v_1 y_1' + v_2 y_2'$$

$$\text{Assume } v_1' y_1 + v_2' y_2 = 0$$

$$\begin{aligned} y_p' &= v_1 y_1' + v_2 y_2' \\ y_p'' &= v_1' y_1' + v_1 y_1'' + v_2' y_2' + v_2 y_2'' \end{aligned}$$

$$\begin{aligned} y_p'' + p y_p' + q y_p &= v_1 [y_1'' + p y_1' + q y_1] \\ &\quad + v_2 [y_2'' + p y_2' + q y_2] \\ &\quad + v_1' [y_1'] + v_2' [y_2'] = f(x) \end{aligned}$$

$$\begin{aligned} v_1' y_1 + v_2' y_2 &= 0 \rightarrow v_1' = -\frac{y_2}{y_1} v_2' \\ v_1' y_1' + v_2' y_2' &= f(x) \rightarrow v_2' \left[y_2' - \frac{y_2}{y_1} y_1' \right] = f(x) \\ &\quad \text{Wronskian } W(y_1, y_2) v_2' \left[\frac{y_1 y_2' - y_2 y_1'}{y_1} \right] = f(x) \end{aligned}$$


$$v_2' = \frac{y_1 f(x)}{W(y_1, y_2)}$$

$$v_1' = -\frac{y_2 f(x)}{W(y_1, y_2)}$$

$$y_p = -y_1(x) \int \frac{y_2(x) f(x)}{W(y_1, y_2)} dx + y_2(x) \int \frac{y_1(x) f(x)}{W(y_1, y_2)} dx$$

[

.. 1



Electric Circuit

Simple relationships between V & current I

$$I = \frac{dQ}{dt} \quad (Q = \text{charge})$$

Simple circuit elmts

Resistor (R) 
 ↖ resistance

$$V_R = IR$$

large resistance $R \rightarrow$ small current I for fixed voltage V

Capacitor (C) 
 ↖ capacitance

$$V_C = \frac{Q}{C}$$

large capacitance $C \rightarrow$ large charge Q for given applied voltage V

Inductor (L) 
 ↖ inductance

$$V_L = L \frac{dI}{dt}$$

Current generates a B field & time varying B field generates an E field

Kirchoff's Laws

(1) Voltage drops round any closed circuit must be zero

$$V = V_R + V_C$$

$$(V - V_C - V_R = 0)$$



(2) Current can't be created or destroyed so total current going into a node or out of a node must be the same

going ...

Parameters are: Applied Voltage V , Resistance R , & C

We want an eqn for V_c

$$V = V_R + V_c$$

$$V_R = IR$$

$$V_c = \frac{Q}{C}$$

we want to eliminate V_R
from the eqn

$$V = IR + V_c$$

$$\frac{dQ}{dt} = \frac{d}{dt}(CV_c) = C \frac{dV_c}{dt}$$

$$V = RC \frac{dV_c}{dt} + V_c$$

$$\frac{dV_c}{dt} = \frac{1}{RC} (V - V_c) ; \quad V_c(0) = 0$$

$$\int \frac{1}{V - V_c} dV_c = \int \frac{1}{RC} dt$$

$$-\ln|V - V_c| = \frac{t}{RC} + A$$

$$|V - V_c| = e^{-\frac{t}{RC} - A} \Rightarrow V - V_c = K e^{-\frac{t}{RC}}$$


$$V_c(t) = V - K e^{-\frac{t}{RC}}$$

$$V_c(0) = V - K e^{-\frac{0}{RC}} = 0$$

$$V - K = 0$$

$$V = K$$

$$V_c = V(1 - e^{-\frac{t}{RC}})$$




$$V_c\left(\frac{1}{2}\right) = \frac{1}{2} V = V(1 - e^{-\frac{T_{1/2}}{RC}})$$

$$\frac{1}{2} = 1 - e^{-\frac{T_{1/2}}{RC}}$$

$$\frac{1}{2} = e^{-\frac{T_{1/2}}{RC}}$$

$$-\frac{T_{1/2}}{RC} = \ln \frac{1}{2} = -(\ln 2)$$

LRC circuit

Want an eqn for V_C = voltage drop across capacitor in terms of the parameters V, R, C, L

$$V = V_R + V_C + V_L$$

$$V_R = I R \quad \text{not } I_R \text{ b/c } I \text{ is } =$$

$$V_C = \frac{Q}{C}$$

$$V_L = L \frac{dI}{dt}$$

$$\begin{aligned} \frac{dV_C}{dt} &= \frac{d}{dt} \left(\frac{Q}{C} \right) = \frac{1}{C} \frac{dQ}{dt} = \frac{I}{C} = \frac{1}{RC} V_R = \frac{1}{RC} (V - V_C - V_L) \\ &= \frac{1}{RC} (V - V_C - L \frac{dI}{dt}) \end{aligned}$$

$$V_C = \frac{Q}{C}$$

$$\frac{dV_C}{dt} = \frac{dQ}{dt} \frac{1}{C} = \frac{1}{C} I$$

$$C \frac{d^2 V_C}{dt^2} = \frac{dI}{dt}$$

$$\frac{dV_C}{dt} = \frac{1}{RC} (V - V_C - L C \frac{d^2 V_C}{dt^2})$$

$$= \frac{1}{RC} V - \frac{1}{RC} V_C - \frac{1}{R} \frac{d^2 V_C}{dt^2}$$

$$\frac{d^2 V_C}{dt^2} + \frac{R}{L} \frac{dV_C}{dt} + \frac{1}{LC} V_C = \frac{1}{LC} V$$

particular soln: $V_C = V$

homogeneous soln $V_C = V_h$

$$V_C = V + V_h$$

$$\frac{d^2 V_h}{dt^2} + \frac{R}{L} \frac{dV_h}{dt} + \frac{1}{LC} V_h = 0$$

$$r^2 + \frac{R}{L} r + \frac{1}{LC} = 0$$

$$r_{\pm} = -\frac{R}{2L} \pm \sqrt{\left(\frac{R}{2L}\right)^2 - \frac{1}{LC}}$$

$$\frac{1}{LC} > \frac{R^2}{4L^2} \Rightarrow \text{decaying oscillations}$$


$$\frac{1}{LC} = \frac{R^2}{4L^2} \Rightarrow \text{critically damped}$$

$$\frac{1}{LC} < \frac{R^2}{4L^2} \Rightarrow \text{overdamped}$$



Systems of 1st order ln eqns

Ex



$$m_1 \frac{d^2 x_1}{dt^2} = F_1 = -k_1(x_1 - a_1) + k_2(x_2 - x_1) - (a_2 - a_1)$$

$$m_2 \frac{d^2 x_2}{dt^2} = F_2 = k_2(x_1 - x_2) - (a_2 - a_1) + k_3((L - x_2) - (L - a_2))$$

equilib

$$\text{Let } x_1 = a_1 + y_1 \Rightarrow y_1 = x_1 - a_1$$

$$x_2 = a_2 + y_2 \Rightarrow y_2 = x_2 - a_2$$

$$\frac{d^2 x_1}{dt^2} = \frac{d^2 y_1}{dt^2}$$

$$\frac{d^2 x_2}{dt^2} = \frac{d^2 y_2}{dt^2}$$

$$m_1 \frac{d^2 y_1}{dt^2} = -k_1 y_1 + k_2(y_2 - y_1)$$

$$m_2 \frac{d^2 y_2}{dt^2} = k_2(y_1 - y_2) - k_3 y_2$$

$$m_1 \frac{d^2 y_1}{dt^2} = -(k_1 + k_2)y_1 + k_2 y_2$$

$$m_2 \frac{d^2 y_2}{dt^2} = -(k_2 + k_3)y_2 + k_2 y_1$$

$$y_1 = \frac{m_2}{k_2} \frac{d^2 y_2}{dt^2} + \frac{(k_2 + k_3)}{k_2} y_2$$

$$m_1 \frac{d^2}{dt^2} \left(\frac{m_2}{k_2} \frac{d^2 y_2}{dt^2} + \frac{(k_2 + k_3)}{k_2} y_2 \right) = -(k_1 + k_2) \left(\frac{m_2}{k_2} \frac{d^2 y_2}{dt^2} + \frac{k_2 + k_3}{k_2} y_2 \right) + k_2 y_2$$

Variables are:

$$y_1, v_1 = \frac{dy_1}{dt}, \quad y_2, v_2 = \frac{dy_2}{dt}$$

$$\begin{bmatrix} \frac{dy_1}{dt} \\ \frac{dv_1}{dt} \\ \frac{dy_2}{dt} \\ \frac{dv_2}{dt} \end{bmatrix} = \begin{bmatrix} v_1 \\ -\frac{(k_1+k_2)}{m_1} y_1 + \frac{k_2}{m_1} y_2 \\ v_2 \\ -\frac{(k_2+k_3)}{m_2} y_2 + \frac{k_2}{m_2} y_1 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 & 1 & 0 & 0 \\ \frac{-(k_1+k_2)}{m_1} & 0 & \frac{k_2}{m_1} & 0 \\ 0 & 0 & 0 & 1 \\ \frac{k_2}{m_2} & 0 & \frac{-(k_2+k_3)}{m_2} & 0 \end{bmatrix}}_M \underbrace{\begin{bmatrix} y_1 \\ v_1 \\ y_2 \\ v_2 \end{bmatrix}}_{\vec{y}}$$

$$\frac{d\vec{y}}{dt} = M\vec{y}$$

Seek solns in the form $\vec{y} = \underbrace{\vec{y}_0}_{\substack{\uparrow \\ \text{const. vector}}} e^{\lambda t}; \quad \vec{y}' = \lambda \vec{y}_0 e^{\lambda t}$

$$\lambda \vec{y}_0 e^{\lambda t} = M \vec{y}_0 e^{\lambda t}$$

$$\lambda \vec{y}_0 = M \vec{y}_0$$

λ is Eigenvalue of matrix

Can a material fail under pure compression? Yes.

Coordination # & Ionic radii.

As cation # gets bigger, coordination # gets bigger
r_{cation} #

Site Selection 2

Stoichiometry

Charge Neutrality (Net charge in structure should be zero)
If all of 1 type of sites is full the remainder have to go somewhere else.

Ex. FCC unit cell has 4 O_H & 8 T_D sites.

Suppose for a specific ceramic: i) cations form FCC sublattice, ii) each unit cell has 6 cations & iii) cations prefer O_H sites.

What fraction of O_H & T_D sites filled?
is

4 anions in FCC

$$1 + 12\left(\frac{1}{4}\right) = 4 \text{ octahedral sites}$$


8 T_D sites

All O_H site: full, 100%

$$2 \text{ cations in } T_D \text{ sites. } \frac{2}{8} = 25\%$$

What would be stoichiometry for above ceramic? What are possible values of ionic charges on each cation/anion?

4 anions / 6 cations



Site Selection 3

Bond Hybridization.

significant covalent bonding



Significant covalent bonding
 - hybrid orbitals can have
 significant impact of significant covalent bond character present.

Predicting the structure of FeO

look up r_{cation} & r_{anion}

Calculate $\frac{r_c}{r_a}$

look up $\frac{r_c}{r_a}$ in relation to coordination # & structure

Rock Salt Structures: Some concepts can be applied to ionic solids in general.


MgO + FeO also have NaCl structure.

Cations prefer O_h sites

AX Crystal Structures

Zinc blende, CsCl

Cesium Chloride



$$\frac{r_{\text{Cs}^+}}{r_{\text{Cl}^-}} = 1.939 \quad \therefore \text{cubic sites preferred}$$

Zinc Blende

ultra-complicated
cube

$$\frac{r_{\text{Zn}^{2+}}}{r_{\text{O}^{2-}}} = 1.529 = O_4??$$

$r_{\text{O}^{2-}}$

No! It's not!

Predicted O_4 but observed T_d .

Due to bonding hybridization,
Zinc prefers T_d

Fluorite structure

another ultra-
complicated diagram

Calcium Fluoride (CaF_2)

Cations in cubic sites

$\text{UO}_2, \text{ThO}_2, \text{ZrO}_2, \text{CeO}_2$

anti-fluorite structure - cations &
anions reversed


c. n. 1

Simple cubic

anions reversed

If you reverse Fluorite, it doesn't work. Works for CsCl, but not for Fluorite, called anti-fluorite.

Perovskite
 BaTiO_3



$\text{Ti}^{4+} \rightarrow \#6$ ●
 $\text{Ba}^{2+} \rightarrow \#12$ ⊗
 $\text{O}^{2-} \rightarrow \#24$ ○

Ceramics are more brittle than metals b/c they have slip systems. too much energy is needed to move one anion past another in ionic, and so slippage is difficult.

Ceramics
$$\rho = \frac{n'(\sum A_c + \sum A_a)}{V_c N_A}$$

always compounds so need to calculate $A_c + A_a$

Metals
$$\rho = \frac{n' A}{V_c N_A}$$

Silicate Ceramics

Most common elmts on Earth are Si & O

SiO_2 structures are quartz, cristobalite, & tridymite

Strong Si-O bond leads to a strong high melting material

We can also form amorphous structures.

A lot of glass is Si-O. - Silica Gels SiO_2 , Si^{4+} & O^{2-} not in well-ordered lattice
 - Charge balanced by H^+ to form OH^- at dangling bonds.
 high surface area $>200 \text{ m}^2/\text{g}$

Carbon Forms

Carbon black-amorphous-surface area ca. $1000 \text{ m}^2/\text{g}$

Diamond

- tetrahedral carbon
- hard - no good slip planes
- brittle - can cut it
- large diamonds - jewelry
- small diamonds
 - often man-made, used for cutting tools & polishing
- diamond films - tools, medical devices, etc. (coatings)

Graphite

- layer structure - aromatic layers
- weak van der Waals forces between layers
- planes slide easily, good lubricant.
- stable form of Diamond

Fullerenes & Nanotubes

- wrap graphite sheet by curving into a ball or tube
- Buckminsterfullerenes (Bucky Ball)
 - Like a soccer ball C_{60} - also C_{70} + others

Defects in Ceramic Structures

Substitutional Impurity

- will substitute for electronically most similar state. Cation for cation site

Interstitial impurity

- impurity must be relatively small compared to anion
- anion interstitials rare.

Frenkel Defect pairs

- cation out of place. cation vacancy + cation interstitial


Shottky Defect pairs

- paired set of cation & anion vacancies.

Equilib conc. of defect pairs $\sim e^{-Q_d/2kT}$

Impurities

Impurities must satisfy charge balance - electroneutrality



Take out 2 Na's to satisfy
 Ca^{2+} w/ Ca^{2+} impurity

Resulting geometry is a cation vacancy.

Ceramic Phase diagrams

Measuring elastic Modulus

Linear Algebra reviews

$$-2x_1 + 3x_2 - x_3 = 4$$

$$-x_1 + x_2 + 2x_3 = 2$$

$$x_1 - 5x_2 + 3x_3 = -1$$

$$\underbrace{\begin{bmatrix} -2 & 3 & -1 \\ -1 & 1 & 2 \\ 1 & -5 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_{\vec{x}} = \underbrace{\begin{bmatrix} 4 \\ 2 \\ -1 \end{bmatrix}}_{\vec{b}}$$

You can only find a soln if the determinant $\neq 0$

If $A\vec{x} = \vec{b}$ has a unique soln
 $\vec{x} = A^{-1}\vec{b}$ if determinant of $A \neq 0$



Consider a homogeneous problem

$$A\vec{x} = \vec{0}$$

A = $n \times n$ matrix

This has a unique soln if $\det(A) \neq 0$ which is
 $\vec{x} = \vec{0}$

There are nonzero (nontrivial) solns to $A\vec{x} = \vec{0}$ only
 if $\det(A) = 0$

$$A = \begin{bmatrix} 1 & 2 & -4 \\ 2 & 1 & 1 \\ -1 & 3 & -11 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 1(-11-3) - 2(-22+1) - 4(6+1) \\ &= -14 + 42 - 28 \\ &= 0 \end{aligned}$$

Since $\det(A) = 0$, then there are
 non zero solns.

$$\begin{bmatrix} 1 & 2 & -4 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & 1 \end{bmatrix} \quad \begin{bmatrix} -1 & 3 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -4 \\ 2 & 1 & 1 \\ -1 & 3 & -11 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -4 \\ 0 & -3 & 9 \\ 0 & 5 & -15 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -3 \\ 0 & 1 & -3 \end{bmatrix}$$

$$\rightarrow \begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{upper triangular form but bottom is all zeros.}$$

2nd row -2 (1st row)
 3rd row $+1$ (1st row)

$$\begin{aligned} x_1 + 2x_2 - 4x_3 &= 0 \\ x_2 - 3x_3 &= 0 \end{aligned}$$

x_3 arbitrary

$$x_2 = 3x_3$$

$$\begin{aligned} x_1 &= -2x_2 + 4x_3 \\ &= -2(3x_3) + 4x_3 = -2x_3 \end{aligned}$$

Soln to $A\vec{x} = \vec{0}$

$$\text{is } \vec{x} = c \begin{bmatrix} -2 \\ 3 \\ 1 \end{bmatrix}$$

Linear Algebra Review

Def. of lin. ind. of vectors $\rightarrow$ A set of k vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ is lin. dependant if there exist constants $c_1, c_2, \dots, c_k$ not all 0 s.t. $c_1\vec{x}_1 + c_2\vec{x}_2 + \dots + c_k\vec{x}_k = \vec{0}$.
 if the only soln to $c_1\vec{x}_1 + c_2\vec{x}_2 + \dots + c_k\vec{x}_k = \vec{0}$ is $c_1 = c_2 = \dots = c_k = 0$ then the vectors are lin. independent.

Let $\vec{x}_1, \dots, \vec{x}_k$ be n -dimensional vectors

if $k < n$, $\Rightarrow$ could either be lin. ind. or dep.

if $k > n$, $\Rightarrow$ must be lin. dep.

Focus on $k=n$

n vectors, n dimensional, $k=n$

Rewrite $c_1 \vec{x}_1 + c_2 \vec{x}_2 + \dots + c_n \vec{x}_n = \vec{0}$ in matrix form as follows

$$\begin{bmatrix} | & | & & | \\ \vec{x}_1 & \vec{x}_2 & \dots & \vec{x}_n \\ | & | & & | \end{bmatrix} \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

$\underline{\underline{X}}$

$$\underline{\underline{X}} \underline{\underline{C}} = \underline{\underline{0}}$$



if $\det(\underline{\underline{X}}) \neq 0 \Rightarrow$ unique soln $\underline{\underline{C}} = \underline{\underline{0}}$

if $\det(\underline{\underline{X}}) = 0 \Rightarrow$ nontrivial soln for $\underline{\underline{C}}$ exists

if $\det(\underline{\underline{X}}) \neq 0 \Rightarrow$ vectors $\vec{x}_1, \dots, \vec{x}_k$ are lin ind.

if $\det(\underline{\underline{X}}) = 0 \Rightarrow$ vectors are lin dep.

$$\det(\underline{\underline{X}}) = 0 \Rightarrow \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ -1 \end{bmatrix} \text{ are lin dep}$$

$$\text{we solved } \underline{\underline{X}} \underline{\underline{C}} = \underline{\underline{0}} \Rightarrow \underline{\underline{C}} = C \begin{bmatrix} -2 \\ 3 \\ 1 \end{bmatrix}$$

$$\text{there exist constants } c_1 \vec{x}_1 + c_2 \vec{x}_2 + c_3 \vec{x}_3 = \vec{0}$$

$$\begin{array}{ccc} & | & \\ -2 & 3 & 1 \end{array}$$

$$-2 \vec{x}_1 + 3 \vec{x}_2 + \vec{x}_3 = \vec{0}$$

1st order

Linear systems of n homogeneous deqs

$$* \frac{d\vec{x}}{dt} = \underline{\underline{P}}(t) \vec{x}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \underline{\underline{P}} \text{ is an } n \times n \text{ matrix}$$

This has a superposition principle:

if $\vec{x}_1(t)$ & $\vec{x}_2(t)$ solve the eqn * then
so does $\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$ for any c_1, c_2

check $\frac{d\vec{x}_1}{dt} = \underline{\underline{P(t)}} \vec{x}_1$ & $\frac{d\vec{x}_2}{dt} = \underline{\underline{P(t)}} \vec{x}_2$

$$\begin{aligned} \frac{d\vec{x}}{dt} &= \frac{d}{dt}(c_1 \vec{x}_1 + c_2 \vec{x}_2) = c_1 \frac{d\vec{x}_1}{dt} + c_2 \frac{d\vec{x}_2}{dt} \\ &= c_1 \underline{\underline{P(t)}} \vec{x}_1 + c_2 \underline{\underline{P(t)}} \vec{x}_2 \\ &= \underline{\underline{P(t)}} (c_1 \vec{x}_1 + c_2 \vec{x}_2) \\ &\quad Q \in \mathbb{R} \end{aligned}$$

Eigenvalues

Monday, May 14, 2007
8:59 AM

Let $\underline{A}$ be an $n \times n$ matrix. λ is an eigenvalue of $\underline{A}$ and $\underline{v}$ is its associated eigenvector if $\underline{A}\underline{v} = \lambda\underline{v}$, $\underline{v} \neq \underline{0}$,
Here λ is a scalar & $\underline{v}$ is an n dimensional vector

$$(*) \quad \frac{d\underline{x}}{dt} = \underline{P}(t) \underline{x}$$

$\underline{x}$ is an n -dim. vector

$\underline{P}$ is an $n \times n$ matrix

The gen. soln of $*$ involves n lin. ind. solns
 $\underline{v}_1(t), \underline{v}_2(t), \dots, \underline{v}_n(t)$

$$\underline{x}(t) = c_1 \underline{v}_1(t) + c_2 \underline{v}_2(t) + \dots + c_n \underline{v}_n(t)$$

Want a simple test for whether $\underline{v}_1, \underline{v}_2, \dots, \underline{v}_n$ are lin. ind.

If we have the gen soln then we know we can solve initial value prob $*$ & $\underline{x}(t_0) = \underline{x}_0$

$$\underline{x}(t_0) = c_1 \underline{v}_1(t_0) + c_2 \underline{v}_2(t_0) + \dots + c_n \underline{v}_n(t_0) = \underline{x}_0$$

Rewrite in matrix form

$$c_1 \underline{v}_1 + \dots + c_n \underline{v}_n = \underbrace{\begin{bmatrix} | & | & & | \\ \underline{v}_1(t_0) & \underline{v}_2(t_0) & \dots & \underline{v}_n(t_0) \\ | & | & & | \end{bmatrix}}_{\underline{V}} \underbrace{\begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}}_{\underline{c}} = \underbrace{\begin{bmatrix} x_{0,1} \\ \vdots \\ x_{0,n} \end{bmatrix}}_{\underline{x}_0}$$

Solve $\underline{V} \underline{c} = \underline{x}_0$ for $\underline{c}$

We can always do this ($\underline{c} = \underline{V}^{-1} \underline{x}_0$), if
 $\det(\underline{V}) \neq 0$

$$\text{Wronskian } W(t) = \det \begin{bmatrix} \underline{v}_1 & \underline{v}_2 & \underline{v}_3 & \dots & \underline{v}_n \\ | & | & | & & | \end{bmatrix}$$

if $W(t) \neq 0 \Rightarrow$ solns are lin. ind.
 if $W(t) = 0 \Rightarrow$ solns are lin. dep.

Can show that the W is always zero or never zero, hence it's enough to check @ 1 pt

$$y'' + ay + by' + cy = 0$$

$$W = \det \begin{bmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{bmatrix}$$

$$y_1 = x_1$$

$$y_1' = x_2$$

$$y_1'' = x_3$$

$$x_1' = x_2$$

$$x_2' = x_3$$

$$x_3' = -ax_3 - bx_2 - cx_1$$

Wronskian w //
be

$$W = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

ex

$$\frac{dx_1}{dt} = 4x_1 - 5x_2$$

$$\frac{dx_2}{dt} = 2x_1 - 3x_2$$

$$x_1(0) = 8 \quad x_2(0) = 5$$

$$\vec{x} = \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}$$

$$\frac{d\vec{x}}{dt} = \begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\frac{d\vec{x}}{dt} = \hat{A} \vec{x}; \quad \vec{x}(0) = \begin{bmatrix} 8 \\ 5 \end{bmatrix}$$

$$\hat{A} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$$

If instead we had just $\frac{dx}{dt} = ax$,
 $x(0) = x_0$
 $x = x_0 e^{at}$

But anyway,

$$\vec{x}(t) = e^{\hat{A}t} \vec{x}_0$$

↑
this is a matrix And we don't really know what that is.

Instead, let's look for a soln $\vec{x} = e^{\lambda t} \vec{v}$ $\frac{d\vec{x}}{dt} = \underline{A} \vec{x}$

$\lambda = \text{scalar}$

$\vec{v} = \text{const. vector}$

$$\frac{d\vec{x}}{dt} = \lambda e^{\lambda t} \vec{v}$$

$$\frac{d\vec{x}}{dt} = \underline{A} \vec{x}$$

$$\vec{v} = \underline{A} e^{\lambda t} \vec{v}$$

$$\boxed{\lambda \vec{v} = \underline{A} \vec{v}} \leftarrow \text{need to solve this eigenvalue prob.}$$

First find eigenvalues $\Rightarrow$ find eigenvectors

Rewrite this eqn

$$(\underline{A} - \lambda \underline{I}) \vec{v} = \vec{0}$$

↑
n x n identity matrix

$$\underline{I} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\underline{I} \vec{v} = \vec{v}$$

Eigenvalue/eigenvector problem $\underline{A} \vec{v} = \lambda \vec{v}$

Find eigenvalues:

$(\underline{A} - \lambda \underline{I}) \vec{v} = \vec{0}$ has a non-zero soln only if

$$\text{Det}(\underline{A} - \lambda \underline{I}) = 0 \leftarrow \text{this is an eqn for } \lambda$$

ex

$$\frac{d\vec{x}}{dt} = \underline{A} \vec{x}, \vec{x}(0) = \begin{bmatrix} 8 \\ 5 \end{bmatrix}$$

$$\underline{A} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$$

Find eigenvalues of $\underline{A}$

$$\text{Det} \left(\begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right)$$

$$= \text{Det} \begin{bmatrix} 4-\lambda & -5 \\ 2 & -3-\lambda \end{bmatrix}$$

$$= (4-\lambda)(-3-\lambda) + 10 = 0$$

$$\lambda^2 - \lambda - 2 = 0$$

characteristic polynomial

$$(\lambda - 2)(\lambda + 1) = 0$$

2 eigenvalues; $\lambda_1 = 2, \lambda_2 = -1$

$$\lambda_1 = 2: \quad A \vec{v}_1 = 2 \vec{v}_1 \Rightarrow (A - 2I) \vec{v}_1 = \begin{bmatrix} 2 & -5 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

$$\lambda_1 = -1: \quad A \vec{v}_2 = -\vec{v}_2 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Find eigenvectors by solving
 $(A - \lambda I) \vec{v} = \vec{0}$

$$2y_1 - 5y_2 = 0$$

$$\vec{v}_1 = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

$$(A - \lambda_2 I) \vec{v}_2 = \begin{bmatrix} 5 & -5 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}; \vec{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$A = n \times n$ vector

λ = scalar (eigenvalue of A)

$\vec{v}$ = n dimensional vector (eigenvector)

$$A\vec{v} = \lambda\vec{v}, \vec{v} \neq \vec{0}$$


$$1) (A - \lambda I)\vec{v} = \vec{0}$$

2) $A - \lambda I$ is "a singular matrix" - cannot be inverted

$$3) \det(A - \lambda I) = 0 \Leftarrow \text{eqn for eigenvalue } \lambda$$

Geometric interpretation of eigenvectors ($A\vec{v} = \lambda\vec{v}$; $A\vec{v}$ & $\vec{v}$ are either parallel or antiparallel provided $\lambda \neq 0$ & λ is real)

$$A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$$




$$A\vec{v} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$$

$$A\vec{v} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ -3 \end{bmatrix}$$

These are not eigenvectors b/c the vectors don't pt. in same direction as multiplier


$$A \begin{bmatrix} 5 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 5 \\ 2 \end{bmatrix} = \begin{bmatrix} 20 - 10 \\ 10 - 6 \end{bmatrix} = \begin{bmatrix} 10 \\ 4 \end{bmatrix} = 2 \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$




eigenvalue eigenvector of A

When you multiply by the matrix you get a vector pointing in same direction

$$A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix} = - \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



$$\underline{\hat{A}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix} = - \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Eigenvalues of the identity matrix $\underline{\hat{A}} = I$

$$\frac{d\vec{x}}{dt} = \underline{\hat{A}} \vec{x} \quad ; \quad \underline{\hat{A}} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$$

$$\vec{x} = c_1 e^{2t} \begin{bmatrix} 5 \\ 2 \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \lambda = -1$$

Check: $\frac{d\vec{x}}{dt} = 2c_1 e^{2t} \begin{bmatrix} 5 \\ 2 \end{bmatrix} - c_2 e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

$$\underline{\hat{A}} \vec{x} = c_1 e^{2t} \underbrace{\underline{\hat{A}} \begin{bmatrix} 5 \\ 2 \end{bmatrix}}_{2 \begin{bmatrix} 5 \\ 2 \end{bmatrix}} + c_2 e^{-t} \underbrace{\underline{\hat{A}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}_{-1 \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \quad \checkmark$$

Def 5 $\underline{\hat{A}} = n \times n$ matrix

$$\det(\underline{\hat{A}} - \lambda \underline{\hat{I}}) = c(\lambda)$$

$c(\lambda)$ is an n th order polynomial in λ called "characteristic polynomial"

$$c(\lambda) = 0 \quad \text{"characteristic eqn"}$$

ex $\underline{\hat{A}} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

eigenvalues & eigenvectors

eigenvalues λ $\det(\underline{\hat{A}} - \lambda \underline{\hat{I}}) = \det \begin{bmatrix} -\lambda & 1 \\ 1 & -\lambda \end{bmatrix} = \lambda^2 - 1 = 0$

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

$$(\underline{\hat{A}} - \lambda \underline{\hat{I}}) \vec{v} = \vec{0}$$

$$\lambda_1 = 1$$

$$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -x+y \\ x-y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\lambda_2 = -1$$


$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x+y \\ x+y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{v}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\hat{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\vec{x} = c_1 e^{-t} \begin{bmatrix} 1 \\ -1 \end{bmatrix} + c_2 e^t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\hat{A} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$$



reflects vectors across $y=x$


ex $\hat{B} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ eigenvalues $\text{Det} \begin{bmatrix} -\lambda & 1 \\ 1 & -\lambda \end{bmatrix} = \lambda^2 + 1 = 0$

$$\lambda_1 = i, \lambda_2 = -i$$

$$\hat{B} \vec{v} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}$$

rotates
vectors
by $\frac{\pi}{2}$

CCW
No real
eigenvectors



every pt is rotated,
⊥ to original.

Complex eigenvalues don't lead to real
eigenvectors → complex eigenvectors

$$e^{i\frac{\pi}{2}}$$

entries in $\hat{A}$ are real

the matrix $\hat{A}$ has real entries

characteristic polynomial $c(\lambda)$ has real coefficients

if $\lambda = a + ib$ is a soln of $c(\lambda) = 0$ (a, b , real #s)
then so is $\lambda^* = a - ib$. λ^* is complex conjugate of λ

if $\vec{v}$ = eigenvector of A associated w/ complex eigenvalue $\lambda = a + ib$
 $\Rightarrow \vec{v}$ is complex

$\vec{v}^*$ = complex conjugate of $\vec{v}$ is the eigenvector for λ^*

ex complex roots to characteristic eqn

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$\det(A - \lambda I) = \det \begin{bmatrix} -\lambda & -1 \\ 1 & -\lambda \end{bmatrix} = \lambda^2 + 1 = 0 \Rightarrow \lambda = \pm i$$

$$(A - \lambda I) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\lambda_1 = i: \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -ix - y \\ x - iy \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\lambda_1 = i; \vec{v}_1 = \begin{bmatrix} i \\ 1 \end{bmatrix} \quad \lambda_2 = -i; \vec{v}_2 = \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

$$\lambda_2 = -i \begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ix - y \\ x + iy \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

For complex eigenvalues $\lambda_1 = a + ib$
 $\lambda_2 = a - ib$, eigenvectors can be
 chosen so that $\vec{v}_2 = \vec{v}_1^*$

$$\frac{d\vec{x}}{dt} = A\vec{x}$$

$$\vec{x} = c e^{it} \begin{bmatrix} i \\ 1 \end{bmatrix} + c^* e^{-it} \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

$c = \text{complex \#} = a + ib$

$c^* = \text{complex conjugate}$

($\bar{c}$ is another notation for c^*, c)

$$\begin{aligned} \vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} &= (a + ib)(\cos t + i \sin t) \begin{bmatrix} i \\ 1 \end{bmatrix} \\ &\quad + (a - ib)(\cos t - i \sin t) \begin{bmatrix} -i \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} ia \cos t - a \sin t - b \cos t - ib \sin t \\ a \cos t + ia \sin t + ib \cos t - b \sin t \end{bmatrix} \\ &\quad + \begin{bmatrix} -ia \cos t - a \sin t - b \cos t + ib \sin t \\ a \cos t - ia \sin t - ib \cos t - b \sin t \end{bmatrix} \\ &= \begin{bmatrix} -2a \sin t - 2b \cos t \\ 2a \cos t - 2b \sin t \end{bmatrix} \end{aligned}$$

$$[2a \cos t - 2b \sin t]$$

$$\vec{x} = A \begin{bmatrix} \sin t \\ -\cos t \end{bmatrix} + B \begin{bmatrix} \cos t \\ \sin t \end{bmatrix} \quad \begin{pmatrix} -2a = A \\ -2b = B \end{pmatrix}$$

Simple test for whether the vectors are real or not:

$$\text{Check that } \vec{x} = \vec{x}^*$$

This only happens if there's no imaginary part.

$$\vec{x} = c e^{it} \vec{v}_1 + c^* e^{-it} \vec{v}_1^*$$

$$\vec{x}^* = c^* e^{-it} \vec{v}_1^* + c e^{it} \vec{v}_1 \quad \checkmark$$

We're solving

$$\frac{d\vec{x}}{dt} = A\vec{x}$$

$$\vec{x}(0) = \vec{x}_0$$

Soln: $\vec{x}(t) = n$ -dim vector

$A = n \times n$ matrix

Case 1

A has n distinct real eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$
with associated eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$

$$\vec{x}(t) = c_1 e^{\lambda_1 t} \vec{v}_1 + c_2 e^{\lambda_2 t} \vec{v}_2 + \dots + c_n e^{\lambda_n t} \vec{v}_n$$

where we choose $c_1, \dots, c_n$ so that $\vec{x}_0 = c_1 \vec{v}_1 + \dots + c_n \vec{v}_n$

Case 2

A has n distinct roots: for each complex conjugate pair $\lambda = a + ib, \lambda^* = a - ib$ there is an associated soln

$$c e^{at} e^{ibt} \vec{v}_1 + c^* e^{at} e^{-ibt} \vec{v}_1^* \quad \begin{array}{l} \text{eigenvector } \vec{v}_1 \\ \text{eigenvector } \vec{v}_1^* \end{array}$$

repeated roots of characteristic eqn

ex 1

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 1-\lambda & 0 \\ 0 & 1-\lambda \end{bmatrix} = (1-\lambda)^2 = 0 \Rightarrow \lambda = 1$$

$$\lambda_2 = 1$$

$$\begin{bmatrix} 1-\lambda & 0 \\ 0 & 1-\lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

repeated
eigenvalue

any vector is an eigenvector

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 \text{ is also an eigenvector}$$

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = 1$$

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

choose these b/c they are lin ind.

ex?

$$\underline{B} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

lower triangular matrix

$$\text{Det}(\underline{B} - \lambda \underline{I}) = \text{Det} \begin{bmatrix} 1-\lambda & 0 \\ 1 & 1-\lambda \end{bmatrix} = (1-\lambda)^2 = 0$$

"defective"

$$\underline{B} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = 1$$

$\lambda = 1$ is a repeated eigenvalue

$$(\underline{B} - \lambda \underline{I}) \vec{v} = \vec{0} \quad ; \quad \lambda = 1$$

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow x = 0$$

$$x \text{ must} = 0 \quad \vec{v}_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

There is only one eigenvector in this problem

$$\vec{v}_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \text{ only one.}$$

Eigenspaces

- The set of all eigenvectors associated w/ an eigenvalue λ is a subspace of our n -dimensional space, called the eigenspace of λ .

Subspaces are lines, planes, origin, anything where you take 2 vectors & their lin. combination is also in that subspace.

Set of all eigenvectors forms a subspace.

Lin combo of 2 eigenvectors w/ same eigenvalue, you get an eigenvector in same eigenspace.

The # of times λ is repeated as a root of $d(\lambda) = 0$ is called the algebraic multiplicity of λ

called the algebraic multiplicity of λ .
The dimension of the eigenspace of λ (# of lin. ind. eigenvectors)
is called the geometric multiplicity of λ .

Phase plane analysis (read Sections 6.1-6.2)

$$\left. \begin{aligned} \frac{dx}{dt} &= F(x, y) \\ \frac{dy}{dt} &= G(x, y) \end{aligned} \right\} (*)$$

2 1st-order autonomous eqns. Don't depend on ind. variable.
 F & G do not depend on t

Plan: (Qualitative) Plotting of soln curves $(x(t), y(t))$ in the (x, y) plane

(x, y) plane = Phase plane

Soln curves can be parametrized by t :

$$(x, y) = (x(t), y(t))$$

The vector field $(F(x, y), G(x, y))$ defines the tangent lines to the soln curves.

Simplest type of curve is a pt. (x^*, y^*) for which $F(x^*, y^*) = G(x^*, y^*) = 0$ Critical pts: (x^*, y^*) of (F, G)

Corresponds to an equilib soln of $*$ since

$$\frac{dx^*}{dt} = \frac{dy^*}{dt} = 0$$

Simple examples

$$\frac{dx}{dt} = x \quad \frac{dy}{dt} = ky \quad k \neq 0$$

$$\begin{aligned} x &= x_0 e^t \\ y &= y_0 e^{kt} \end{aligned}$$

$(0, 0)$ is a critical pt. It is the only critical pt. If $k=0$, then critical pts are $x=0, y=\text{anything}$.

Method 1 for getting $y = y(x; x_0, y_0)$ eliminate t
Solve $x = x_0 e^t$ for t
 $t = \ln \frac{x}{x_0}$ provided $x \neq 0$

$$y = y_0(e^t)^k = \frac{y_0}{x_0^k} x^k$$

Method 2

$$\frac{dx}{dt} = x \quad \frac{dy}{dt} = ky$$

$$\frac{dy}{dx} = \left(\frac{dy}{dt} \right) / \left(\frac{dx}{dt} \right) = \frac{ky}{x}$$

$$\int \frac{dy}{y} = \int k \frac{dx}{x}$$

$$\ln y = k \ln x + C$$


$$y = e^{k \ln x} e^C = A x^k$$

$k > 1$


$k = 1$

$0 < k < 1$


$k < 0$




origin is
unstable
critical pt
"unstable node"



origin is an
"unstable"
node



unstable
node



still unstable
node.

Saddle pt
b/c it is
stable in
one pt

$$m \frac{d^2x}{dt^2} = -kx$$

"phase portrait"

A critical pt (x^*, y^*) of the autonomous system (*) is stable provided that if the initial condition (x_0, y_0) is sufficiently close (x^*, y^*) then $(x(t), y(t))$ remains close to the critical pt (x^*, y^*) for all $t \geq 0$.

Formally, let $\vec{X}(t) = (x(t), y(t))$, $\vec{X}_0 = (x_0, y_0)$, $\vec{X}^* = (x^*, y^*)$

$\vec{X}^*$ is stable provided that for each $\epsilon > 0$, there exists a $\delta > 0$ s.t. if $|\vec{X}_0 - \vec{X}^*| < \delta$, then $|\vec{X}(t) - \vec{X}^*| < \epsilon$

$$\forall t > 0$$


If a C.p. is not stable $\rightarrow$ it is unstable.

If a C.p. is stable and $\vec{x}(t) \rightarrow \vec{x}^*$ as $t \rightarrow \infty$
(provided $\vec{x}_0$ close enough to $\vec{x}^*$) then $\vec{x}^*$ is
asymptotically stable.

$$m \frac{d^2x}{dt^2} = -kx$$

$$\frac{dx}{dt} = y$$

$$\frac{dy}{dt} = -\frac{k}{m}x$$




Tues 11-1 Fri 1-3 (not on wed)

Phase portrait for linear spring

$$m \frac{d^2x}{dt^2} = -kx$$

$$\frac{dx}{dt} = y = \text{velocity of spring}$$

$$\frac{dy}{dt} = -\frac{k}{m}x = \text{acceleration of spring}$$



When velocity is neg,
x is dec. So
it has to be CW,
not CCW.

$$\boxed{\omega < 1}$$

$0 = x^2 = y^2$ is a stable
C.P. but not asymptotically
stable.

We can obtain the phase portrait by considering the eqn for energy conservation: (no damping)

$$\text{total } E = KE + PE = \frac{1}{2} m \underset{\substack{\uparrow \\ v}}{y}^2 + \frac{1}{2} k x^2$$


$$x(t) = A \cos(\omega t - \alpha) \quad ; \quad \omega = \sqrt{\frac{k}{m}} \quad \frac{1}{2} m y_0^2 + \frac{1}{2} k x_0^2 = E_0 = \frac{1}{2} m y^2 + \frac{1}{2} k x^2$$

$$y(t) = \frac{dx}{dt} = -A \omega \sin(\omega t - \alpha)$$

$$x^2 + \frac{y^2}{\omega^2} = A^2 [\cos^2(\omega t - \alpha) + \frac{\omega^2}{\omega^2} \sin^2(\omega t - \alpha)] = A^2 \checkmark$$

With small damping (b),

$$m \ddot{x} = -b \dot{x} - kx$$




$x=y=0$ is an asymptotically stable C.P.

$$\frac{dx}{dt} = F(x, y)$$

$$\frac{dy}{dt} = G(x, y)$$

examine behavior of solns near critical pt $(x, y) = (x_0, y_0)$

$$\Rightarrow F(x_0, y_0) = G(x_0, y_0) = 0$$



Is (x_0, y_0) stable? Unstable? Asymptotically stable?

let $x(t) = x_0 + u(t)$, where $|u_0| < \epsilon$ $u_0 = u(0)$

$y(t) = y_0 + v(t)$, where $|v_0| < \epsilon$ $v_0 = v(0)$

Taylor expand $F(x, y)$ & $G(x, y)$ about (x_0, y_0)

$$F(x_0 + u, y_0 + v) = \overbrace{F(x_0, y_0)}^{0 \text{ b/c } x_0, y_0 = \text{equilib}} + \frac{\partial F}{\partial x}(x_0, y_0)u + \frac{\partial F}{\partial y}(x_0, y_0)v + \dots$$

$$f(x+h) = f(x_0) + f'(x_0)h + \frac{1}{2}f''(x_0)h^2 + \dots$$

$$G(x_0 + u, y_0 + v) = \overbrace{G(x_0, y_0)}^0 + \frac{\partial G}{\partial x}(x_0, y_0)u + \frac{\partial G}{\partial y}(x_0, y_0)v + \dots$$

$$\frac{dx}{dt} = \frac{d}{dt}(x_0 + u) = \frac{du}{dt}, \quad \frac{dy}{dt} = \frac{d}{dt}(y_0 + v) = \frac{dv}{dt}$$

Linearized eqns of motion (neglect nonlinear terms in u, v)

$$\begin{aligned} \frac{du}{dt} &= F_x(x_0, y_0)u + F_y(x_0, y_0)v \\ \frac{dv}{dt} &= G_x(x_0, y_0)u + G_y(x_0, y_0)v \end{aligned} \quad \left| \quad \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \underbrace{\begin{bmatrix} F_x & F_y \\ G_x & G_y \end{bmatrix}}_{J(x_0, y_0)} \begin{pmatrix} u \\ v \end{pmatrix} \right.$$

Jacobian

Example

Example

Jacobian

$$\frac{dx}{dt} = 3x - x^2 - xy$$

$$\frac{dy}{dt} = y + x^2 - 3xy$$

Find all C.P.'s + determine the linearized eqns. near each C.P.

Find C.P.

$$\frac{dx}{dt} = \frac{dy}{dt} = 0 \text{ at } (x_0, y_0)$$

$$3x - x^2 - xy = 0$$

$$y + y^2 - 3xy = 0$$

$$x(3 - x - y) = 0$$

$$y(1 + y - 3x) = 0$$

$$(x_0, y_0) = (0, 0)$$

$$3 - x - y = 0$$

$$1 + y - 3x = 0$$


$$4 - 4x = 0$$

$$x = 1$$

$$(x_0, y_0) = (1, 2)$$

$$(x_0, y_0) = (0, -1)$$

$$(x_0, y_0) = (3, 0)$$




$$J(x, y) = \begin{bmatrix} f_x & f_y \\ g_x & g_y \end{bmatrix} = \begin{pmatrix} 3 - 2x - y & -x \\ -3y & 1 + 2y - 3x \end{pmatrix}$$

$$J(0, 0) = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}$$


$$\frac{du}{dt} = 3u$$

$$\frac{dv}{dt} = v$$



this way b/c it's going faster in the u direction b/c of the $3u$.

C.P. is unstable.




Linearized Eqns

Wednesday, May 23, 2007
9:03 AM

Example

$$\frac{dx}{dt} = 3x - x^2 - xy \quad \frac{dy}{dt} = y + y^2 - 3xy$$

4 C.P.'s $(x, y) = (0, 0), (0, -1), (3, 0), (1, 2)$



$$\frac{dx}{dt} = F(x, y) = F(x_0 + u, y_0 + v) =$$

$$\frac{dy}{dt} = G(x, y) = G(x_0 + u, y_0 + v)$$

C.P. $(x, y) = (x_0, y_0)$

$$\text{let } \begin{cases} x = x_0 + u \\ y = y_0 + v \end{cases} \quad \left. \begin{matrix} \\ \end{matrix} \right\} |u|, |v| \ll 1$$

$$= \cancel{F(x_0, y_0)} + \frac{\partial F}{\partial x}(x_0, y_0)u + \frac{\partial F}{\partial y}(x_0, y_0)v + \dots$$

$$\cancel{G(x_0, y_0)} + \frac{\partial G}{\partial x}(x_0, y_0)u + \frac{\partial G}{\partial y}(x_0, y_0)v + \dots$$

$$\frac{dx}{dt} = \frac{d}{dt}(x_0 + u) = \frac{du}{dt}$$

$$\frac{dy}{dt} = \frac{d}{dt}(y_0 + v) = \frac{dv}{dt}$$

u^2, v^2 etc
much smaller
than other
terms u, v
so ignore.

Linearized Eqns


$$\frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = J(x_0, y_0) \begin{pmatrix} u \\ v \end{pmatrix}$$

↑
Jacobian matrix

$$J(x_0, y_0) = \begin{bmatrix} \frac{\partial F}{\partial x} & \frac{\partial F}{\partial y} \\ \frac{\partial G}{\partial x} & \frac{\partial G}{\partial y} \end{bmatrix}$$


$$J = \begin{pmatrix} 3 - 2x - y & -x \\ -3y & 1 + 2y - 3x \end{pmatrix}$$

$$(0,0): \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$



$$\left. \begin{aligned} \frac{du}{dt} &= 3u \Rightarrow u = u_0 e^{3t} \\ \frac{dv}{dt} &= v \Rightarrow v = v_0 e^t \end{aligned} \right\} u = u_0 \left(\frac{v}{v_0} \right)^3$$

Inc. in u more rapidly than v



blow up of little circle

$$(0,-1): \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 4 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

$$\det \begin{pmatrix} 4-\lambda & 0 \\ 3 & -1-\lambda \end{pmatrix} = (4-\lambda)(-1-\lambda) = 0$$

$$\begin{aligned} \lambda_1 &= 4 \\ \lambda_2 &= -1 \end{aligned}$$

$$\begin{pmatrix} u \\ v \end{pmatrix} = c_1 e^{4t} \vec{v}_1 + c_2 e^{-t} \vec{v}_2$$

this is pos, so any non zero val. of c_1 will make soln blow up. So unstable c.f.

$$(5-\lambda) \vec{v}_1 = \vec{0}$$


$$\begin{bmatrix} 0 & 6 \\ 3 & -5 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 3w_1 - 5w_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{v}_1 = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$$


$$(5-\lambda) \vec{v}_2 = \vec{0}$$

$$\begin{bmatrix} 5 & 0 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 5w_1 \\ 3w_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Saddle pt




pt

$(3, 0)$

$$\frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} -3 & -3 \\ 0 & -8 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

$$\begin{pmatrix} u \\ v \end{pmatrix} = c_1 e^{-3t} \vec{v}_1 + c_2 e^{-8t} \vec{v}_2$$

This is stable. goes to 0.




$$(1, 2) \quad \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} -1 & -1 \\ -6 & 2 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$$

$$\det \begin{pmatrix} -1-\lambda & -1 \\ -6 & 2-\lambda \end{pmatrix} = (-1-\lambda)(2-\lambda) - 6 = \lambda^2 - \lambda - 8 = 0$$

$$\lambda = \frac{1 \pm \sqrt{1+32}}{2}$$

$$\lambda = \frac{1}{2} \pm \frac{\sqrt{33}}{2}$$

saddle pt



Lin. autonomous homogeneous eqn

$$\frac{dx}{dt} = ax + by$$

$$\frac{dy}{dt} = cx + dy$$

$(x, y) = (0, 0)$ is an isolated C.P.


$$(ad - bc \neq 0)$$

Let $\lambda_1, \lambda_2 =$ eigenvalues of $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$

If λ_1, λ_2 have neg. real part $\rightarrow$ C.P. is
asymptotically
stable



If either λ_1 or λ_2 has pos. real
part, C.P. is unstable.

If $\lambda_1, \lambda_2 = \pm i\beta$ ($\beta \neq 0$) stable but
not asymptotically
stable



linear prob w/ $\lambda_1, \lambda_2 = \pm i\beta$

nonlin.



slow spiral in,
S

Stability can often be determined by linearizing (*) about $(x, y) = (x_0, y_0)$

$$\frac{dx}{dt} = F(x, y) \quad \frac{dy}{dt} = G(x, y)$$

$(x, y) = (x_0, y_0)$ is a c.p. if $F(x_0, y_0) = G(x_0, y_0) = 0$

$$x = x_0 + u$$


$$y = y_0 + v$$

$$\begin{pmatrix} \frac{du}{dt} \\ \frac{dv}{dt} \end{pmatrix} = S(x_0, y_0) \begin{pmatrix} u \\ v \end{pmatrix} \quad \left| \quad S = \begin{pmatrix} \frac{\partial F}{\partial x} & \frac{\partial F}{\partial y} \\ \frac{\partial G}{\partial x} & \frac{\partial G}{\partial y} \end{pmatrix} \right.$$

Focus on eigenvalues of matrix about c.p.

eigenvalues of $S(x, y)$	Stability prop. of c.p.	Phase portrait near (x_0, y_0)
$\lambda_1, \lambda_2 < 0$	asymptotically stable node	
$\lambda_1 = \lambda_2 < 0$	asymptotically stable node / spiral	
$\lambda_1 < 0 < \lambda_2$	unstable saddle	
$\lambda_1 = \lambda_2 > 0$	unstable node / spiral	
$\lambda_1, \lambda_2 > 0$	unstable node	
$\lambda_1, \lambda_2 = a \pm ib \ (a < 0)$	stable spiral	
$\lambda_1, \lambda_2 = a \pm ib \ (a > 0)$	unstable spiral	
$\lambda_1, \lambda_2 = \pm ib \ (b \neq 0)$	stable/unstable, center pt or spiral	

Phase portrait for the pendulum





small θ approx $\rightarrow \sin \theta \approx \theta$

$$\ddot{\theta} = -\frac{g}{L} \theta \leftarrow (\text{c.f. lin spring } \ddot{x} = -\frac{k}{m} x)$$

$\dot{\theta} = v$ = ang. velocity

$$\dot{v} = -\frac{g}{L} \sin \theta$$






phase portrait from conservation of energy

$$\text{Energy} = KE + PE \\ = \left(\frac{1}{2} I \omega^2 = \frac{1}{2} m l^2 \dot{\theta}^2 \right) + (mgh = mg(l - l \cos \theta))$$

$$\frac{E}{m l^2} = \frac{1}{2} \dot{\theta}^2 + \frac{g}{l}(1 - \cos \theta) = \frac{1}{2} y^2 + \frac{g}{l}(1 - \cos \theta)$$

Soln curves are level surfaces of constant energy



Example probs

Tuesday, May 29, 2007
9:02 AM

$$\vec{x}' = (\quad) \vec{x}, \quad \lambda = -1, 3, 3$$

$$\vec{v}_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$9a + 11b - 4c = 0$$

$$b = 0 \Rightarrow \begin{pmatrix} 4 \\ 9 \end{pmatrix}$$

$$a = 0 \Rightarrow \begin{pmatrix} 0 \\ 4 \\ 2 \end{pmatrix}$$

$$c = 0 \Rightarrow \begin{pmatrix} 2 \\ -9 \end{pmatrix}$$

#30. 6.2)

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} y-1 \\ x^2-4 \end{pmatrix}$$

Find c.p.'s ; $x' = 0$
 $y' = 0$

$$x' = 0 \quad y-1=0 \Rightarrow y=1$$

$$y' = 0 \quad x^2-4=0 \Rightarrow x = \pm 2$$

c.p.'s $(-2, 1), (2, 1)$



Stability:

$$J(x, y) = \begin{pmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \\ \frac{\partial g}{\partial x} & \frac{\partial g}{\partial y} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 2x & -1 \end{pmatrix}$$

$$J(-2, 1) = \begin{bmatrix} 0 & 1 \\ -2 & -1 \end{bmatrix}$$

$$4 - \lambda^2 = 0$$

$$\begin{bmatrix} -\lambda & 1 \\ -2 & -1-\lambda \end{bmatrix} = 0$$

$$\lambda(1+\lambda)+2=0$$

$$\lambda^2 + \lambda + 2 = 0$$



$$\lambda = \frac{-1 \pm \sqrt{1-8}}{2}$$

Stable spiral

Stable spiral



$$\frac{d^2\theta}{dt^2} + \frac{g}{l} \sin\theta = 0$$



1

$$x' = 2x + 3y + 5 \quad y' = 2x + y - 2t$$

$$\vec{x}' = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix} \vec{x} + \begin{bmatrix} 5 \\ -2t \end{bmatrix}$$

$$\vec{x}_h = A \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} e^{4t} + B \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-t} \quad \begin{array}{l} \text{(soln to} \\ \vec{x}' = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix} \vec{x} \\ \text{eigenval} \\ \text{prob} \end{array}$$

$$\vec{x} = \vec{x}_h + \vec{x}_p$$

$$\vec{x}_p = \begin{bmatrix} a+bt \\ c+dt \end{bmatrix}$$

$$\vec{x}_p' = \begin{bmatrix} b \\ d \end{bmatrix}$$

$$\begin{bmatrix} b \\ d \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} a+bt \\ c+dt \end{bmatrix} + \begin{bmatrix} 5 \\ -2t \end{bmatrix}$$

4 eqns for a, b, c, d
equate coefficients in
front of same types
of terms

$$\begin{bmatrix} 2a + 2bt + 3c + 3dt + 5 \\ 2a + 2bt + c + dt - 2t \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 3 & 0 \\ 2 & 0 & 1 & -1 \\ 0 & 2 & 0 & 3 \\ 0 & 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} -5 \\ 0 \\ 0 \\ 2 \end{bmatrix}$$

t terms
must cancel
out
 $2bt + d - 2 = 0$
 $2bt + 3dt = 0$

$$\vec{x}_{n+1} = \vec{x}_n + \frac{h}{2} (\vec{k}_1 + 2\vec{k}_2 + 2\vec{k}_3 + \vec{k}_4)$$

$$\text{where } \vec{k}_1 = \vec{f}(t_n, \vec{x}_n), \quad \vec{k}_2 = \vec{f}\left(t_n + \frac{h}{2}, \vec{x}_n + \frac{h}{2}\vec{k}_1\right), \\ \vec{k}_3 = \vec{f}\left(t_n + \frac{h}{2}, \vec{x}_n + \frac{h}{2}\vec{k}_2\right), \quad \vec{k}_4 = \vec{f}(t_{n+1}, \vec{x}_n + h\vec{k}_3)$$

$$y'' + y^2 = t \quad y(0) = 0, \quad y'(0) = 0$$

1st step of Runge-Kutta, determine y at next step, $y_1 = y(h)$.

$$y'' + y^2 = t$$

$$w = y'$$

$$\vec{w} = \begin{bmatrix} y' \\ w \end{bmatrix} = \begin{bmatrix} t - y^2 \\ t - y^2 \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} y \\ w \end{bmatrix}, \vec{x}' = \begin{bmatrix} x_2' \\ t - x_1^2 \end{bmatrix} = \begin{bmatrix} x_1' \\ x_2' \end{bmatrix}$$

Plug into Runge Kutta formulas

$$\vec{x}' = \vec{f}(t, \vec{x})$$

$$\vec{f} = \begin{bmatrix} x_2 \\ t - x_1^2 \end{bmatrix}$$

$$y(0) = 0$$

$$y'(0) = 0 \Rightarrow \vec{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{k}_1 = \vec{f}(t_n + \vec{x}_n) \quad \Delta = 0$$


$$= \vec{f}(0, \begin{bmatrix} 0 \\ 0 \end{bmatrix}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\vec{k}_2 = \vec{f}(t_n + \frac{h}{2}, x_n + \frac{h}{2} \vec{k}_1) = \vec{f}\left(\frac{h}{2}, \begin{bmatrix} 0 \\ 0 \end{bmatrix}\right) =$$


Match direction fields to eqns

$$\frac{dy}{dx} = x^2 - y - 2$$

$$\frac{dy}{dx} = y - x + 1$$



$$\frac{dy}{dx} = x - y$$



$$\frac{dy}{dx} = x + y$$

$$\frac{dx}{dt} = x - y \quad \frac{dy}{dt} = x + 3y - 4$$

C.P. (1,1)



A



B

D

$$\frac{dx}{dt} = 1 - y^2 \quad \frac{dy}{dt} = x + 2y$$

C.P. (-2, 1) (2, -1)

A

$$\frac{dx}{dt} = 2x - 2y - 4 \quad \frac{dy}{dt} = x + 4y + 3$$

C.P. (1, -1) B

$$\frac{dx}{dt} = 2x - y \quad \frac{dy}{dt} = x - 3y$$

C

Solve: $\frac{dx}{dt} = x(5-x)$, $x(0) = 3$

$$\text{Ans } \frac{\frac{15}{2} e^{5t}}{1 + \frac{3}{2} e^{5t}} = x$$

$$xy' + 5y = 7x^2, \quad y(1) = 5$$

$$y = x^2 + \frac{4}{x^5}$$

$$y'' - 4y = 2e^{2x}$$


$$y_1 = Ae^{2x} + Be^{-2x}$$

$$y_p = \frac{1}{2}xe^{2x}$$

$$y = Ae^{2x} + Be^{-2x} + \frac{1}{2}xe^{2x}$$

$$x'' + 4x = 5\sin(3t) \quad x(0) = x'(0) = 0$$

$$x = \frac{3}{2}\sin 2t - \sin 3t$$



$$x' = 2x + 3y + 5 \quad y' = 2x - y - 2t$$

$$x_h = A \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} e^{4t} + B \begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{-t}$$

$$x_p = \begin{pmatrix} \frac{1}{8} + \frac{3}{2}t \\ -\frac{2}{4} - t \end{pmatrix}$$

$$x = x_h + x_p$$

$$\frac{dx}{dt} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \vec{x} \quad ; \quad \vec{x}(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\lambda = \{1, 1\}$$

$$\vec{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^t + \begin{bmatrix} t \\ 1 \end{bmatrix} e^{-t}$$

$$\vec{x} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \vec{x}$$

$$(1-\lambda)(1-\lambda) = 0$$

$$\lambda = 1$$

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow$$

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \Rightarrow \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\Rightarrow x = \begin{pmatrix} 1 \\ 0 \end{pmatrix} t + \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^t$$

$$x = A \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^t + B \begin{bmatrix} t \\ 1 \end{bmatrix} e^t$$

$$\vec{x}' = A \vec{x}$$

$$A v_1 = \lambda v_1 \Rightarrow (A - \lambda I) \vec{v}_1 = 0$$

$$A v_2 = \lambda v_2 + v_1 \Rightarrow (A - \lambda I) \vec{v}_2 = v_1$$

Practice Final

$$(4) \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}' = \begin{bmatrix} -\frac{1}{10} & 0 & \frac{1}{10} \\ \frac{1}{10} & -\frac{1}{10} & 0 \\ 0 & \frac{1}{10} & -\frac{1}{10} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad \begin{bmatrix} x_1(0) \\ x_2(0) \\ x_3(0) \end{bmatrix} = \begin{bmatrix} 100 \\ 6 \\ 0 \end{bmatrix}$$

$$(5) \text{ Variation of Parameters ... } u_1 = \int \sin x dx, u_2 = \int -\frac{\sin^2 x}{\cos x} dx$$

$$(6) \vec{x}_{nh} = \begin{bmatrix} h^2/2 & -h^2/46 \\ h^2/6 & \end{bmatrix}$$

- (7) (2, 3) stable spiral
 (0, 0) saddle
 (5, 0) saddle pt

long term behavior $\rightarrow$ 29,000 prey
 30,000 predator