

Linear Algebra

Tuesday, September 19, 2006
7:50 AM

Course webpage: Blackboard www.ea1.mccormick.northwestern.edu

- course info 3 links gen. info.

4 lectures / week Tues - Fri.

Some Mon off, some lab (comp. lab)

MG 47

Quizzes Oct 12 Oct 26 Nov 16

At 7 PM

Final exam on Tues Dec 5 1500-1700

Weekly HW due through course management site

Midnight Wed deadline

There's a course outline & reading info section - WHERE HW

Case studies - Economics, encryption, digital
audio

Course Management Systems

Linear Algebra - Study of vectors and their transformations.

Vector - element of a vector space.

3 equations, 3 unknowns, solve. And how that applies to other stuff.

EA 2 & 3 - More linear Algebra, statics, dynamics,
Physics

Sectors of economy

A: Manufacturing

B: Energy

C: Transportation

Final demand (external)

A: 5 units

B: 10 units

C: 8 units

How much should each sector produce?

Intermediate demand - for a sector to produce their final demand, they must produce units to satisfy each other.

More data is necessary.

To produce 1 unit of A output, we use

	1 A units
input	2 B units

To produce 1 unit of A output, we use

input $\left\{ \begin{array}{l} .1 \text{ A units} \\ .2 \text{ B units} \\ .3 \text{ C units} \end{array} \right.$

Do research on other 2 sectors

$\begin{array}{c} \underline{A} \quad \underline{B} \quad \underline{C} \end{array} \leftarrow \text{output}$

A 0.1 0.1 0.3

B 0.2 0.1 0.06

C 0.3 0.2 0.1

X_A

X_B

X_C

↑
input

consumption matrix

Vectors, Column Vectors

Wednesday, September 20, 2006
6:56 AM

	output					
	A	B	C			
A	0.1	0.1	0.3	} consumption matrix	A	5 units Man.
B	0.2	0.1	0.06		B	10 units Energy
C	0.3	0.2	0.1		C	8 units Transp.
input	↑ intermediate demand					

Unknowns

x_A = # of units produced by A
 x_B " " by B
 x_C " " by C

We want each sector to produce the total demand to maximize the profit.

Production Level = intermediate demand + final demand

$$x_A = 0.1x_A + 0.1x_B + 0.3x_C + 5$$

$$x_B = 0.2x_A + 0.1x_B + 0.06x_C + 10$$

$$x_C = 0.3x_A + 0.2x_B + 0.1x_C + 8$$

Linear equations b/c they form straight lines, and this is b/c there are no exponents greater than 1

Vector - Ordered list of numbers

The order matters - 3, 5, 9 is not the same as 5, 9, 3

Column Vector

$$\begin{bmatrix} 3 \\ 5 \\ 9 \end{bmatrix} \begin{bmatrix} 9 \\ 5 \\ 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

$\mathbb{R}^3$ } ordered list of 3 real #s
 $\mathbb{R}^2$
 $\mathbb{R}^4$
 belong to $\mathbb{R}^3$
 Real #s

We can multiply vectors by scalars

Scalar Multiplication

$$3 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}$$

↓
Scalar

Vector Addition

$$\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix} = ?? \text{ Not DEFINED}$$

Linear combination

$$3 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

coefficients, "weights"

put weights into a vector

$$\begin{bmatrix} 3 \\ 4 \\ 2 \end{bmatrix}$$

Same computation:

$$\begin{bmatrix} 1 & 0 & 1 \\ 2 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

Matrix

Vector

Multiplication

A

b

Ab

of columns of A must be the same as the # of entries in b.

The matrix is always first, on the left

$$\begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = ?? \text{ MAKES NO SENSE}$$

Final DEMAND

$$\begin{bmatrix} 5 \\ 10 \\ 8 \end{bmatrix} \in \mathbb{R}^3$$

Production Levels

$$X = \begin{bmatrix} x_A \\ x_B \\ x_C \end{bmatrix}$$

Consumption Matrix

$$C = \begin{bmatrix} 0.1 & 0.1 & 0.3 \\ 0.2 & 0.2 & 0.06 \\ 0.3 & 0.2 & 0.1 \end{bmatrix}$$

rows
3x3 3

$$X = \underbrace{CX}_{\text{intermediate demand}} + d$$

$$X - CX = d$$

Matrix Addition

$$\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 3 & 2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ 2 & 0 \end{bmatrix}$$

We need to write x as $A_1 x$. C would be A_2 .

$$(A_1 + A_2)b = A_1 b + A_2 b$$

$$x = Cx + d \quad \text{HW on Blackboard due Wed}$$

Cx = linear combination
Matrix(vector)
Result = intermediate demand
 $x, d \in \mathbb{R}^3$
vectors in $\mathbb{R}^3$

$$x = \begin{bmatrix} x_A \\ x_B \\ x_C \end{bmatrix} \text{ unknowns}$$

$$d = \begin{bmatrix} 5 \\ 10 \\ 8 \end{bmatrix}$$

C is 3×3 matrix

Matrix $x - Cx = d$

vector $x = x_A \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + x_B \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_C \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

$$(I - C)x = d$$

$$x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} x$$

identity matrix "I"

$$\boxed{Ax = b}$$

Standard form

$$\boxed{\begin{matrix} A \text{ is } m \times n \\ x \in \mathbb{R}^n \\ b \in \mathbb{R}^m \end{matrix}}$$

b is a linear combination of A , so it has m entries since A has m rows.

Matlab !!! Start playing w/ it.

Variable $\begin{bmatrix} x \\ 0 \\ 5 \\ 2 \end{bmatrix}$

$$\begin{bmatrix} x \\ 0 \\ 5 \\ 2 \end{bmatrix}$$

In Matlab, we create variables

Names start w/ letters

Case sensitive

this _variable_ works. But that's a long name for a variable.

assignment statements

~

$$x = 3$$

variable on left

data on right

$$x = 3 + x$$

$$\begin{bmatrix} x \\ 3 \end{bmatrix}$$

this assignment says "take the ~"

$3 = x$ ERROR!

$$x = 3 + x$$

$$\overset{x}{\boxed{3}}$$

This assignment says "take the previous variable and add 3"
So if the variable was 3, the new variable would be 6.

$$\text{Also } x = x + 3$$

Command Window is where you can enter in commands 1 by 1 and Matlab will execute them.

`CLEAR` variable name it gets rid of the variable.

`CLEAR ALL` gets rid of everything

* means multiplication

^ means exponent

; means Matlab won't tell you that it got to the command, but it will carry out what it was told.

you don't need spaces, it just makes it easier to read

$x = [3; 5; 2]$ tells Matlab that x is a vector

$x = [.1 .1 .2; .3 .0 .1; .3 .11 .03]$ 3x3 Matrix

$i = [1 \ 0 \ 0; 0 \ 1 \ 0; 0 \ 0 \ 1]$ identity matrix.

OR

$i = \text{eye}(4)$ gives us a 4x4 identity matrix

Colon Operators, Subarrays

Friday, September 22, 2006

7:02 AM

Quiz Options

Oct. 26th

Thurs

Oct. 12th

is another quiz

Nov. 16th

is yet another

5 - 6:30 PM

6 - 7:30 PM if Chem moves to 8 PM

8:30 - 10 PM X - hockey, doesn't work

Input

Data representation

1x1 is a scalar, and is the smallest unit in Matlab
same as 4 or ['a']

You give names to scalars & matrices, and those are variables

Functions - later. Solve Prob.

Display/Save results

- integer values displayed as integers.

- doubles displayed as w/ 4 decimal digits

2.45678

2.45678

- doubles that are too large or too small are displayed in scientific notation.

Assignments.

radius = 15.5 you don't have to declare variables

radius_copy = radius;

circumference = 2*pi*radius pi is predefined.

Predefined variables

pi; ans (stores result); i, j (sqrt(-1)); Inf (infinity)

NaN (doesn't work); eos; date; clock.

More assignments

x(1,2) = 5; % partial def. - all other values filled w/ 0

x

1	2

y(4) = 0.1

Since y is already assigned, say as

1	2
3	4

 it will replace

row-major - we store the #1 in the 1st row, then #2, #3, #4, #5 & 6
column major - means #1 in the column, then #2, #3, #4, #5, 6
 The matrix we specified is column major.
 It will replace the 2 with 0.1.

$$\text{data} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

$\text{data}(2) = 9$, changes the 4 into a 9.

$d2 = [\text{data} \text{ data}]$ takes 2 copies of the matrix, puts them side-by-side, and makes one big matrix

$d3 = \text{data}'$ % transpose $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^T$ transpose does this: $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$

$\text{dir } 1 = [2 \ 3]'$ shorthand

$\text{dir } 2 = [2 \ 3]$ % same as $\text{dir } 1$

$I = \text{eye}(4)$ 4x4 identity matrix

$I = \text{eye}(2, 4)$ $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$

$>> \text{ones}(4, 5)$
 $\begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}$

format compact means save a few lines.

$\text{size}(\text{ans})$ gives you the size of the matrix

if you did $x = \text{ones}(4, 5);$

$y = \text{zeros}(\text{size}(x))$ So you can be lazy.

Colon Operator

Shorthand notation that allows you to generate

$d = 1:3$

you get $d =$

1 2 3

$d = a:c$

you get $d =$

#s between a & c

$d = 'a': 'c'$

you get $d =$

abc

$\text{size}(d) = 1 \quad 3$ b/c matlab thinks abc is a 1×3 matrix
 $x = (1:0.1:2)^* 2$ means you take everything between 1, 0.1, & 2 and make a matrix, then multiply by 2

$y = [x; x*2]$ Matrix

$x = 1:0.1:2; \quad y = [x; x*2];$
 $\downarrow$

1	1.1	1.2	1.3	
2	2.2	2.3	2.4	...

Subarrays You can use the colon operator to access chunks of what you're working on.

$x = (1:3)$ is $(10, 20, 30)$ if x originally was $[10, 20, 30, 40]$
 $x = (1:2:4)$ is $(10, 30)$
 $x = (1:)$ means get everything in the 1st row

- Way to enter data: shortcuts, :

- Way to refer to data so we can use it in calculations
- Way to perform the calculations
- Way to display results

$$x = Cx + d$$

$$x - Cx = d$$

$$(I - C)x = d$$

$$\text{eye}(\text{size}(C))$$

$$(I - C)^{-1} (I - C)x = (I - C)^{-1} d$$

$$Ix = (I - C)^{-1} d$$

$$x = (I - C)^{-1} d$$

`x = input('please enter number x:');`

This prompts the user to type something in, and then assigns what they type to `x`.

`x = input('please enter table x:');`

`name = input('please enter your name:');`

MAT-file preserves variable names, types, sizes
can only be read by
MATLAB typical ext = .mat

ASCII file data saved as ASCII strings separated by spaces.

variable info lost

can be read by any program/human

Loading files: `load filename.mat`

`load filename.dat` ASCII

`load filename.mat x y` only those variables

$\text{inv}(x)$ is the inverse of x .

array operations & matrix operations

↓
.. ..

Do not always produce

element by element basis
& between matrix & scalar

the same result.

$$? \begin{bmatrix} 0 & 1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} ?0 & ?1 \\ ?2 & ?3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 0 & 4 \end{bmatrix} \quad \text{Array operation}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 3 & 4 \end{bmatrix} \quad \text{Standard multiplication.}$$

$$\begin{aligned} (1 \cdot 1) + (2 \cdot 3) &= 7 && \text{1st row 1st column} \\ (1 \cdot 2) + (2 \cdot 4) &= 10 && \text{1st row 2nd column} \\ (0 \cdot 1) + (1 \cdot 3) &= 3 && \text{2nd row 1st column} \\ (0 \cdot 2) + (1 \cdot 4) &= 4 && \text{2nd row 2nd column} \end{aligned}$$

$$(n \times m)(m \times p) = n \times p ?$$

* is array multiplication
* is multiplication.

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + 1 = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 1 \\ 7 & 5 \end{bmatrix} - \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} 2 & -2 \\ 3 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \cdot * 2 = \begin{bmatrix} 2 & 4 \\ 6 & 8 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 1 \\ 7 & 5 \end{bmatrix} \cdot * \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} 8 & 3 \\ 28 & 25 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \cdot / 2 = \begin{bmatrix} .5 & 1 \\ 1.5 & 2 \end{bmatrix}$$

$$2 ./ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ .67 & .5 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 1 \\ 7 & 5 \end{bmatrix} ./ \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} 2 & .3 \\ 1.75 & 1 \end{bmatrix}$$

$$A / B = A(B^{-1})$$

Standard

$$A \setminus B = B / A \quad \text{array}$$

or

$$A \setminus B = B / A \quad \text{matrix}$$

So we have soln. $x = \text{inv}(\text{eye}(\text{size}(C)) - C) * d$
disp(x) shows x.

msg = ['The production vector is']

disp(mg50); disp(x)
the production vector is

number 1 x_m

number 2 x_A

number 3 x_s

↙ # of spaces

fprintf('Agr production is %4.2f'), x(1)

↑ # of x's displayed like

fprintf('%4.2f, %4.2f', x(1), x(2)) 1.22 instead of
1.2257987

num2str: is how you create a string.
 floor(x) is the biggest integer smaller than x.
 ceil(x) is the smallest integer larger than x.
 cos(x), sin(x), sqrt(x)

max(4 6 6) gives you the largest # in a matrix.
 = 6

plotting w/ Matlab - get data, (like record highs + lows for Sept.)
 legend needed for once.

2 plots on the same figure
 Diff color for each one.

- Create 2 vectors - one for x values, one for y values.
 - Must have same length.

- "plot" is the function

plot(x, y) $x.^2$ b/c 2 is a scalar.

plot y as a function of x x is a vector

w/ linespec being a string

that describes the style (e.g. dotted), color, & marker
 symbol of the lines. "plot(x, y, linespec)"

eg. $x = [-5:1:5]$; $y = x.^3$; plot(x, y, 'bo--')
 b for blue o for circular markers, -- for dashed
 line.

- "plot(x1, y1, linespec1, x2, y2, linespec2)"

plot both functions in the same figure.

- "plot(... 'PropertyName', PropertyValue, ...)"

properties ex. line width, etc.

plot(x, y1, 'bo--', x, y2, 'k*:', 'linewidth', 2);

You can't have 2 'linewidths' in the same
 command. This is not the method for putting
 properties in the same figure.

- "loglog" is similar to "plot", but used for exponential
 functions.

- "xlabel('some text')"

- "ylabel('some text')"

- "legend('explanation for x', 'exp for y')"

- title ('some text')

- toggle grid lines by 'grid on' or 'grid off'

M-files are files you upload, usually .txt files, and Matlab executes the files for you. Could be typical script or function.

.mat file is a Matlab file, can only operate using Matlab.

load "name.mat"

"hold;" means for the file to stay, for the graph to not be deleted and for Matlab to plot your following line on the same graph.

legend ('high', 'low')

legend ('high', 'low', 0)
↑ position.

"grid" gives you a grid.

xlabel ('days'), ylabel ('temp'),

To Matlab help
'Axes' or 'Axis'
tells you about this.

title ('Highs and Lows. Source: weather.com')

↑
we want to create a 2nd line
here. So insert ' on end of 'Highs and Lows'
and ①, and 10① and insert ' on beginning
of 'Source...' 10 is 'next line'.

And insert brackets on inside of ().

to save this variable information

"save data.mat specific-variable1 specific-variable2"

or just this ↵

'%d' is for integers

x_0 - start off w/ this

$$x = \underbrace{C x}_{\text{intermediate demand}} + d \quad \text{external demand}$$

NO equilibrium if x on right $= x_0$. $x^{(0)}$ gives you $x^{(1)}$

Now if you want to approximate again, you get

$$x^{(2)} = C x^{(1)} + d$$

$$x^{(3)} = C x^{(2)} + d$$

$$x^{(4)} = C x^{(3)} + d$$

$$\boxed{x^{(n)} = C x^{(n-1)} + d}$$

Iterative Method

Execute 1 + calculations repeatedly until some condition has been satisfied.

you end up w/
an approximation that's
very close to the real
thing.

$$x^* = C x^* + d$$

compare x 's - if the diff
between them is really small
they are just about the same.

↓
print results

$x = \text{input}(\quad)$

$y = \text{input}(\quad)$

do it again ← if not small enough

if y is 0, then fprintf("division by zero not allowed")
otherwise, (x / y) .

$x ==$

check whether
 x has the same value
as y

$1 == 5$

$0 = \text{false}$ in Matlab.

$'a' == 'a'$

$1 = \text{true}$

$[10 \ 1 \ 2] == [10 \ 3 \ -2]$

$[1 \ 0 \ 0]$ because 1st elmt. is
true & other 2 are false

$'a' == 'A'$

If/Then - Sequel to Either/Or?
Knowledge = Power

$x \sim y$ test inequality
 0 = equal
 1 = not equal

Boolean values
 True / False values.

$x < y$ less than
 $x \leq y$ less than/equal to
 $x > y$ greater than
 $x \geq y$ greater than/equal to

relational operators.

Allow us to create simple expressions

if to day == 'Sunday'
 then weather == 'sunny'
 else have a picnic
 sleep in.

we want to combine them
 They both have boolean values.

short circuit evaluation

logical operators — && : and
 ↓
 supports short circuit evaluation
 ONLY WORKS FOR SCALARS

$(x \leq 5) \&\& (y == 10)$
 T && T → T, 1
 everything else → 0

Logical operators — & : and
 & : and

if t = Sunday && w = sunny
 then picnic
 else sleep

$x \parallel y$
 $x \mid y$
 if both x, y are 0 →
 result 0
 all other cases →
 result 1

if t = Sunday | w = sunny
 then picnic
 else sleep

diff is that if it's Sun, then we
 have a picnic no matter what the
 weather is. But if it's false, we
 check the weather and we get
 a picnic if it = sunny.

Xor inclusive or
 xor (x, y)
 this is the Either/Or
 operator.
 "Xor"


x	y	result
1	0	1
0	1	1
0	0	0
1	1	0

~ NOT operator → unary operator
x, ~x the opp. of x

if (~('tank = full')) if the () expression is true, the ~
fill up negates it. same as if (tank ~='full')
fill up.

Relational: ==, ~=, <, <=, >, >=
Logical &, &, ||, |, xor, ~

If/Then book



x || y & 8 < z < 4 ||
x + y * z PEMDAS
x + y - z left-right

ischar(A) is character
isfloat

isinteger

isinf is infinity

isnumeric(A) array/matrix

logical(x) convert # into boolean.

4 || 0

True - 1, b/c if it's
not 0, it's true.

If, Else, End

Friday, September 29, 2006
7:01 AM

$$f(x) = \begin{cases} x^2, & \text{if } 1 \leq x \leq 10 \\ x-1, & \text{other value} \end{cases}$$

`x = input('please enter x')`

```
if 1 <= x <= 10
    true / \ false
    y = x^2   y = x-1
           \ /
            disp(x)
```

```
if x >= 1 & x <= 10
    y = x.^2
else
    y = x-1
end
disp(y)
```

```
if score > 90
    grade = 'A'
elseif score > 80
    grade = 'B'
elseif score > 70
    grade = 'C'
else
    grade = 'F'
end
```

equivalent of having more if statements in the else statement.

IF condition

stat1
stat2
stat3

end

```
if tank = 'empty'
    true / \ false
    fill up   end
              drive
```

NUMBERS

SWITCH

```
if x == y1
    then stmt1
elseif x == y2
    then
        stmt2
elseif x == y3
    then
        stmt3
else
    ...
```

→

```
switch(x)
case y1
    stmt1
case y2
    stmt2
case y3
    stmt3
otherwise
    stmt4
```

x must be
a character
for this
to work.
or a #.


```


else
    stmt4
end
end
end
end

```

Back to plotting

figure(1) go on w/ plot commands > same window
 figure(2) puts another plot in.

subplots  2x1 table.

subplot(2,1)  2x2 table
 subplot(2,2,4)
 plot(...)

axis([x_{min}⁰ x_{max}³¹ y_{min}⁵² y_{max}¹⁰⁰]) tells you to start w/ 4th quadrant. Not like math.
 default: [0 0 0 0] specify differently.

xlabel('Day') xlabel('\bf\fontsize{12} Day')

The program sometimes calculates things incorrectly even though they look right. The easiest way to fix it is to do it on paper, but the best way to do it is to get Matlab to figure it out for you. If you tell Matlab to do your code line-by-line, you can find the statement that caused the stray value.

Debugger 'step'
 step in (a function)
 step out

Breakpoints - execute everything up to a point and then wait

M-file - you can create a file and then insert it into Matlab. It'll work only on matlab. You have to click on New → M-file. Then save

Expression is anything that has a value. $\sin(x)$ is an expression. If you're asked to write an expression about 2 matrices, you can make them a & b and leave it at that - you don't need a specific a or b .
 $A == B$ is usually found in if/then statements.

$$\begin{aligned} X &= C * X + d & X^0 &= d \\ X^1 &= C * X + d \\ X^2 &= C * X^1 + d \\ X^3 &= C * X^2 + d \\ &\text{until } X^n = X^{n-1} \end{aligned}$$

for → when you know how many times you want to iterate. use counter.
 while → event based. we don't know how many times it'll iterate. wait for an event to exit the loop.

for $\text{index} = \text{expr}$ ← typically first : incr : last
 statement 1
 statement 2 } body of loop
 statement n
 end

ex.

```
for ii = 1:2:10
    disp(ii);
end
```


displays

1
3
5
7
9

always a scalar in this example

ex.

for $v = [1:3; 4:5]$



line it all up nicely.

Never mess
w/ v inside
the loop like
this →

```
disp('-----');
disp(v);
v = v, * 2;
disp('changed to');
disp(v);
```

displays →

```
-----
  1
  4
changed to
  2
  8
-----
  3
  5
changed to
  6
 10
```

while loops

```
while expr
    statement1
    statement2
    statement n } body of loop
end
```

load econ_data.mat

$x = [10; 10; 10; 10]$ % initial estimate

error = 1; % initialization

while (abs(error) > 10^{-5})

old_x = x;

$x = C * x + d$;

error = old_x - x;

end

disp('The production vector is: ');

disp(x);

(if we picked $d = x$, and make error =

old_x = zeros(size(x));

thresh = input(...);

if you put
iterations = 500,
then you can
change while
to obey the
amount of iterations

```
while (abs(old_x - x) > thresh)
    old_x = x
    x = C * x + d
end
```

while (abs(old_x - x) > thresh & iterations < 500

old_x = x;

$x = C * x + d$

↑

counter

$x = C * x + d$
 iterations = iteration + 1
 end

this is a counter.
 Value doesn't change
 magically. You need
 to update x.

A break statement stops the loop completely.

```

myguess = ceil(6 * rand(1));
while 1 % 1 means true, & it'll never change.
    x = input('Guess a number: ');
    if (x == myguess)
        disp('You found it!');
        break; % means -
    else
        disp('Sorry, that's not it. ');
    end
end
  
```

A continue goes to the end of the statement, but goes back through the loop again.

Never test for equality between floating point #s.

```

x = .1;
while (x ~= 1)
    disp(x);
    x = x + .1;
end
  
```

displays →

0.1
 0.2
 0.3
 0.4
 0.5
 0.6
 0.7
 0.8
 0.9
 1.0
 1.1
 1.2
 1.3
 ...

Wasteful

```

ii = 1:5
data(ii) = some-value
end
  
```

better

```

data = zeros(5);
for ii = 1:5
    data(ii) = some-value
end
  
```

eek!
 it'll go
 forever!

size, then $A(B)$ returns the values of A whose corresponding index in B is 1. The values are returned in COLUMN major format, not in row major format as the lecture notes and our class example implied. For example:

$A = \begin{bmatrix} 10 & 7 & 20 \\ 19 & 9 & 18 \end{bmatrix}$

If $B = A > 10$ then B is

$\begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$


There are ones at indices 2, 5 and 6.

$A(B)$ is $\begin{bmatrix} 19; \\ 20; \\ 18 \end{bmatrix}$

Vana

Arrays, solving probs

Wednesday, October 04, 2006
6:55 AM



data(1) = 19
data(2) = 38
data(3) = 54

We should avoid
enlarging an array
inside the loop
To avoid this, tell
it you need 1000
array.

ex. tic; Sets stop watch
for k = 1:500
for m = 1:500
data(k,m) = 1;
end
end
elapsed_time = toc; tells us how much time has passed
fprintf('Elapsed time with no preallocation: %6.4f seconds\n', elapsed_time);
clear all;
tic;
data = zeros(500);
etc

```
1 x 10000  
for k = 1:10000  
    CV(k) = cos(v(k))  
end
```

} takes too much time

cos(v) in beginning is better.

preallocating is always better.

index arrays

```
arr(1:3)  
arr([1 5 9])  
arr(2:end, 1:end)
```

Find out what elements of an array satisfies some property.

$$\begin{bmatrix} 1 & 0 & -2 \\ 6 & -4 & -5 \end{bmatrix}$$

$A > 0$

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \leftarrow \text{logical array}$$

$A(B)$ will extract the elements that gave you a true (1)

$$\begin{bmatrix} 1 \\ 6 \end{bmatrix}$$

And if it were $\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix}$

then you get $\begin{bmatrix} 1 \\ 4 \end{bmatrix}$

$B = A > 0$

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

You can determine if any elements exist that satisfy the property.

$$\begin{bmatrix} 1 & -2 & 4 \\ -1 & 3 & 9 \end{bmatrix}$$

$A < 0$

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$\text{negs} = A < 0$

if any($A(\text{negs})$)

then...

$A(\text{negs})$

$$\begin{bmatrix} -2 \\ -1 \end{bmatrix}$$

A ball is thrown from position (0,0) w/ $v_0 = 20 \text{ m/s}$ at $\theta = 0^\circ$. Write program that plots trajectories for the ball for all θ from 5° to 45° in 10° steps. Program should also determine horizontal distance traveled by ball before it reaches the ground again for all θ from 0° to 90° in 10° increments. Also determine θ that maximizes range of the ball, a plot that trajectory in a diff color w/ a thicker line

top $\rightarrow$ down

evaluate ranges

plot ranges

$\rightarrow$ determining θ for max range.

evaluate trajectories

plot trajectories

You don't write the whole program in one go.

$$v_0 = 20 \text{ m/s}$$


$$v_x = v_0 \cos \theta \quad v_y = v_0 \sin \theta$$

$$y = \frac{v_y}{g} t - \frac{1}{2} g t^2$$

$$V_0 = 20 \text{ m/s}$$

$$V_x = V_0 * \cos \theta \quad \left. \begin{array}{l} V_y = V_0 * \sin \theta \end{array} \right\} \text{velocity}$$

$$V_y = V_0 * \sin \theta$$



$$y(t) = y_0 + V_{y0}t + \frac{1}{2}gt^2$$

$y_0 = 0$ at this problem

$$y(t) = V_{y0}t + \frac{1}{2}gt^2$$

$$x(t) = x_0 + V_{x0}t$$

$x_0 = 0$ at this problem

$$x(t) = V_{x0}t$$

$$\text{radians} = \text{degrees} * (\pi / 180)$$

We want to calculate the ranges (the x's) from $0^\circ - 90^\circ$

$$0 \quad 1 \quad 2 \quad \dots \quad 90$$

$$x(t) = V_{x0}t$$

$$\text{we know } 0 = 0 + V_{y0}t + \frac{1}{2}gt^2$$

$$V_{y0} = V_{y0} * \sin \theta$$

$$V_{y0} = 2 * \sin \theta$$

$$\frac{y = -\frac{2V_{y0}^2}{g}}{g} \rightarrow V_{x0} = 2 * \cos \theta$$

$$x = V_{x0}t$$

In Matlab-English hybrid:

$$g = 9.81$$

$$V0 = 20$$

$$\text{angles} = 0 - 90$$

$$V_y = V0 * \sin(\text{angles}, * \text{ratio})$$

$$V_x = V0 * \cos(\text{angles}, * \text{ratio})$$

Ranges for all angles (sample prob)

Thursday, October 05, 2006
7:00 AM

Back to yesterday's problem: Find ranges for all $45^\circ - 90^\circ$

angles = 0:90;
rads = angles .* (pi/100); % converts to degrees

v0_x = v0 .* cos(rads);
v0_y = v0 .* sin(rads)

time-to-land = (-2/y) .* v0_y;
range = v0_x .* time-to-land;

Now we want a table:

angle	range
0	0.00
1	8.48
2	—
...	...
90	—

So: fprintf(*\n Angle
for (k=1:91)
fprintf(*%2d %8.4f\n', angles(k), range(k));

Annotations:
- * is for floating point
- %2d is integer
- %8.4f is 8 spaces, 4 decimal digits
- \n is new line

Now we want to find & print angle for max range
Because we set up angle(0) = range(1), angle(1) = range(2)
We can find the Δ.

[maxrange index] = max(range);
fprintf('Max range %6.2 fm at %d degrees\n\n')

Now we want trajectory
For each θ , find out where it was at time t
So we use a for loop.

for angle = 5:10:85
rad = angle * pi/180;
v0_x = —
v0_y = —
time-to-land = —

time-interval = time-to-land/99;
time = 0:time-interval:time-to-land
... x = v0_x * t ...

? How long will it take to land?
? split in intervals - for each moment in time, find location of the ball.
Save to a table



```

current_x = v0_x * t * times;
current_y = v0_y * t * times;
if angle <= max_int_x - 1
    plot(current_x, current_y, 'linewidth', ...

```

```

else
    plot(current_x, current_y, ' - - -')
end
hold on;
% we want to plot everything on
% the same figure, so if you put
% that there, it'll start in a new
% window next time. You don't
% want to lose the previous
% trajectory.
end
plot
end
plot
end

```


end
hold off;

A double every 60 min

B double every 90 min

BACTERIA

Examine how they double, save info about their #s
every 3 hours. Plot growth.



x: 0 1 2 3 4 5 6 ... 24
y_A y₁ y₂ y₃
y_B y₁ populations at x times (hrs)

for 0 min ... 24 * 60 min
note: if it's a multiple of 60 min,
then A reproduces.
else if it's a multiple of 90 min,
then B reproduces.

else if it's a multiple of 3 hours
then save populations in our table.

end

plot.

rem () remainder function == 0
then double population.
check every 30 min, = increment.

num A = 1;
num B = 1;

```
for hicker = 30 : 30 : 600 - min
```

```
if rem(hicker, 60) == 0
```

```
    numA = numA * 2;
```

```
end
```

```
if rem(hicker, 90) == 0
```

```
    numB = numB * 2;
```

```
end
```

```
if rem(hicker, 180) == 0
```

```
    population(1, index) = numA;
```

```
    population(2, index) = numB;
```

```
    index = index + 1;
```


```
end
```

← loop - goes to next
available position


```
figure(1);
```

```
plot(1:9, population(1, :));
```

```
hold on;
```



log plot,
not normal
b/c you can't
see it
otherwise.



Exam covers everything up to & not including reduction. Located in Tech Auditorium

Matlab:

Solve $Ax = b$

Method 1: $x = A \setminus b$


Method 2: $\text{inv}(A) * b$ → Gaussian Elimination (Matlab uses this)
 > not equivalent in Matlab's eyes.

$$x - 2y = 0$$

$$x + y = 1$$

$$y = 1 - x$$


$$y = \frac{x}{2}$$




you try to see if they intersect.

but if they were
 then 0, & if they
 were the same line, there
 are ∞ solutions.

3x3 systems intersect. Planes coinciding, being 11, or 3 planes intersect but not all 3 form a pt.



very close together, even though it has a unique solution, but that will appear that it has an ∞ of solns. $\therefore$ b/c we use floating pt. #s.



pretend that's the graph for $x - 2y = 0$
 $x + y = 1$

$$-x + 2y = 0$$

$$x + y = 1$$

$$3y = 1$$

$$y = \frac{1}{3}$$

By adding them,
 we're moving them,
 rotating them without
 leaving the intersection
 point.

You multiply everything
 by a nonzero #.

FBI

$$x + 0y + 0z = x_1$$

$$0x + y + 0z = y_1$$

$$0x + 0y + z = z_1$$

Solving systems of linear equation

$$2x + 3y + z = 10$$

$$-x + z = -1$$

$$3x - y + 2z = -7$$

pivot

$$\begin{bmatrix} 1 & 0 & 0 & 5 & 1 \\ 0 & 1 & 0 & 5 & 2 \\ 0 & 0 & 1 & 5 & 3 \end{bmatrix}$$

Start w/ augmented matrix:

$$\begin{bmatrix} 2 & 3 & 1 & 10 \\ -1 & 0 & 1 & -1 \\ 3 & -1 & 2 & -7 \end{bmatrix} \rightarrow \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \end{bmatrix}$$

leading values
 echelon matrix

Rules

0 If there are any rows of all 0's, they

UW 153.1

Once you have the
echelon form, you can do
a few more things like
multiplying & adding to make
#s 0.

- ① If there are any rows of all 0's, they
will be at the bottom.
- ② Everything in the column below
a leading value (1st # after 0)
is 0
- ③ Every leading # is somewhere
to the right of the leading #
above it. * (to the right of
the last #.

$$\begin{bmatrix} 2 & 3 & 1 & 10 \\ -1 & 0 & 1 & -1 \\ 3 & -1 & 2 & -7 \end{bmatrix} \rightarrow [(row 2)3] + row 3$$

$$\begin{bmatrix} 2 & 3 & 1 & 10 \\ -3 & 0 & 3 & -3 \\ 3 & -1 & 2 & -7 \end{bmatrix} \text{ intermediate step}$$

$$\begin{bmatrix} 2 & 3 & 1 & 10 \\ -1 & 0 & 1 & -1 \\ 0 & -1 & 5 & -10 \end{bmatrix} \text{ and you get this.} \rightarrow [(row 2)2] + row 1$$

$$\begin{bmatrix} 2 & 3 & 1 & 10 \\ -2 & 0 & 2 & -2 \\ 0 & -1 & 5 & -10 \end{bmatrix} \text{ intermediate}$$

$$\begin{bmatrix} 0 & 3 & 3 & 8 \\ -1 & 0 & 1 & -1 \\ 0 & -1 & 5 & -10 \end{bmatrix} \text{ and you get this.} \text{ swap rows 1 + 2}$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & 3 & 3 & 8 \\ 0 & -1 & 5 & -10 \end{bmatrix} \rightarrow [(row 3)3] + row 2$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & 3 & 3 & 8 \\ 0 & -3 & 15 & -30 \end{bmatrix} \text{ int.}$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & 0 & 18 & -22 \\ 0 & -1 & 5 & -10 \end{bmatrix} \rightarrow \text{swap rows 2 + 3}$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & -1 & 5 & -10 \\ 0 & 0 & 18 & -22 \end{bmatrix} \rightarrow \text{divide row 3 by 18}$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & -1 & 5 & -10 \\ 0 & 0 & 1 & -11/9 \end{bmatrix} \rightarrow (row 3) - 5 + row 2$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & -1 & 0 & -10 + \frac{55}{9} \\ 0 & 0 & 1 & -11/9 \end{bmatrix} \rightarrow (row 2) \times (-1)$$

$$\begin{bmatrix} -1 & 0 & 1 & -1 \\ 0 & 1 & 0 & 10 - \frac{55}{9} \\ 0 & 0 & 1 & -11/9 \end{bmatrix} \rightarrow [(row 3) \times (-1)] + row 1$$

$$\begin{bmatrix} 1 & 0 & 0 & 11/9 - 1 \\ 0 & 1 & 0 & 10 - \frac{55}{9} \\ 0 & 0 & 1 & -11/9 \end{bmatrix} \text{ and those are the answers, } x, y, z.$$

$$\begin{bmatrix} 1 \\ 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

0 = 0 is a true statement,
One free variable that can take any value, *.

Multiplying Matrices

Tuesday, October 10, 2006

7:04 AM

$Ax = b$
 $x = A^{-1}b$ Not a good way. Like doing this: $7x = 21$
 $x = 7^{-1} \cdot 21$

instead, do this: $x = 21/7$. $x = b/A = A \backslash b$

3 types of operations you can apply to the rows of an augmented matrix
 - Interchange

$[A \ B]$ $\rightarrow$ $[I \ b^*]$ is the solution

$\begin{bmatrix} * & * & * \\ * & * & * \\ * & * & * \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$

When doing this by hand, you can swap rows. But when writing a program, you can't have it do random ops, it's hope for the best. You

$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$

$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} s_1 \\ s_2 \\ s_3 \end{bmatrix}$ need a sequence

Mult. 1st row by $-a_{21}/a_{11}$
 & add 2nd row. Adds a 0 in position (2, 1)

$a_{11} \left(-\frac{a_{21}}{a_{11}} \right) = -a_{21}$
 $-a_{21} + a_{21} = 0$

$\begin{matrix} a_{11} & a_{12} & a_{13} \\ 0 & (a_{22} + a_{12}(-\frac{a_{21}}{a_{11}})) & (a_{23} + a_{13}(-\frac{a_{21}}{a_{11}})) \\ a_{31} & a_{32} & a_{33} \end{matrix}$ This is a for loop

Mult. 1st row by $-a_{31}/a_{11}$ & add to a_{31}

$\begin{bmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & * \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$

$$\begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & -1 & -4 \\ 0 & 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -5 \\ 0 & 1 & 0 & -4 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -4 & 0 & 8 \\ 2 & -3 & 2 & 1 \\ 5 & -10 & 4 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -4 & 0 & 8 \\ 0 & 5 & 2 & -15 \\ 0 & 10 & 4 & -39 \end{bmatrix} = \begin{bmatrix} 1 & -4 & 0 & 8 \\ 0 & 5 & 2 & -15 \\ 0 & 0 & 0 & -9 \end{bmatrix}$$


this says $0x + 0y + 0z = -9$
this means $\emptyset$.

When you have $0x + 0y + 0z = 0$

then you only have 2 equations & 3 unknowns left, so you choose the variable where fault it is and you write the other variables in terms of the unknown.

This gives you an infinite amount of solutions.

Math Club



As $u \rightarrow \text{Math club}$,
 $\text{fun} \rightarrow \infty$.

$$.00001x + 2y = 2$$

$$x + y = 2$$

$$\begin{bmatrix} .00001 & 2 & 2 \\ 1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} .00001 & 2 & 2 \\ 0 & .99999 & 1.99999 \end{bmatrix} \rightarrow \begin{bmatrix} .00001 & 2 & 2 \\ 0 & 1 & .99999 \end{bmatrix}$$

$$\rightarrow \begin{bmatrix} .00001 & 0 & 0^* \\ 0 & .99999 & .99999 \end{bmatrix} \quad 0^* \approx 0, \text{ approx soln } x=0, y=1. \text{ Incorrect.}$$

Rounding errors cause the wrong answer.

Swap instead

$$\begin{bmatrix} .00001 & 2 & 2 \\ 1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 2 \\ .00001 & 2 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 2 \\ 0 & .99999 & 1.99999 \end{bmatrix}$$

$$\rightarrow \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & .99999 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 1^* \\ 0 & 1 & .99999 \end{bmatrix}$$

$$x^* = 1 \quad x^* = 1$$



Manufacturing
Agriculture
Services

$$C = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$d = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix}$$

$$x = Cx + d$$

1 unit M consumes .1 units M

1 unit A consumes .6 units M

x_2 units of A consumes $.6x_2$ M

x_1 units of M consumes $.1x_1$ M

x_3 units of S consumes $.6x_3$ M

$$x_1 = .1x_1 + .6x_2 + .6x_3 + d_1$$

same thing as

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + d_1 = x_1$$

another way to view the matrix is

$$\begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \end{bmatrix} \overset{A_1}{*} x_1 + \begin{bmatrix} a_{12} \\ a_{22} \\ a_{32} \end{bmatrix} \overset{A_2}{*} x_2 + \begin{bmatrix} a_{13} \\ a_{23} \\ a_{33} \end{bmatrix} \overset{A_3}{*} x_3 + \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

linear combination of A_1, A_2, A_3 (column vectors)

$$c_1 A_1 + c_2 A_2 + c_3 A_3$$



$$x = Cx + d$$

so 3 diff ways to solve.

Linear combination

Matrix

Linear system of equations




every time you multiply a vector by a scalar, you move its end pt. along the original line.

$$-5 \begin{bmatrix} 3 \\ 6 \end{bmatrix} = \begin{bmatrix} -15 \\ -30 \end{bmatrix}$$

$$-1 \begin{bmatrix} 3 \\ 6 \end{bmatrix} = \begin{bmatrix} -3 \\ -6 \end{bmatrix}$$

$$2 \begin{bmatrix} 3 \\ 6 \end{bmatrix} = \begin{bmatrix} 6 \\ 12 \end{bmatrix}$$


$$\begin{bmatrix} 3 \\ 2 \end{bmatrix} + 1 = \begin{bmatrix} 3 \\ 2 \end{bmatrix} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$



$$\begin{bmatrix} 3 \\ 2 \end{bmatrix} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

$$c(v_1 + v_2 + v_3 + v_4) \quad \text{or} \quad av_1 + bv_2 + cv_3 + dv_4$$

$$2v_1 + v_2$$



you have a grid when you add them together

↑ represents linear combination of vectors w/ some coefficients

$$\begin{matrix} v_1 & v_2 & v_3 & \dots & v_p \\ n \times 1 \end{matrix}$$

$$c_1 v_1 + c_2 v_2 + c_3 v_3 + \dots + c_p v_p$$

span subset of $\mathbb{R}^n$

does w belong in this span?

$$\text{span} \{v_1, v_2, \dots, v_p\} = \text{grid above}$$

$$v_1 = \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} \quad v_2 = \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix}$$

$$v_3 = \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix} \quad y = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix}$$

$$x_1 v_1 + x_2 v_2 = w$$

$$x_1 \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} + x_2 \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix}$$

we want to find h so that y is in the span v_1, v_2, v_3

$$\begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} x_1 + \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix} x_2 + \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix} x_3 = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix}$$

$$\begin{bmatrix} 1 & 5 & -3 \\ -1 & -4 & 1 \\ -2 & -7 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix}$$

$$\begin{bmatrix} 1 & 5 & -3 \\ 0 & 1 & -2 \\ -2 & -7 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \\ -8 + h \end{bmatrix}$$

$$\begin{bmatrix} 1 & 5 & -3 \\ 0 & 1 & -2 \\ 0 & 3 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \\ -8 + h \end{bmatrix}$$

↓

$$\begin{bmatrix} 0 & 5 & -6 \\ 1 & 5 & -3 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -1 \\ -5+h \end{bmatrix}$$

$$0 = -5 + h$$

$$5 = h$$

If h is not 5, the system will not have a solution. So y is not in the span. As long as $h=5$, y is in the span.

Let A be an $m \times n$ matrix.

- Columns of " A " span in the augmented matrix
- $Ax=b$ has a solution for every b in the augmented matrix
- Every b in the augmented matrix is a linear combo. of the columns of A
- A has a pivot position in every row

A is the coefficient matrix, not the augmented.

If one of these is true, the others are. If one is not true, the others aren't.

If you want to prove something is not true, then the others will be not true as well.

Could a set of 3 vectors in $\mathbb{R}^4$ span all of $\mathbb{R}^4$?

$$\begin{bmatrix} 1 & 1 & 1 \\ -1 & 1 & 1 \\ -1 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} * \\ * \\ * \\ * \end{bmatrix} \leftarrow b$$

$\uparrow$
 v_3

So no, b/c every ~~part~~ ^{row} does not have a pivot position.

4x5 matrix: is it possible to delete a column & still have the remaining ones span $\mathbb{R}^4$?

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

$\uparrow$
 v_3

While
Loops

$$x_{n+1} = Cx_n + d$$

$$x_1 = Cx_0 + d$$

$$x_2 = Cx_1 + d$$

$$x_3 = Cx_2 + d$$

$$\text{so } x_n = Cx_n + d$$

Sum

Thursday, October 12, 2006
7:01 AM

$$\frac{1}{x^0} + \frac{1}{x^1} + \frac{1}{x^2} + \frac{1}{x^3} + \dots + \frac{1}{x^n}$$

$$\begin{aligned} \text{sum} &= 0 \\ \text{sum} &= \text{sum} + \frac{1}{x^1} \\ \text{sum} &= \text{sum} + \frac{1}{x^2} \\ &\vdots \end{aligned}$$

$$\text{sum} = \text{sum} + 1/x^n \text{ k}$$

$$\text{sum} = 0$$

for is used when we know how many times we want to iterate

for index = 0:12 index = counter

$$\text{sum} = \text{sum} + 1/x^n \text{ index}$$

end

for ii = 1:2:5

A(ii) = x

ii = 1, ii = 3, ii = 5
% the 1st, 3rd, & 5th element will = x. ← of array A

end

while (condition)

==

end

% the moment condition ≠ true, it stops.
Condition needs to change. It needs
to become false or the loop will continue
forever.

$$\text{sum} = 0$$

$$x = 1$$

while x > 0

$$\text{sum} = \text{sum} + \frac{1}{x};$$

end

% This will go on forever

$$\begin{aligned} 0 &\neq \frac{1}{1} = \frac{1}{1} \\ \frac{1}{1} + \frac{1}{2} &= \frac{3}{2} \end{aligned}$$

$$\text{sum} = 0$$

$$x = 1$$

while x > 0

$$\text{sum} = \text{sum} + \frac{1}{x};$$

if x == 1 break;

end

% this would execute only once

$$\text{count} = y$$

$$\text{sum} = 0$$

$$x = 1$$

while x > 1

$$\text{sum} = \text{sum} + \frac{1}{x};$$

```

sum=0
x=1
while x>1
    sum=sum + 1/x;
    if x==1 break;
end
↑
go on = input('Continue? y/n ');

```

```

A = [ ]
[m n] = size(A);
for r = 1:m
    for c = 1:n
        A(r,c) = 0
    end
end

```

```

r=1
    c=1
        A(1,1)=0
    c=2
        A(1,2)=0
    c=3
        ...
    c=n
        A(1,n)=0
r=2
    c=1
        A(2,1)=0
    c=2
        A(2,2)=0
    etc.


```

% This is inefficient because you're accessing the matrix row by row. In Matlab, matrices are stored in column format. You're jumping around in memory.

% You could have eliminated the inner loop by using a colon.
 For r = 1:m
 A(r, :) = 0
 end

A([2 4], :)
 A([2 2], :)
 m x n matrix A(6)

2nd + 4th row completely.
 Access the 2nd row twice
 Counts six its in each column



ones(5)' 5x5 matrix of all 1's

ones(5,4)

ones(5,2) ...

`eye(size(B))` identity matrix of size that's been specified

$$A(1:2:5, :) = B(1:2:5, :)$$

it will copy rows 1, 2, & 5 ^{from A} completely into B.

only if $\text{size}(A) = \text{size}(B)$

$$A(1:2:5, 4:\text{end})$$

4 to the last column, 1st, 3rd, & 5 elements in the row. So (1,4) to end of column.
(3,4) to end of its column.

$$A = \text{zeros}(5)$$

$$A([1\ 3], [2\ 5]) = [0\ 1; 1\ 2]$$

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

0	0	0	0	1
0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	0

$$A = \text{zeros}(4, 3); \quad A(1:2:5, :) = 1:2:5; \quad x = A(4, 3)$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Sizes do not compute.

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

There is no 4th row now.

`break` takes you out of a loop

`continue` skips the iteration but goes back to the top

`return` terminates the program outside of the loop

If you're doing an array operation, you need the dot.

Every time you have `x = %6.2f`, `y = %d`

This prints a # that takes up 6 spaces, 2 of which are after the decimal point, and then `%d`, an integer.

1. Arithmetic
2. Relational
3. Negations

what
Chapman says

1. Negations
2. Arithmetic
3. Relational

The right way

$$\begin{array}{c|c} \sim 1 & 0 \text{ negative (st)} \\ \text{not} & \\ 0 & \sim 1 \end{array}$$

If a plane is spanned by vectors $\vec{v}$ & $\vec{u}$, then every vector $\vec{b}$ located in this plane can be expressed as a linear combination of $\vec{v}$ & $\vec{u}$


$$\vec{b} = x_1 \vec{v} + x_2 \vec{u}$$

$$\vec{b} = x_1 \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} + x_2 \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} v_1 & u_1 \\ v_2 & u_2 \\ v_3 & u_3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

If you want to know whether a certain vector lies in the hyperplane spanned by $\vec{v}$ & $\vec{u}$, find coefficients x_1 & x_2 so the system

$$\begin{bmatrix} v_1 & u_1 \\ v_2 & u_2 \\ v_3 & u_3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$



More practical: Open Leontief model if the econ probos we have

1 Closed Leontief model. Small ^{alien} civilization completely self sufficient. Nothing comes from outside, nothing goes out.

The farming alien consumes 40% of the food, 31% of the clothes, 44% of the energy.

The tailor alien consumes 25% of the food, 3% of the clothes, & 20% of the energy.

The coal mining alien consumes 35% of the food 33% of the clothes, & 36% of the energy

What should be the price of food, clothes, & energy so that the income of each person exactly equals the output? You want everything produced to be consumed.

rfj

$$\begin{aligned} .40f + .31c + .44e &= f \\ .25f + .36c + .20e &= c \\ .35f + .33c + .36e &= e \end{aligned}$$

$$\begin{aligned} -.60f + .31c + .44e &= 0 \\ .25f + -.64c + .20e &= 0 \\ .35f + .33c + -.64e &= 0 \end{aligned}$$

$$\begin{bmatrix} -.60 & .31 & .44 \\ .25 & -.64 & .20 \\ .35 & .33 & -.64 \end{bmatrix} \begin{bmatrix} f \\ c \\ e \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

reduce matrix

$$\begin{bmatrix} \dots \end{bmatrix} \begin{bmatrix} f \\ c \\ e \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$Cx = x \rightarrow \begin{bmatrix} f \\ c \\ e \end{bmatrix}$$


$$Cx - Ix = 0$$

$$(C - I)x = 0$$

Homogeneous systems where right side = 0.

System has a trivial soln, 0. But, if the equation has ~~at~~ at least 1 free variable, we'll have a nontrivial soln.

$$\vec{x} = e \cdot \vec{v} \text{ where } \vec{v} = \begin{bmatrix} 1.1275 \\ 1 \end{bmatrix}$$



$$\vec{x} = \text{span}(\vec{v})$$

$$x = e \cdot \sqrt{\text{represents a line}}$$

$$\vec{x} = c_1 \vec{v} + c_2 \vec{v} \text{ // plane}$$


$$\begin{bmatrix} 1 & 0 & 1.6 \\ 0 & 1 & .4 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -3 \\ 0 \end{bmatrix}$$

$$x_1 + 1.6x_3 = -4$$

$$x_2 + .4x_3 = -3$$

$$x_3 \text{ is free}$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 - 1.6x_3 \\ -3 - .4x_3 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -3 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1.6 \\ -.4 \\ 1 \end{bmatrix}$$





Networks



or



Merchandise goes in, goes



TEN-THIRTEEN

1013



The yin
gray!

$$x_{13}$$

$$x_{12}$$

$$x_{23}$$

$$x_{24}$$

$$x_{25}$$

$$x_{35}$$

$$x_{45}$$

$$x_{46}$$

$$x_{56}$$

$$x_{13} + x_{23} = x_{35}$$

$$x_{24} = x_{45} + x_{46}$$

$$x_{46} + x_{56} = 12$$

$$x_{23} + x_{35} + x_{45} = x_{56} + 3$$

$$x_{12} + 5 = x_{23} + x_{24} + x_{25}$$

$$x_{10} = x_{12} + x_{13}$$



So we build an order. Pick an equation.

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_{13} \\ x_{12} \\ x_{23} \\ x_{24} \\ x_{25} \\ x_{35} \\ x_{45} \\ x_{46} \\ x_{56} \end{bmatrix} = \begin{bmatrix} 3 \end{bmatrix} x_{13} + x_{35} + x_{45} - x_{56} = 3$$

$$\left[\text{Really huge matrix} \right] x = \begin{bmatrix} 0 \\ 0 \\ 12 \\ 3 \\ -5 \\ -10 \end{bmatrix}$$



Spanning Tree

Tuesday, October 17, 2006
7:06 AM





distance
Stand for roads
Which roads should we pave?

This forms a spanning tree - a graph that has no cycles.



Max. capacity at each node,
find max. amount of product
you can generate utilizing
as many of the routes & capacities
as often.



$$\begin{aligned} ① & x_{12} + x_{14} = 10 \\ ② & x_{14} + x_{23} = x_{12} \\ ④ & x_{43} + x_{42} = x_{14} \\ ② & x_{12} + x_{23} = x_{14} \end{aligned}$$

x_{12}	x_{14}	x_{24}	x_{23}	x_{42}	x_{43}	x_{34}
1	1	0	0	0	0	10
0	0	0	1	0	0	0

$$\begin{aligned} \textcircled{3} \quad & x_{23} + x_{43} = 6 \\ \textcircled{5} \quad & x_{14} + x_{34} + x_{24} = 4 \end{aligned}$$

$$\begin{aligned} & 0 \quad 0 \quad - \quad - \quad - \quad - \end{aligned}$$



so you continue this. If you have negative flow, you just subtract that from the appropriate equation.

After you find parametric flows,


$$\begin{aligned} 6 + x_{24} - x_{43} &= x_{12} \\ 9 - x_{24} + x_{43} &= x_{14} \\ 6 - x_{43} &= x_{23} \\ x_{24} &= \text{free} \\ x_{43} &= \text{free} \end{aligned}$$

basic soln

$$\begin{bmatrix} x_{12} \\ x_{14} \\ x_{23} \\ x_{24} \\ x_{43} \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \\ 6 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} x_{24} + \begin{bmatrix} -1 \\ 1 \\ -1 \\ 0 \\ 1 \end{bmatrix} x_{43}$$



So if you have info about the money, you want to take advantage of the cheaper soln.



So now you don't want 6 from 2 to 3 you want



When you add/remove flow it affects other edges that form a cycle.



Now it's not a spanning tree ~~because~~ but it's an improved soln even if it has cycles




4-5-7

examine how the flow
along the cycle will change
if you push more flow along
the chosen edge.

Add ϵ up and see if you
lose or gain money.





The basic soln is a spanning tree
 If you add the middle edge, you get a cycle.

$2(6+t) + 3(6+t) + 5(4-t) = 11$ if you end up w/ less \$, then it's good - you saved \$. If more \$, don't do it. It's a waste of time & money.

If you subtract enough from one branch, you get a new soln w/out that branch included. In the end, you should end up w/ a spanning tree. If you move the columns around, you end up w/ a diff spanning tree.

if _____
 sort()

sort being a function you come up w/.

if _____
 sort()

$A = [\quad]$

If you loaded this up and you previously had an A, it would override the previous A. Same is not true for functions. What happens in a function stays in a function. Like Vegas. Or Risa.

WHAT HAPPENS ON

RISA

STAYS ON RISA

WHAT HAPPENS IN

$f(x)$

name.m
function [outarg1, outarg2 ...] = name [inarg1...]
 ? Name description
 ? Help text
 ? Comments
 Statements

WRITE THE WORD "FUNCTION"

$T(x)$
 stays in $F(x)$  sounds like a good t-shirt,

outarg1, outarg2 are placeholders. They hold the final results in your program.

function [N,M] = size(A)

(rows, columns) = size(A)

>> help size

>> size

>> look for average

help lines with no blank lines in between

No

% help text

It will see this as
 the beginning of
 the function.

% Comments
 Statements

There are a lot of comments that go into this type of work.

temp = (temp - 32) * 5/9

% Convert F to C.

NOT % temp - 32 times 5/9.

[n,m] = size(A);

% verify it's a column vector

if m ~= 1

error('Input must be a column vector')

end

% compute the average

sum = 0;

for ii = 1:n

sum = sum + A(ii)

end

avg = sum/n;

$$\text{avg} = \text{sum}/n;$$

% input arg - column vector
 % beginning | avg - (output arg) avg. of values

parameter passing

$$v = [1; 2; 3; 4]$$

$$x = \text{average}(v)$$

plugs v into A , but doesn't actually change v .

MATLAB sorts

Thursday, October 19, 2006
7:03 AM

$n = \text{size}(A)$
 $[n \ m] = \text{size}(A)$
↑ ↑
rows columns

Sorting means taking a sequence of elmb. & ordering them


- #s from largest to smallest
- names from A-Z
- etc.

Insertion sort is not very good for big arrays, but if the array is small, or if the array is more or less sorted. You go along until you find one that needs to be behind/in front of current ones.

Another method is to split the pile into smaller piles and then even smaller ones. Until it's done.


Bucket sort is when you have lots of piles and you put each one in the right pile as you go.

Selection sort is the worst method. You go through the whole thing & find which one should be 1st. Then you find the one that should be done.



Insertion sort

20 | 78, 45, 5, 20
20 78 | 45, 5, 20
20 45 78 | 5, 20
5, 20, 45, 78 | 20
5, 20, 20, 45, 78 |



CAN WE GET PAST 17?

Start with wall at position 2: # of elmts. in array.

el = element at the wall % to the right

Starting at (wall-1) move left as long as we haven't hit the edge & haven't found a place for el
Shift el to the right
repeat

$A(\text{index}) = \text{saved element}$

function [sorted] = isort(v, direction)

% isort sorts a vector using insertion sort in ascending or descending order.

% 'a' for ascending ...

% More info for user.

- checks # args % (verify correct # of args)
- if not correct

`nargchk` checks # args % (verify correct # of args)
`nargchk(min # args, max # args, nargin)` # of args specified
`message = nargchk(1, 2, nargin)`
`error(message);`
 if there is no problem, error message will be empty string.

% we set the direction to be 'a' by default. Has to be 'a' or 'd' or print error
 % message.

CAN WE GET THROUGH 289?

```

if nargin < 2
    direction = 'a';
end
if (direction ~= 'a' & direction ~= 'd')
    error('Invalid direction. Legal values: ...')
end
  
```

```

[rows cols] = size(v);
if ~(rows == 1 || cols == 1)
    % if rows == 1 & cols == 1
    % if (rows == 1 | cols == 1)
    error('Input array should be a vector. You suck!')
end
  
```

1 or 2 input args
 arg1 : row/col vector
 arg2 : (opt.) 'a' 'd'
 1 output : sorted vector
 Algorithm = insertion sort

function [sorted] = isort(v, 5)

↙
 this is different than
 this ↘

$v = [5 \ 2 \ 1 \ 6 \ 3]$
 $\text{sorted } v = \text{isort}(v)$
 $\text{sorted } v2 = \text{isort}(v, 'd')$

msg = narginchk(1, 2, nargin)

↑
 1 to 2 inputs. If you wanted 1-4 inputs, range
 would be 1-4.

If # of args are valid, it will contain a blank line.
 if # of args is wrong, it will print out an error message

if nargin == 1,
 direction = 'a'
end
 if the user doesn't specify, then the
 argument is automatically a.

if direction ~= 'a' & direction ~= 'd'
 error('Legal values are...')
end

[rows cols] = size(v)

if ~[rows == 1 | cols == 1] % if rows ~= 1 or cols ~= 1
 error('Input should be row or column vector')
end

Repeating the operation - once you go through and insert elements, you can go through again. for loop.

for wall = 2 : length(v)

element = v(wall);

walker = wall - 1

while (element < v(walker)
& & walker > 0)


v(walker + 1) = v(walker)

walker = walker - 1

end

v(walker + 1) = element %b/c that's the last empty slot

end



% saving elmt by wall

% to check each value, we need a while loop
b/c we don't know how far it will have

to go,

% elmt is less than item at the walker &
walker hasn't hit the edge



Sorted = v

% assigns v to the output argument & that will
become the value of the function.

% but we're only sorting ascending

% & allows each thing
to be evaluated separately.
if one is false, so is
the other,

% & only tests once at
a time w/ no regard
for the other.

System Consistency

Tuesday, October 24, 2006
7:09 AM

linear search

Start at beginning
keep checking
if you find it
return success
all the way to the end
return failure

function [found] = linsearch(A, target)

% LINSEARCH searches for target in matrix A

% help text

blank line

% comments

message = msgchk(2, 2, nargin)

error(message);

found = 0 % initially you haven't found it

[m n] = size(A)

numelnts = m * n

for ii = 1: numelnts;

if target == A(ii)

found = 1

break

else
continue

end

% useless in this case.


end

% you can have found = 0 at the bottom if we change to break to return. Return stops execution of the function.


for loops nested is another option.

one for loop to go through all cols, one to go through all rows

A(ii) tells you what element you're on.



A single vector spans a line of the same slope. Multiply the vector by a scalar to get a new point



$$\vec{u} + \vec{v} = \vec{w}$$

$$2\vec{u} + \vec{v} = \vec{w}_1$$

Linear dependence of vectors

$$x_3 \vec{w} = x_1 \vec{u} + x_2 \vec{v} \rightarrow x_1 \vec{u} + x_2 \vec{v} - x_3 \vec{w} = 0$$

$\vec{w}$ is a linear combination of $\vec{u}$ & $\vec{v}$.

If at least one is a linear combination of another, you say the whole set of vectors is linearly dependent.

$$c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_p \vec{v}_p = 0$$

where one set of weights $\{c_1, c_2, \dots, c_p\}$ where weights, at least

$$v_i = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$v_1 = \begin{bmatrix} 1 \\ -\frac{2}{3} \end{bmatrix}$$

$$v_2 = \begin{bmatrix} 4 \\ \frac{5}{6} \end{bmatrix}$$

$$v_3 = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

Are they linearly dependent?

See if the system is consistent

$$c_1 \begin{bmatrix} 1 \\ -\frac{2}{3} \end{bmatrix} + c_2 \begin{bmatrix} 4 \\ \frac{5}{6} \end{bmatrix} + c_3 \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{pmatrix} c_1 & c_2 & c_3 \end{pmatrix} \begin{bmatrix} 1 & 4 & 2 \\ -\frac{2}{3} & \frac{5}{6} & 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$0,0,0$ is a trivial soln. we're looking for a non-trivial soln.

we're hoping for a free variable

$$\begin{bmatrix} 1 & 4 & 2 & 0 \\ 0 & -\frac{2}{3} & -\frac{5}{6} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} c_1 + 2c_3 &= 0 \\ c_2 + c_3 &= 0 \\ 0c_3 &= 0 \end{aligned}$$

$$x_3 \vec{w} = x_1 \vec{v} + x_2 \vec{v} \rightarrow x_1 \vec{v} + x_2 \vec{v} - x_3 \vec{w} = \vec{0}$$

For linear independency, you should expect a pivot in every column. You want exactly one soln.

Independence means $0,0,0$ is the only soln.

If you can find out of a collection of vectors, one of them is a linear combination of the others, then it is dependent.

$$\begin{bmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 0 & -1 & -1 \end{bmatrix}$$

$$\vec{v} \quad \vec{v} \quad \vec{w}$$

$$\vec{w} = \vec{v} + \vec{v}$$

$$1\vec{w} - 1\vec{v} - 1\vec{v} = \vec{0}$$

To prove they're independent, none of them are a linear combo of others.

$$\begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} ? \\ ? \end{bmatrix}$$

yes, linear combo

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 8 \\ 3 \end{bmatrix}$$

You can take anything times 0 vector.

$$5\vec{v} + 0\vec{v} + 0\vec{v} = \vec{0}$$

$$\{c_1, c_2, c_3\} \neq 0$$

A $n \times n$ for every $b \in \mathbb{R}^n$ $Ax=b$ has at most 1 soln

why must the cols of A be linearly independent?


$$Ax=b \text{ even if } b=0$$

$Ax=0$, system has at most one soln. This has, at very least a trivial soln (all 0's). Therefore that soln has to be that

be that.
If coefficients are all 0, cols of A are linearly independent
by definition.


$$Ax=b$$

we have a vector x , we apply a transformation A , & we view vector b .




$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -x_2 \\ -x_1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$




$$\begin{matrix} 3+4=7 \\ 0+2=2 \end{matrix}$$

$$\begin{bmatrix} 7 \\ 2 \end{bmatrix}$$



You pulled $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$ into $\begin{bmatrix} 7 \\ 2 \end{bmatrix}$

SE TAG CHOICES, INC.



all x 's go over...
infinite # of resulting transformations.
If no x 's go over, no soln to set.

Because of the 0, the 2 stays the same.

Height doesn't change
called a shear transformation

$$\begin{bmatrix} 0 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \rightarrow \begin{bmatrix} x_1 \\ x_2 \\ 0 \end{bmatrix}$$

$\mathbb{R}^3$ $\mathbb{R}^3$

The vector was projected onto 2 axes only, so you get 2 things filled in, not 3.

$$\underbrace{T(x)}_{Ax} = b$$

Same thing, diff way to write.

$$T(x) = Ax \quad \text{any vector } x \text{ in } \mathbb{R}^n \text{ Domain} \quad b = \text{any vector in } \mathbb{R}^m, \text{ codomain}$$

$T(x)$ = "image of x " All images together = range. Not necessarily = to codomain, but that is a possibility

$$T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v}) \text{ for all vectors } \vec{u} + \vec{v} \text{ in domain of } T$$

$$T(k\vec{u}) = k T(\vec{u}), \quad k = \text{scalar}$$

$$\begin{aligned} T(x) &= Ax + b \quad b \neq 0, \\ T(\vec{u} + \vec{v}) &= A(\vec{u} + \vec{v}) + b \\ &= \underbrace{A\vec{u}}_{\text{not } T(\vec{u})} + \underbrace{A\vec{v}}_{T(\vec{v})} + b \end{aligned}$$

$$T(\vec{u}) + T(\vec{v}) = A\vec{u} + b + A\vec{v} + b \quad \text{not the same as}$$

$\therefore$ This particular transformation is not linear.

JUNIPER
SACRIFICE
by
CONNOR
O'MALLEY



$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right)$$

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} x_1 + \begin{bmatrix} 0 \\ 1 \end{bmatrix} x_2\right)$$

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} x_1\right) + T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix} x_2\right)$$

$$x_1 T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + x_2 T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$$

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{pmatrix} 5 \\ -7 \end{pmatrix}$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \cdot \underbrace{\begin{bmatrix} 5 & 3 \\ -7 & 2 \end{bmatrix}}_A$$



how does the transformation affect x_1, x_2 ?
what is the standard transformation matrix A ?

Standard matrix for the transformation that will rotate a vector $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ by θ or ψ (pos)

counter-clockwise



counter-clockwise



$$\Delta 1 \quad \sin \psi = \frac{y}{1} \quad \sin \psi = y$$

$$\cos \psi = \frac{x}{1} \quad \cos \psi = x$$

$$\Delta 2 \quad \sin \psi = \frac{-x}{1} \quad \sin \psi = -x$$

$$\cos \psi = \frac{y}{1} \quad \cos \psi = y$$


$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} \cos \psi \\ \sin \psi \end{bmatrix}$$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} -\sin \psi \\ \cos \psi \end{bmatrix}$$

$$A = \begin{bmatrix} \cos \psi & -\sin \psi \\ \sin \psi & \cos \psi \end{bmatrix}$$

Horizontal shear from $\mathbb{R}^3$ to $\mathbb{R}^2$

$$\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 + kx_2 \\ x_2 \end{bmatrix}$$



$$Ax = b$$

$$Ay = b$$

You can have 2 vectors that will both give you b .

"Onto" Mapping means every vector in b is the image of at least 1 x . We cover all of $\mathbb{R}^m$, in other words, we span all of $\mathbb{R}^m$.

Columns of A span $\mathbb{R}^m$. $\mathbb{R}^n \rightarrow \mathbb{R}^m$

which means $Ax = b$ always has a soln.

Each vector b in $\mathbb{R}^m$ is the image of at least one vector x in $\mathbb{R}^n$. Must be a pivot position in each row of A .

If one of Theorem 4 conditions are satisfied, then you can

Say 'yes, it's $\mathbb{R}^2$ or whatever it is.

If it's true for every b , you can pick 0.

$Ax=0$ For every b , there is at most one x that matches it. They already have at least one soln if $A \neq 0$, and so this system has the trivial soln.

This definition tells us there is exactly one soln, & in this case it's trivial.

~~Every~~ Every column has a pivot, so columns are linearly independent.

$$T(x_1, x_2, x_3) = (x_1 - 5x_2 + x_3, x_2 - 6x_3)$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 - 5x_2 + x_3 \\ x_2 - 6x_3 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & -5 & 1 \\ 0 & 1 & -6 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

one-to-one or onto?

Onto

$\mathbb{R}^3$ onto $\mathbb{R}^2$

For every b there is an x that gives us the b .
So the system has a soln.



Every row has a pivot so soln is onto.

Not one-to-one

Transformations

Friday, October 27, 2006
7:28 AM

$$Ax = b$$



$$T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$$

$$T(c\vec{u}) = c T(\vec{u})$$

$$T(\vec{u}) + T(\vec{v}) = 3\vec{u} + 3\vec{v}$$

$$T(c\vec{u}) = 3(c\vec{u}) = 3(3\vec{u}) = 9\vec{u}$$

$$c T(\vec{u}) = 3 T(\vec{u})$$

X-FILES

FOX
MULDER

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} 4x_1 - 2x_2 \\ 3|x_2| \end{bmatrix}$$

$$x_2 < 0 \rightarrow -x_2$$

$$x_2 > 0 \rightarrow x_2$$

$$c \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} cx_1 \\ cx_2 \end{bmatrix}$$


Has to be true for every scalar. So
come up with one that disproves it.

$$c = -1$$

$$T\left(\begin{bmatrix} -x_1 \\ -x_2 \end{bmatrix}\right) = \begin{bmatrix} 4(-x_1) - 2(-x_2) \\ 3|-x_2| \end{bmatrix} = \begin{bmatrix} -4x_1 + 2x_2 \\ 3x_2 \end{bmatrix}$$

$$c T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = - \begin{bmatrix} 4x_1 - 2x_2 \\ 3|x_2| \end{bmatrix} = \begin{bmatrix} -4x_1 + 2x_2 \\ -3|x_2| \end{bmatrix}$$

So it doesn't
work. Transformation
isn't linear.



unit vectors

$$T(x)$$

$$Ax = b$$

$$A [T(e_1) \dots T(e_n)]$$

columns of the
identity matrix

$$x = Ix = [e_1 e_2 \dots e_n] \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

$$Tx = T(Ix) = T([e_1 e_2 \dots e_n])$$

$$= T([e_1 x_1 + e_2 x_2 + \dots e_n x_n])$$

$$= T(e_1 x_1) + T(e_2 x_2) + \dots + T(e_n x_n)$$

$$= T(e_1) x_1 + T(e_2) x_2 + \dots + T(e_n) x_n$$

X-FILES

DANA
SCULLY


FOX
MULDER

$$= T(e_1)x_1 + T(e_2)x_2 + \dots + T(e_n)x_n \\ = [T(e_1) \ T(e_2) \ \dots \ T(e_n)] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Standard matrix of the transformation
A

p. 90

8. $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
through horizontal x_1
through line $x_1 = x_2$



Reflections go like
this, final position
is $(-1,0)$

Shear $e_2 \rightarrow e_2 + 3e_1$

$$T(e_1) = e_1$$

$$T(e_2) = e_2 + 3e_1$$

$$\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

general form is $\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$ for vertical

Standard of Transformation

Tuesday, October 31, 2006
8:02 AM

```

for ii = 1: size(A, 1)
    if ii == jj
        A(ii, col)
    end
end
for li = 1: size(B, 1)
    A(col, li)
end

```

if $ii = 1$, $ii = jj$
 jj is row
 col is column

for each column
 find potential pivot
 for row where pivot is
 check whether it's preceded by all 0's $0:col-1 = 0$
 if yes,
 pivot

$$A = \begin{bmatrix} 1 & -3 & 2 & 0 & -1 & 4 \\ 3 & -5 & -9 \end{bmatrix}$$

Find image of $[1 \ 1 \ 1]'$

← standard matrix of transformation

$$Ax = b$$

← domain

$$T: \mathbb{R}^n \rightarrow \mathbb{R}^m \leftarrow \text{codomain}$$

$$x \in \mathbb{R}^n$$

$$T(x) = b \in \mathbb{R}^m$$

linear transformation




image is individual b

whole image collection = range

If you're given A & x (as in above), multiply A by x to get b .

$$\mathbb{R}^3 \rightarrow \mathbb{R}^2$$

$$\begin{bmatrix} 1 & -5 & 7 \\ -3 & 7 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -5 & -7 \\ 0 & -8 & -16 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ -8 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -5 & -7 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ -8 \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 0 & 1 & 2 & 2 \end{array} \right] \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 - 3x_3 \\ 2 - 2x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} + \begin{bmatrix} -3 \\ -2 \end{bmatrix} x_3$$

Does such an x exist?

Is this x unique?

Is every b the image of at most 1 x ? In other words, are columns of A linearly independent? This would mean one-to-one.

If every b is the image of at least 1 x , it's onto.

If every b is the image of at most 1 x , it's one-to-one.

It's possible for a transformation to be both, if the transformation is the image of exactly one x , then it is onto & one-to-one.

If you can prove for $Ax=0$ that you have a nontrivial solution, you've found one b that breaks property. Transformation is not one-to-one.

$$T: \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$A = m \times n$$

$$\begin{bmatrix} x_1 & \dots & x_n \\ \vdots & & \vdots \\ x_m \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \rightarrow \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$$


$$m \times n \quad n \times 1 \rightarrow m \times 1$$

For every b , we have a solution in our system. That means m should equal the # of pivots

If we know something is one-to-one, we know that if there is a solution, it will be unique. n pivots

It's possible for m or n to be the same. If you know the transformation is onto (or one-to-one), do you know if the

transformation has to be $m \times n$?



1. Reflect one way
2. Reflect other way

$$T_1(x) = b$$

$$A_1 x = b$$

$$T_2(b) = c$$


$$A_2 b = c$$

$$A_2(A_1 x) = c$$

$$(A_2 \cdot A_1) x = c$$

$$\text{so } A = A_2 \cdot A_1$$

$$A = (A_k \cdot A_{k-1} \cdot \dots \cdot A_1)$$



CAPTAIN
JONATHAN
ARCHER



Associative

Distributive

Not commutative

$AB = AC$ does not necessarily
mean $B = C$

$$AB \neq BA$$

$AB = 0$ does not necessarily mean $A = 0$ or $B = 0$

$$(A^T)^T = A$$

$$(A + B)^T = A^T + B^T$$

$$(cA)^T = cA^T$$

$$(AB)^T = B^T A^T$$

Not every matrix can be inverted.

For a matrix X to be inverted, $(A^{-1})A$
 $= I_n$

$$(A^{-1})^{-1} = A$$


$$(AB)^{-1} = B^{-1}A^{-1}$$

$$(A^{-1})^T = (A^T)^{-1}$$

$$(A^T)(A^T)^{-1} = I$$

$$(A^T)(A^{-1})^T = (A^{-1} \cdot A)^T = I^T = I$$

onto $n \geq m$
 1-1 $m \geq n$
 onto & 1-1 $m = n$



$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ this gives you } 1-1 \text{ \& onto}$$

$$T(x) = Bx = [b_1 \ b_2 \ \dots \ b_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = b_1x_1 + b_2x_2 + \dots + b_nx_n$$

$$T(T(x)) = C(Bx) = Cb_1x_1 + Cb_2x_2 + \dots + Cb_nx_n$$

$\uparrow$ $m \times p$ matrix $\uparrow$ $p \times n$ matrix $\times$ $\uparrow$ $n \times 1$ vector $\rightarrow$ $m \times 1$ vector

$$= C[b_1 \ b_2 \ \dots \ b_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = (CB)x$$

$$xy = xz$$

$$\frac{1}{x}xy = \frac{1}{x}xz$$

$$y = z$$

$xy = xz$ what if x is 0?
 $\frac{1}{x}$ doesn't exist. This breaks the rule

$$A^{-1}A = I$$

$$(A^{-1})AB = AC(A^{-1})$$

$$IB = IC$$

$$B = C$$

Not all matrices have an inverse.
 If the inverse exists, you can divide
 & get $B = C$. If not, you're SOL.

X-FILES

MULDER
 SCULLY

SKINNER
 DOGGETT

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$ad-bc \neq 0$$

$$A^{-1} = B$$

A is invertible if there exists B st. $AB = BA = I_n$

$$Ax = b$$

$$x = A^{-1}b$$

$$\text{inv}(A) \times b = x$$

A singular matrix near no inverse.
 Non-singular means it has an inverse.

$$(A^{-1})^{-1} = A$$

$$(AB)^{-1} = B^{-1}A^{-1}$$

$$(A^{-1})^T = (A^T)^{-1}$$

$$E \cdot A = (A^{-1})A = I$$

so A is the inverse of E

Doing ref is like transformations. In the end, you end up with I .



If we can find the individual transformations, we can find the original matrix.

$$\begin{bmatrix} 0 & 0 & 1 \\ 3 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} 7 & 8 & 9 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

only works if you multiply on the left



An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n .

1st prove that if A is invertible, it is ^{row} equivalent to I_n .

$\text{ref}(A)$ has a pivot in every row so $= I_n$

Then prove that if A is row equivalent to I_n , it can be inverted.

If A is row equivalent to I_n , there exist elementary matrices $E_1, E_2, \dots, E_p$ such that $E_p \dots E_2 E_1 A = I_n$ (those intermediate matrices you get after transformations).

Elementary matrices are invertible, so $(E_p \dots E_2 E_1)^{-1}$ exists

Left-multiply both sides of equation by it:

$$(E_p \dots E_2 E_1)^{-1} (E_p \dots E_2 E_1) A = (E_p \dots E_2 E_1)^{-1} I_n$$

$$I_n A = (E_p \dots E_2 E_1)^{-1} I_n$$

$$A = (E_p \dots E_2 E_1)^{-1}$$

$[A \ I]$ perform row reduction

If A is row equivalent to I ,

$[A \ I]$ is row equivalent to $[I \ A]$

Otherwise, A does not have an inverse.

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array} \right]$$



Determinants

Thursday, November 02, 2006
8:03 AM

Determinants:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = a_{11}[a_{22}a_{33} - a_{23}a_{32}] - a_{12}[a_{21}a_{33} - a_{23}a_{31}] + a_{13}[a_{21}a_{32} - a_{22}a_{31}]$$

If $Ax = b$ Goes from $\mathbb{R}^n \rightarrow \mathbb{R}^m$
 $A^{-1}b = x$ Goes from $\mathbb{R}^m \rightarrow \mathbb{R}^n$

If $Ax_1 = b$
 $Ax_2 = b$ $x_1 \neq x_2$
 A is not 1-1.


$T^{-1}(b) = \begin{cases} x_1 \\ x_2 \end{cases}$ So getting 2 answers for this one thing doesn't make sense. b cannot be mapped to 2 things. Transformations cannot look like this.

MULDER • 97 DIUM

If a transformation is not onto, it doesn't work. Transformations only work for 1-1 or onto or both.

$$1 \Rightarrow 2 \Rightarrow 5 \Rightarrow 4 \Rightarrow 3 \Rightarrow 1$$

If there were 5 statements, and you can prove if 1 is true, 2 exists, etc. If you prove all of this, you don't have to do anything else. You know $5 \Rightarrow 2$, or $4 \Rightarrow 5$, or $1 \Rightarrow 5$ etc. Come up w/ a proof to put them in a $\odot$.



Condition # is the stability of a matrix. Depending on round-off error, we can't depend on our solution. It could be wrong. Ill conditioned is a matrix w/ large round-off error, and a large condition #, and therefore it is close to being singular.

$$x = (x, d)$$

$$\begin{bmatrix} d_1 \\ d_2 \end{bmatrix}$$

We're assuming for one sector is d_i is

$$\begin{bmatrix} d_1 \\ \vdots \\ d_n \end{bmatrix} \text{ instead } d_i + \theta.$$

$$X = (I - C) \backslash d$$

$$X = \begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix} \begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_n \end{bmatrix}$$

$$X^{-1} = \begin{bmatrix} s_{11} & \dots & s_{1n} \\ \vdots & & \vdots \\ s_{n1} & \dots & s_{nn} \end{bmatrix} \begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_i + \theta \end{bmatrix}$$

$$= \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \end{bmatrix} \left(\begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_i \\ d_{i+1} \\ \vdots \\ d_n \end{bmatrix} + \begin{bmatrix} \theta \\ 0 \\ \vdots \\ 0 \end{bmatrix} \right)$$

$$= \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_i \\ d_{i+1} \\ \vdots \\ d_n \end{bmatrix} + \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$


$$= \text{II} + \begin{bmatrix} 0 & 0 & s_{i1}\theta & 0 & 0 \\ 0 & 0 & s_{i2}\theta & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & s_{in}\theta & 0 & 0 \end{bmatrix}$$

$$= \text{II} + \begin{bmatrix} s_{i1} \\ s_{i2} \\ \vdots \\ s_{in} \end{bmatrix} \theta$$

$$= (I - C)^{-1} d + \theta * [\text{i+th col of } (I - C)^{-1}]$$

If θ was positive, we produce a little more (external demand slightly higher) & overall production is higher for all sectors.

If θ was neg, overall production less b/c demand $\downarrow$.



$$MP = C$$

$$C \rightarrow P?$$

$$(MP)M^{-1} = CM^{-1}$$

$$P = CM^{-1}$$

So to do this we need a matrix that's invertible.

Use ASCII code to convert message to a sequence of #'s.
Have to represent whitespace as well

Have to represent whitespace as well

Arrange #s into a matrix N .

Use linear transformation to transform N into another matrix.
 N 's inverse must exist or have integer entries

$s = \text{'This is a test!'}$

$\text{double}(s)$ converts to ASCII
 $\text{char}(n)$ converts to characters.

M is the key

$n \times n$ $n \times ?$

[]

If $n \times n = 5 \times 5$, & we have 13 char, 5×3 means we have some extra entries.
add a few spaces

$n\text{numb} = \text{reshape}(\text{double}(s), r, c);$

$r \times c$ matrix
 $\Downarrow$
3 5

$n\text{numb} = \begin{bmatrix} 84 & 115 & 115 & 32 & 115 \\ 104 & 32 & 32 & 116 & 116 \\ 105 & 105 & - & - & - \end{bmatrix}$

diff matrix,
this one's 3×5 .
If you had a 3×3
original, it would
work.

$n\text{numb} - 32 =$
 $\Downarrow$
 $= \text{spaces.}$

So then you get 0-94, w/
95 possible codes.

Come up w/ an M .

$M * (n\text{numb} - 32) = \begin{bmatrix} \text{bunch of} \\ \text{big \#s} \end{bmatrix}$ So you have to
bring them down to
the original range 0-94.

So scale everything based on multiples of 95.

$n\text{code} = \text{mod}(M * (n\text{numb} - 32), 95)$ So then add 32 to get
back to ASCII range.

$n\text{code} = \text{mod}(M * (n\text{numb} - 32), 95) + 32$

3×3 $M \cdot S \xrightarrow{\text{encryp fion key}} \text{cipher text } (c)$

$$M^{-1}c \Rightarrow S$$

(3×3) 1×13
 can't multiply, so must
 reshape $\quad$.

We have 13 elems &
 1 has to become a
 3, so it has to be
 3×5 to contain
 all integers.
 $3 \times \text{cols} \geq 13$

1. Find an M M must be invertible. Inverse only contains integers,
 M only contains integers.

2. Figure out how to reshape $\text{numb} = \text{reshape}(\text{double}(s), 3, 5)$

3. Pad the text w/ extra spaces $\text{ncode} = \text{mod}(m \times (\text{numb} - 32), 95) + 32$
 $\text{scode} = \text{reshape}(\text{char}(\text{ncode}), 1, 15)$

4. Multiply M (encryption key) & scale accordingly $\rightarrow$

To get the value back, we do the same thing

$\text{scode} = 1 \text{ blah}$
 $\text{ncode} = \text{reshape}(\text{double}(s), 3, 5)$
 $\text{numb} = \text{mod}(\text{inv}(m) \times (\text{ncode} - 32), 95) + 32$
 $\text{orig} = \text{reshape}(\text{char}(\text{numb}), 1, 15)$

$$\begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad 1(1-0) - 0(x-0) + 0(0-x)$$

This matrix can be inverted.

$$\begin{bmatrix} 1 & x & x \\ 0 & 1 & x \\ 0 & 0 & 1 \end{bmatrix} \quad 1(1-0) - x(0-0) + x(0-0)$$

so this can be inverted too.

$$M = (M_1 M_2)$$

$$M^{-1} = M_2^{-1} M_1^{-1}$$

$$(M_2 M_1)^{-1}$$

You come up w/ 2 big prime #s, p & q , of which M
 is the product.

Extra spaces must be manufactured with 15.

$$e \cdot d = p \cdot q$$

public

using m & d , the a/hess create a message.
 Mulder has e , & e is a secret. d is a
 well known alien text key, & now only
 Mulder has $e \cdot d$. and he can have

out that they're wrong.

to create spaces: educational doodling

[plaintext spaces] = plaintext You can regard spaces as integers of ASCII value 32

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad ad - bc \neq 0$$

$$\left[\begin{array}{cc|cc} a & b & 1 & 0 \\ c & d & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{cc|cc} a & b & 1 & 0 \\ 0 & d - \frac{bc}{a} & -\frac{c}{a} & 1 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{cc|cc} a & b & 1 & 0 \\ 0 & ad-bc & -c & a \end{array} \right]$$

row of all 0's,
and so you can
stop there. no
inverse.

$A^T \text{ inv} = A \text{ inv}$ We want to prove that A is invertible.

if this
is invertible,

$$A^T B = B A^T = I \quad B \text{ is the inverse, } A^{-1}$$

$$(A^T B)^T = (B A^T)^T = I^T \quad \text{we just took } T(A^T B)$$

$$A \cdot B^T = B^T \cdot A = I$$

B^T is the matrix C for which $CA = I$, and
therefore A^T is invertible.

$$DA = I$$

↓

$$Ax = 0 \rightarrow x = 0$$

$$D(Ax) = (0)D$$

$$(DA)x = 0$$

$$Ix = 0$$

$$x = 0 \quad \checkmark$$

Vector Space, null Space

Tuesday, November 07, 2006
8:02 AM

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} x$$

$$Ax = b$$

$$c_1 x_1 + c_2 x_2 + \dots + c_n x_n = b$$

A vector space is a collection of objects that can be added or scaled. (Added or multiplied by a vector). Not really a vector in the traditional sense.

A nonempty set V of objects called vectors, on which are defined two operations.

addition $\vec{u} + \vec{v} \in V$ for all $\vec{u}, \vec{v} \in V$

scalar multiplication $c\vec{v} \in V$ for all $\vec{v} \in V$ & $c \in F$ (F is a field such as $\mathbb{R}$.)

Addition is commutative

Addition is associative

Addition has an identity elmt. zero vector $\vec{0} \in V$ s.t. $\vec{u} + \vec{0} = \vec{u}$

Addition has an inverse elmt. for each $\vec{u} \in V$, vector $\vec{v} \in V$ s.t. $\vec{u} + (-\vec{u}) = \vec{0}$

Scalar multiplication is distributive over vector addition.

$$c(\vec{u} + \vec{v}) = c\vec{u} + c\vec{v}$$

Scalar multiplication is distributive over field addition.

$$(c + d)\vec{u} = c\vec{u} + d\vec{u}$$

Scalar multiplication is associative.

Scalar multiplication has an identity elmt.

$V = \{(x, y) : x > 0, y > 0\}$ w/ standard addition & scalar mult. in $\mathbb{R}^2$.

(claimed that this is not a vector space.) $\vec{x}$ & $\vec{y}$ have to be positive, so we can't have an inverse elmt. for addition (no vector would add to get $\vec{0}$)

Also if you mult. by a neg. scalar you get something not in the set.

A set of polynomials w/ exactly degree 'n' & addition/mult.

You end up w/ something not in the set.

Convert signal



Addition makes 2 signals.

Multiplication strengthens signal.



Infinite vector b/c it can expand forever.

Subspace

A subset of vector space that is itself a vector space.

A subset H of a vector space V w/ properties:

- Zero vector of V also in H
- H is closed under vector addition $\vec{u} + \vec{v} \in H$ for every $\vec{u}, \vec{v} \in H$
- H is closed under scalar multiplication $c\vec{u} \in H$ for every $\vec{u} \in H$ & scalar c .

A set is closed if 2 elmts of the set, $\neq$ perform operation on them, is also an element in H .

Vector space $\mathbb{R}^2$ is not a subspace of $\mathbb{R}^3$

$$\mathbb{R}^3 \rightarrow (x, y, z)$$

$$\mathbb{R}^2 \rightarrow (x, y)$$

$\mathbb{R}^2$ is not a subset of $\mathbb{R}^3$ when seen like this

X-FILES

$$V = \{(x, y, z)\}$$

$$H = \{(x, y, 0)\}$$

H is a subset of V now.

$(0, 0, 0)$ belongs in H .

if you add any 2 elmts

$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ 0 \end{pmatrix}$$

So closed under addition.

$$c \begin{pmatrix} x \\ y \\ 0 \end{pmatrix} = \begin{pmatrix} cx \\ cy \\ 0 \end{pmatrix}$$


And so this is still legit

Therefore H is a subspace of V .




A line that does not pass through the origin is NOT a subspace. It must contain $(0, 0, 0)$

Every vector space is a subspace of itself.



Every vector space is a subspace of itself.



All polynomials of the form $p(t) = a \cdot t^2$, $a \in \mathbb{R}$
Is this a subspace of P_n ?

$$V = \{p(t) = at^2, a \in \mathbb{R}\}$$

$$p(t_1) + p(t_2)$$

$$\begin{array}{r} x^3 + x^2 + 1 \\ 2x^3 - 2x^2 - 1 \\ \hline 3x^3 - x^2 \end{array}$$

Scalar mult: $2(x^3 + x^2 + 1) = 2x^3 + 2x^2 + 2$

$$(\vec{p} + \vec{q})(t) = \vec{p}(t) + \vec{q}(t) \quad V = \{ \text{All polynomials of degree at most } n \}$$

$$(c\vec{p})(t) = c\vec{p}(t)$$

$$H = \{p(t) = at^2, a \in \mathbb{R}\}$$

Is H a subspace of V ?

Closed under addition - 2 elmts in H added = elmt in H ?

$$at^2 + bt^2 = (a+b)t^2$$

$$b = -a \quad = 0t^2 \quad 0 \text{ belongs}$$

(closed under multiplication = $c at^2$

$$= (ca)t^2 \quad \text{so this works.}$$

$$H = \{ \text{set of all matrices of the form } \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} \}$$

Is this a subset of $M_{2 \times 2}$?

It contains the 0 elmt for a, b, d

Add 2 matrices of this form will give you somekind of result.

$$\begin{bmatrix} a & b \\ 0 & d \end{bmatrix} + \begin{bmatrix} c & e \\ 0 & f \end{bmatrix} = \begin{bmatrix} a+c & b+e \\ 0 & d+f \end{bmatrix}$$

set of elmts $\{v_1, v_2, \dots, v_k\} \in \mathbb{R}^n$ the set of all linear combinations of v_1, v_2, v_k , is a subspace of $\mathbb{R}^n$.

Furthermore: If $v_1, v_2, \dots, v_k$ are in a vector space V then $\text{Span}(v_1, v_2, v_3, \dots, v_k)$ is a subspace of V .

Set of vectors written in matrix form $\begin{matrix} n \times n \\ V \end{matrix}$ we can determine:

Null space

Column space

Row space

Nul A (set of all solns of equation $Ax=0$)

$$\begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & 1 & 4 & -2 \end{bmatrix} \text{ Parametric vector form.}$$

$$\begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & 1 & 4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \begin{bmatrix} x_1 - 7x_3 \\ 2x_4 - 4x_3 \\ x_3 \\ x_4 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -7 & 0 \\ 0 & 1 & 4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The Null space of an $m \times n$ matrix is a subspace of $\mathbb{R}^n$

- zero vector of $\mathbb{R}^n$ is also in Nul A $Ax_1=0, Ax_2=0$
- Nul A is closed under vector addition $A(x_1+x_2)=A x_1 + A x_2 = 0$
- Nul A is closed under scalar mult.

The Column Space of a matrix A (Col A) is the set of all linear combinations of the columns of A.

Column space of an $m \times n$ matrix is a subspace of $\mathbb{R}^m$

- zero vector of $\mathbb{R}^m$ in Col A
- Col A closed under vector addition
- Col A closed under scalar mult.

$$\text{Col A} = \text{Span} \{a_1, a_2, \dots, a_n\}$$

- range of linear transformation $x \mapsto Ax$ (set of all images)
- set of solns of the system.

Row space of a matrix A, Row A, is set of all linear combos of rows of A

Row space of an $m \times n$ matrix is a subspace of $\mathbb{R}^n$

- same reasons.

$$V = \mathbb{R}^3$$

$$H \subseteq V$$

$$0 \in H$$

H is closed under addition

H is closed under multiplication by scalar.

} These prove the statement.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

$$Ax = 0$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\frac{\text{Nul } A}{Ax=0}$$

$$\frac{\text{Col } A}{}$$

Set of all vectors
that can be written in
this form:

$$y = c_1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + c_2 \begin{bmatrix} 2 \\ 5 \end{bmatrix} + c_3 \begin{bmatrix} 3 \\ 6 \end{bmatrix}$$

$$= \begin{bmatrix} c_1 + 2c_2 + 3c_3 \\ 4c_1 + 5c_2 + 6c_3 \end{bmatrix}$$

$$y = \text{span} \{c_1, c_2, c_3\}$$

$$H = \{p(t) = a + t^2, a \in \mathbb{R}\}$$

$$(p+q)(t) = p(t) + q(t)$$

$$(cp)(t) = c \cdot p(t)$$

Is this a subspace of P_n ?

prove these $\rightarrow$

$0(t)$ always 0

$$0(t) \in H$$

$$p_1(t), p_2(t) \in H \Rightarrow (p_1 + p_2)(t) \in H$$

$$p_1(t) \in H \rightarrow (cp_1)(t) \in H$$

$$\text{Given } A = \begin{bmatrix} 1 & 3 & -2 \\ 4 & 2 & -1 \end{bmatrix}$$

- is $[1; -1; 2]$ in $\text{Nul } A$

- is $[-4; 5]$ in $\text{Col } A$?

$a + t^2$ cannot be 0 for any t . The 0 polynomial

is a polynomial that is 0 no matter what t is. t could be 1 and it's not 0. at^2 would be a 0 polynomial

$\text{Nul } A$: solutions x of $Ax=0$

$\text{Col } A$: linear combinations of cols of A .

$$\begin{bmatrix} 1 & -3 & -2 \\ 4 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \text{ in Nul } A? \text{ All vectors that are a solution to } Ax=0$$

if $= \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, then yes. If something other than $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$, no.


$$\begin{bmatrix} -4 \\ 5 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 4 \end{bmatrix} + c_2 \begin{bmatrix} -3 \\ 2 \end{bmatrix} + c_3 \begin{bmatrix} -2 \\ -1 \end{bmatrix} \text{ in Col } A?$$

$$\begin{bmatrix} -4 \\ 5 \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \begin{bmatrix} 1 & -3 & -2 \\ 4 & 2 & -1 \end{bmatrix}$$

$$A \setminus b = x$$

Simple to solve

I WANT TO BELIEVE



Subspace
Col A Span $\{a_1, a_2, \dots, a_n\}$ columns of A

Every member of the subspace is a linear combination of these

$$\mathbb{R}^2 \quad \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$$

Any vector we pick in $\mathbb{R}^2$ can be expressed as $c_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

$$\begin{bmatrix} 4 \\ -3 \end{bmatrix} = -3 \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

We want to come up w/ the smallest possible # of vectors that span the subspace. Every vector in the subspace is a linear combination of the vectors in the set.

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\text{so } \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) \right\}$$

Looking for a set of vectors linearly independent (so that it isn't a combination of the others), that spans H.

Nul A is set of all solns to $Ax=0$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3x_2 - x_3 \\ x_2 \\ x_3 \end{bmatrix} \text{ free variables, } x_2 \text{ or } x_3$$

$$= \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix} x_2 + \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} x_3$$

2 vectors that describe Nul A. B/c any soln x of $Ax=0$ can be written as a linear combination of these 2 vectors. These make up a basis for Nul A.

$$A = \begin{bmatrix} -2 & 4 & -2 & -4 \\ 2 & -6 & -3 & 1 \\ 3 & 8 & 2 & 3 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 6 & 5 \\ 0 & 2 & 5 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

With this, we know there are 2 non-pivot columns, so 2 vectors will be the basis for A.

$$Ax=0 : \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -6x_3 - 5x_4 \\ 2.5x_3 - 1.5x_4 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -6 \\ 2.5 \\ 1 \\ 0 \end{bmatrix} x_3 + \begin{bmatrix} -5 \\ -1.5 \\ 0 \\ 1 \end{bmatrix} x_4$$

basis for Nul A.

Col A =


basis(Col A) $\stackrel{?}{=} \{a_1, a_2, \dots, a_n\}$ if columns of A are not ...

linearly independent) this

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 4 & 1 \end{bmatrix}$$

this is not a base.

falls apart.



we don't need that other arrow. Just the 2 are needed.

In the $\text{rref}(A)$, each nonpivot column is a linear combination of the pivot columns.

$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ The 3rd column is a linear combination of the previous 2 $\begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

$$x = 5 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$x = 5 \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} - 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 2 \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$$

$$X = 7 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - 1 \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The vector set consisting of pivot cols(A) is a basis (use the pivot columns of A , not of $\text{ref}(A)$).

$\begin{bmatrix} 0 \\ 1 \end{bmatrix}$
 $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$
 $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$

[1], [5]

$$x = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$x = c_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$H \quad \{b_1, b_2, \dots, b_n\}$$

$$x = c_1 b_1 + c_2 b_2 + \dots + c_n b_n \quad \text{these coefficients are unique.}$$

a diff set of coefficients $\{d_1, d_2 \dots d_n\}$

$$x = d_1 b_1 + d_2 b_2 + \dots + d_n b_n$$

$$X = 0$$

$$0 = \underbrace{(d_1 - c_1)}_0 b_1 + \underbrace{(d_2 - c_2)}_0 b_2 + \dots + \underbrace{(d_n - c_n)}_0 b_n$$

So $d = \zeta$ • there is no other coefficient set.

If A is $m \times n$
 then $\text{nul } A$ contains $n \times 1$ vector
 while $\text{col } A$ contains $m \times 1$ vector

$\text{nul } A = \text{sols of } Ax=0$

$\text{col } A = \text{span } \{A \text{'s columns}\}$

$\begin{bmatrix} 1 & 2 & 0 \\ 1 & 2 & 1 \\ 1 & 2 & 2 \end{bmatrix}$ basis
 * lin. independent vectors
 * Any vector can be written as a linear combination

$$v = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} c_1 + \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} c_2 + \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} c_3$$

The number of elmts is called the dimension of the vector space.

All bases of a vector space contain the same # of elmts (dimension)

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix} x_3$$

only one free variable. Dimension of $\text{nul } A$ is 1.

For $\text{col } A$, the basis will be columns in original matrix that correspond to pivot columns. In this case, it's 2 of them.
 Dimension $\text{col } A = 2$.

of pivot columns = dimension of $\text{col } A$

of free variables = dimension of $\text{nul } A$

$\dim \text{col } A + \dim \text{nul } A = \# \text{ of columns in the matrix}$

$$\dim(\text{col } A) = \text{rank } A$$

$$\text{rank}(A) + \dim(\text{nul } A) = n.$$

Invertible Matrix Theorem update

A is invertible *

Columns of A form a basis of $\mathbb{R}^n$

$$\text{col } A = \mathbb{R}^n$$

$$\dim(\text{col } A) = n$$

$$\text{rank } A = n$$

$$\text{nul } A = \{0\}$$

$$\dim(\text{nul } A) = \{0\}$$

* $Ax=0$ has only the trivial solution. So $\text{rref}(A)$ has a pivot in every column. This is why $\text{nul } A = \{0\}$

$$A = \begin{bmatrix} 1 & 2 & 2 & 3 \\ 1 & 1 & 2 & 1 \\ 1 & 3 & 2 & 2 \end{bmatrix}$$

$$\text{rref}(A) = \begin{bmatrix} 1 & 2 & 2 & 3 \\ 0 & -1 & 0 & -2 \\ 1 & 3 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 2 & 3 \\ 0 & -1 & 0 & -2 \\ 0 & 1 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 2 & 3 \\ 0 & -1 & 0 & -2 \\ 0 & 0 & 0 & -3 \end{bmatrix}$$

$\dim(\text{col } A) = 3$
 $\dim(\text{nul } A) = 1$
 basis for $\text{col } A = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix} \right\}$
 Choose original cols from original matrix, corresponding to pivot cols.

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 1 \\ -1 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & 5 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 7 \end{bmatrix}$$

$\dim(\text{col } A) = 3$
 $\dim(\text{nul } A) = 0$
 $\text{rank } A = 3$
 basis for $\text{nul } A = \{ \}$
 basis for $\text{col } A =$

$$v_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad v_2 = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

Vectors span a plane

v_1, v_2 are linearly independent

dimension of S is 2

how many vectors need to be added to $\{v_1, v_2\}$ to make a basis for $\mathbb{R}^3$ & what would they look like?

$$\begin{bmatrix} x_1 \\ x_2 \\ 1 \end{bmatrix} \text{ where } x_3 \neq 0.$$

Every vector x in a vector space V has a unique representation as a linear combination of the elements of basis $B = \{b_1, b_2, \dots, b_n\}$

$$x = c_1 b_1 + c_2 b_2 + \dots + c_n b_n$$

$c_1, c_2, \dots, c_n$ are unique.

$$x = c_1 b_1 + c_2 b_2 + \dots + c_n b_n$$

This allows us to regard the basis as a coordinate system w/ $(c_1, c_2, \dots, c_n)$ the coordinates of vector x (relative to B) we write $[x]_B = (c_1, c_2, \dots, c_n)$ & call this the B -coordinate vector of x .

$$\begin{bmatrix} -2 \\ 5 \end{bmatrix} = 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix} + (-2) \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$



$$x = \begin{bmatrix} -2 \\ 5 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \begin{aligned} c_1 &= -2 \\ c_2 &= 5 \end{aligned}$$

$-2, 5$ are the coordinates of the vector in the new system.

A vector $x = (3, 4)$, vector in $\mathbb{R}^2$.

Another basis of $\mathbb{R}^2 = B = \{(1, 0), (1, 2)\}$

B -coordinates of $x = (3, 4)$ b/c $x = b_1 + 2b_2$



These should match up

Changing the basis hasn't changed anything. It has the same form as $\mathbb{R}^2$. That's called isomorphic to $\mathbb{R}^2$.

To find if x is in H , set up $A \{c_1, c_2\} = x$

$$v_1 c_1 + v_2 c_2 = x$$

$$\begin{bmatrix} v_1 & v_2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = x$$

3×2 matrix 3×1

If you can find c_1, c_2 it would mean x is on the span $\{v_1, v_2\}$, & then you can find the B coordinates of x are c_1, c_2 (b/c B is the basis of H).

$$x = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$


$$x_B = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \cdot x_B = x$$

$$x_B = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}^{-1} x$$

Then you get back the original.

Geometrically, space $H = \text{span}\{v_1, v_2\}$
plane.



$B = \{1+t^2, t+t^2, 1+2t+t^2\}$ is a base for P_2 .

Find coordinate vector of $p(t) = 1+4t+7t^2$ relative to B .

$$1+4t+7t^2 =$$

$$c_1(1+t^2) + c_2(t+t^2) + c_3(1+2t+t^2)$$

$$1+4t+7t^2 = (c_1 + c_3) + (c_2 + 2c_3)t + (c_1 + c_2 + c_3)t^2$$

$$c_1 + c_3 = 1 \quad c_2 + 2c_3 = 4 \quad c_1 + c_2 + c_3 = 7$$

Solve & we're done.

$$\begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix}, \quad B = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \right\}$$

$$x = v_1 + c_1 v_2 + c_2 v_3$$


STUCKERS WOO!

Thursday, November 09, 2006
8:03 AM


$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 2 & 0 & 1 & 0 & 0 \\ 3 & 0 & 0 & 44 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 3 & 0 & 0 & -44 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -44 & 0 \end{bmatrix}$$

Frequency, Time

Friday, November 10, 2006
8:02 AM



sample each time & find frequency (partition).



So we can write it to a CD.

$$\begin{aligned} t &\rightarrow p(t) \\ t_2 &\rightarrow p(t_2) \\ t_3 &\rightarrow p_3 \\ t_4 &\rightarrow p_4 \\ t_n &\rightarrow p_n \end{aligned}$$

Multiply it by a constant c if you want it to sound louder.

$c > 1$ inc. volume

$c < 1$ dec. volume

Z signals, 2 pieces of music

p_n q_n if you add, you mix 2 signals together

$\{p_1, p_2, p_3, \dots, p_n\}$ signal 1

$\{q_1, q_2, q_3, \dots, q_n\}$ signal 2

$r = c_1 p + c_2 q$ new signal, part of the set of signals

$r = \{r_1, r_2, \dots, r_n\}$

basis of vector space $\{b_1, \dots, b_n\}$

p can be written like this $\rightarrow$

r can too

q can as well.

All signals can be written as linear combinations of the bases.

Sampling occurs 44,100 times per second. Times are very close together.

input A matrix

output: are ~~to~~ cols A linearly independent (T/F)

$\dim \text{null } A$

solution to $Ax=0$

function [isind, dimnul, sol] = indep(A)

%INDEP...

% help

% k pivots

% dim nul A = n - k

% dim col A = k

% if k < n

% isind = true

% else isind = false

X-FILES
I WANT TO
BELIEVE

% if A looks like this $\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -x_4 - x_2 \\ x_2 \\ -x_4 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} x_2 + \begin{bmatrix} -1 \\ 0 \\ -1 \\ 1 \end{bmatrix} x_4$$

$$\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 6 \\ 0 \\ 0 \end{bmatrix}$$

For each individual equation, we can express it in matlab

$$m(r, :) \cdot x = 0$$

$$1x_1 + 1x_2 + 0x_3 + 1x_4 = 0$$

Make all free variables 1. We don't know which ones are (assume we don't.)

$$\text{so } 1(1) + 1(1) + 0 + 1(1) = 0$$

$$3 = 0 \quad \emptyset$$

By making $x_1 = 1$, we made a mistake.

We're concentrating on finding x_1 , so

we ~~can~~ have to revise it

$$-(m(r, :) \cdot x - 1) = -1 + 1(1) + 1(1) + 0 + 1(1) = 0$$

this gives us our x_1 value.



examine each row, but start at the diagonal.

function _____

% stuff
% stuff

% check arg.

% convert to rref

% num-pivots = 0

% dimnul = if cols A

% for each row

% start at main diagonal, going right

% check whether it's 1.

% if yes, update num-pivots & dimnul & solution.

% check num-pivots

% if == if of cols A

% isind == true

% else

% isind == false

% end

nargchk(1,1,nargin);

error(msg);

nargchk(1,3,nargout)

error(msg)

M = rref(A)

[rows cols] = size(A)

num-pivots = 0

if(nargout > 1)

dimnul = cols

if(nargout > 2)

% only if user
specifies that
he wants it.

Sol = ones(cols,1)

end
end

for ii = 1:rows

for jj = 1:cols

if (M(ii,jj) == 1)

num-pivots = num-pivots + 1

if nargout > 1

dimnul = dimnul - 1

] if nargout > 2

Sol(jj) = 1 - M(ii,jj)*

ones(cols,1)

end

end

break

end

end

if num-pivots == cols

isind = true

else

isind = false

end

Dot Product


Tuesday, November 14, 2006
8:04 AM

$$\begin{aligned}y &= ax + b \\y_2 &= ax_2 + b \\y_3 &= ax_3 + b \\y_4 &= ax_4 + b\end{aligned}$$

$$y - ax + b = 0$$

If you go out & collect data, & get diff y 's & x 's.

4 equations & 2 unknowns



You're trying to find a .
If the line isn't exactly on the points, then you have an error. It's the distance between the pt. and the line.



You want those distances, overall, to be as small as possible.

Inner Product / dot product

$$\begin{aligned}\vec{u} \cdot \vec{v} &= \vec{u}_1 \vec{v}_1 + \vec{u}_2 \vec{v}_2 + \dots + \vec{u}_n \vec{v}_n \\ \vec{u} \cdot \vec{v} &= \vec{u}^T \vec{v}\end{aligned}$$

$$\begin{aligned}\vec{u} &= \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} & \vec{v} &= \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \\ \vec{u}^T &= [u_1 \ u_2 \ \dots \ u_n] & \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} &= \vec{v}\end{aligned}$$

$$\vec{u} \cdot \vec{u} \geq 0$$

Length / norm of a vector $\vec{v}$ is defined as $\|\vec{v}\| = \sqrt{\vec{v} \cdot \vec{v}}$
 $= \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$

A unit vector is a vector w/ length = 1

Normalizing a unit vector $\vec{v}$ means finding a unit vector $\vec{u}$ that has the same direction as $\vec{v}$





$$\text{distance} = \sqrt{(v_1^2 + v_2^2 + v_3^2 + \dots)^2} = 1,$$



$$\sqrt{\|v\|^2}$$

The distance between 2 vectors $\vec{u}, \vec{v}$ is the length of $\vec{u} - \vec{v}$

$$\text{dist}(\vec{u}, \vec{v}) = \|\vec{u} - \vec{v}\|$$



$$\|\vec{u} - \vec{v}\| = \|\vec{u} + \vec{v}\|$$

If $\perp$.

$$\|\vec{u} - \vec{v}\| = \|\vec{u} + \vec{v}\|$$

$$\|\vec{u} - \vec{v}\|^2 = \|\vec{u} + \vec{v}\|^2$$

$$(\vec{u} - \vec{v}) \cdot (\vec{u} - \vec{v}) = (\vec{u} + \vec{v}) \cdot (\vec{u} + \vec{v})$$

$$\vec{u} \cdot (\vec{u} - \vec{v}) - \vec{v} \cdot (\vec{u} - \vec{v}) = \vec{u} \cdot (\vec{u} + \vec{v}) + \vec{v} \cdot (\vec{u} + \vec{v}) \quad ((\vec{u} + \vec{v}) \cdot \vec{w} = \vec{u} \cdot \vec{w} + \vec{v} \cdot \vec{w})$$

$$\vec{u} \cdot \vec{u} - \vec{v} \cdot \vec{u} - \vec{v} \cdot \vec{u} + \vec{v} \cdot \vec{v} = \vec{u} \cdot \vec{u} + \vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{u} + \vec{v} \cdot \vec{v}$$

$$-\vec{v} \cdot \vec{u} - \vec{v} \cdot \vec{u} = \vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{u}$$

$$-2\vec{u} \cdot \vec{v} = 2\vec{u} \cdot \vec{v}$$


$$0 = 4\vec{u} \cdot \vec{v}$$

$$0 = \vec{u} \cdot \vec{v}$$

so they're $\perp$ when $\vec{u} \cdot \vec{v} = 0$

orthogonal

The zero vector is orthogonal to everything.



$$z^2 = x^2 + y^2$$

$$\text{distance } z = \sqrt{x^2 + y^2}$$

$$\|\vec{u} - \vec{v}\|^2 = \|\vec{u}\|^2 + \|\vec{v}\|^2 = \|\vec{y} - \vec{x}\|^2 = (y-x)(y-x) \quad \text{or } \perp$$

$$y \cdot y - 2y \cdot x + x \cdot x$$

$$\|y\|^2 + \|x\|^2 - 0$$

If a vector $\vec{v}$ is orthogonal to every vector in a subspace H of $\mathbb{R}^n$, then $\vec{v}$ is $\perp$ to H .

The set of all vectors $\vec{v}$ that are $\perp$ to H form the orthogonal complement of H , written $H^\perp$.

I LIKE
X-MEN
I DON'T

the orthogonal complement of H , written $H^\perp$
 $(\text{Row } A)^\perp = \text{Nul } A \leftarrow \text{sols to } Ax=0$
 $\uparrow$
 rowspace (space spanned by rows of a matrix)

I DON'T
LIKE
MATLAB

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$A \qquad \qquad X$

All solutions x of $Ax=0$
 So $(\text{Row } A)^\perp = A \cdot x = 0$ b/c it's $\perp$.

$$\begin{aligned} a_{11}x &= 0 \\ a_{21}x &= 0 \\ a_{31}x &= 0 \end{aligned}$$

$$\begin{aligned} y &= c_1 a_{11} + c_2 a_{21} + c_3 a_{31} \\ y \cdot x &= (c_1 a_{11} + c_2 a_{21} + c_3 a_{31}) \cdot x \\ &= c_1 (a_{11} \cdot x) + c_2 (a_{21} \cdot x) + c_3 (a_{31} \cdot x) \\ 0 &= c_1 (0) + c_2 (0) + c_3 (0) \end{aligned}$$

Any vector you take in
 $\text{Row } A$ is orthogonal
 to x b/c they're all
 0, regardless of c 's.

$(\text{Col } A)^\perp = \text{Nul } A$
 orthogonal complement



$\|u\|\cos\theta$ projection of $\vec{u}$ onto $\vec{v}$

$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos\theta$ when $\vec{u}$ & $\vec{v}$ are
 orthogonal, dot product = 0
 when $\vec{u}$ & $\vec{v}$ are same vector, $\|\vec{u}\| = \|\vec{v}\|$
 are equal.

I WANT TO SEE
X-FILES AGAIN!!!

Let $\vec{v} = (a \ b)$ describe vectors $\perp$ to $\vec{v}$.

$$\begin{bmatrix} x \\ y \end{bmatrix}$$

$$\begin{bmatrix} a \\ b \end{bmatrix}$$

$$ax + by = 0$$

$$ax = -by$$

x is
a multiple
of $-b$

y is a multiple of a

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} \cdot c$$




$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} \cdot c$$



$$\{(x, y) : x^2 + y^2 = 1\}$$

$$\begin{aligned} \|\vec{v}\| &= \sqrt{\vec{v} \cdot \vec{v}} \\ &= \sqrt{x+x \cdot x + y+y} = 1 \end{aligned}$$



0 is not in the set, b/c $0^2 + 0^2 \neq 1$
so it is not a subspace.


Let y be a vector orthogonal to u & v
Show that y is orthogonal to $u+v$

$$\vec{y} \cdot \vec{u} = 0$$

$$\vec{y} \cdot \vec{v} = 0$$

$$\vec{y} \cdot (\vec{u} + \vec{v}) = \vec{y} \cdot \vec{u} + \vec{y} \cdot \vec{v} = 0 + 0 = 0$$

1




$c_1 \vec{u} + c_2 \vec{v} = \text{linear combination vector}$
And if you do all linear comb's
then you get a plane.

If x is both in W & its orthogonal complement, what can you say about x ?

$$\vec{u} \cdot \vec{v} = 0$$

$$\vec{u} \cdot \vec{v} = 0$$

x must be $\vec{0}$ vector.



Let $\vec{u} = (5, -6, 7)$ & W be the set of all vectors $\vec{v}$ in $\mathbb{R}^3$ s.t. $\vec{u} \cdot \vec{v} = 0$. Is W a subspace of $\mathbb{R}^3$?

It's a Nullspace. $Ax = 0$

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta$$

Given a set of vectors $V = (v_1, v_2, \dots, v_n)$ in $\mathbb{R}^n$ such that $v_i \cdot v_j = 0$, for any $2 \leq i, j \in V$

(Right hand rule, 3 directions $\rightarrow$)

Then V is called an orthogonal set.

Ex. $V = ((0, 0, 1), (0, 1, 0), (1, 0, 0))$

$$V = \left\{ \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} \\ 2 \\ 3.5 \end{pmatrix} \right\}$$

Furthermore, V is linearly independent

This makes it a basis of $\mathbb{R}^3$

We call it an orthogonal basis.

We introduce the idea of basis b/c it's more efficient, & b/c we express any vector as a linear combination w/ unique coefficients, we have unique combo of coefficients

Let $\{v_1, v_2, \dots, v_n\}$ be an orthogonal basis for a subspace H of $\mathbb{R}^n$. Then every elmt $x \in \mathbb{R}^n$ is written as linear combo

$$x = c_1 v_1 + \dots + c_n v_n$$


where $c_i = \frac{x \cdot v_i}{v_i \cdot v_i}$

The whole middle set will vanish if you multiply by v_i , except for $c_i v_i$

$$x \cdot v_i = c_1 v_1 \cdot v_i + c_2 v_2 \cdot v_i + \dots + c_n v_n \cdot v_i$$


$$x \cdot v_i = c_i v_i \cdot v_i$$

$$\frac{x \cdot v_i}{v_i \cdot v_i} = c_i$$



$$x = c_1 v_1 + c_2 v_2 \quad \text{where } c_i = \frac{x \cdot v_i}{v_i \cdot v_i}$$

Geometric representation



$c_1 v_1$ is the projection of x on v_1


This is a vector w/ same direction as v_1 , & length $\|x\| \cos(\theta)$

$$\text{From } x \cdot v_1 = \|x\| \|v_1\| \cos(90 - \theta)$$

We get that length of $c_1 v_1$ is $\frac{x \cdot v_1}{\|v_1\|}$

$$c \cdot v_1 = \frac{x \cdot v_1}{\|v_1\|} \cdot \frac{v_1}{\|v_1\|} = \frac{x \cdot v_1}{\|v_1\|^2} \cdot v_1 = \frac{x \cdot v_1}{v_1 \cdot v_1} \cdot v_1$$

Given a nonzero vector v , decompose a vector x into the sum of 2 vectors: one a multiple of v , & another orthogonal to v



$$x = \hat{v} + z$$

where $\hat{v} = c\hat{v}$


& z is vector orthogonal to $\hat{v}$



where $\vec{v} = c\vec{u}$
 $\vec{z}$ is vector orthogonal to $\vec{v}$

$\vec{v}$ & $\vec{z}$ are orthogonal, so will make up a basis - $x - \vec{v} = \vec{z}$

EX. Let $\vec{y} = (1, 7)$ & $\vec{v} = (-4, 2)$. Compute orthogonal projection of y through $\vec{v}$ & the origin.



$$y = \begin{bmatrix} 1 \\ 7 \end{bmatrix} \quad \vec{v} = \begin{bmatrix} -4 \\ 2 \end{bmatrix}$$

$$y = c \cdot \vec{v} + \vec{w}$$

$$\frac{y \cdot \vec{v}}{\vec{v} \cdot \vec{v}} = \frac{y \cdot \vec{v}}{\|\vec{v}\|^2}$$

$$y \cdot \vec{v} = -4 + 14 = 10$$

$$\|\vec{v}\|^2 = 16 + 4 = 20$$

$$c = \frac{10}{20} = \frac{1}{2}$$

$$(1, 7) = \frac{1}{2}(-4, 2) + \vec{w}$$

$$(1, 7) = \boxed{(-2, 1)} + \vec{w}$$


projection

cuz they are stretching over the axes!

Yeah!

free
I wonder why

EX2 Let $y = (2, 3)$ & $\vec{v} = (4, -7)$ write y as the sum of 2 orthogonal vectors, one in $\text{span}(v)$ & one orthogonal to $\vec{v}$.



$$\vec{y} = \vec{v} \cdot w$$

in $\text{span}\{\vec{v}\}$

span of a single vector is a line, $c\vec{v}$

$$V = \text{span}\{b_1, b_2, \dots, b_n\}$$

$$x \in V$$

$$x = c_1 b_1 + c_2 b_2 + \dots + c_n b_n$$

$$\vec{y} = (c\vec{v}) + \vec{w}$$

$$\frac{y \cdot \vec{v}}{\vec{v} \cdot \vec{v}}$$

then mult. this by $\vec{v}$
 solve for $\vec{w}$.

Orthogonal Sets, Dot Products

Thursday, November 16, 2006

8:09 AM

Define orthogonal set $\{v_1, v_2, \dots, v_n\}$ in $\mathbb{R}^n$ as a set of vectors whose pair-wise dot products are zero
(inner product)

If all vectors in an orthogonal set are also unit vectors, then it is called an orthonormal set. The vectors have been normalized.

$$\{v_1, v_2\}$$

$$\left\{ \frac{v_1}{\|v_1\|}, \frac{v_2}{\|v_2\|} \right\} \text{ still a basis, still linearly independent.}$$

If an orthonormal set V spans a subspace H , then V is called the orthonormal basis of H .

The dot product between every vector and itself is 1.

$$U^T = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \\ u_{31} & u_{32} \end{bmatrix} \text{ instead of } U = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \\ u_{31} & u_{32} \end{bmatrix}$$

$$U^T \cdot U = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

If a square matrix U has orthonormal columns, it is invertible & its inverse is just the transpose. We call that an orthogonal matrix.

An $m \times n$ matrix U has an orthonormal columns, iff $U^T U = I$.

If U is an $m \times n$ matrix w/ orthonormal columns, & $\vec{x}, \vec{y}$ are vectors in $\mathbb{R}^n$, then:

$$\|U\vec{x}\| = \|\vec{x}\|$$

transformation $\vec{x} \rightarrow U\vec{x}$ preserves the length of $\vec{x}$.

$$(U\vec{x}) \cdot (U\vec{y}) = \vec{x} \cdot \vec{y}$$

$$(U\vec{x}) \cdot (U\vec{y}) = 0 \text{ iff } \vec{x} \cdot \vec{y} = 0$$

Preserves orthogonality of $\vec{x}$ & $\vec{y}$.

To see if 2 vectors are orthogonal, matrix multiplication

$$U = [u_1 \ u_2 \ u_3 \ \dots \ u_n]$$

$$U^T \cdot U = \text{dot product}$$

We have defined an orthogonal basis for subspace W as a basis for W that is also an orthogonal set.

$$c_i = \frac{x \cdot v_i}{v_i \cdot v_i}$$
 If $B = \{v_1, v_2, \dots, v_m\}$ is an orthogonal basis for subspace W of $\mathbb{R}^n$, those coefficients c are:

Every element x of W can be written as a unique linear combination of the elements of B

$x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$ where $c_i = \frac{x \cdot v_i}{v_i \cdot v_i}$

$z \cdot \hat{y} = 0$ $\leftarrow z \perp \hat{y}$

$z \in W^\perp$

y is decomposed into its components.

$\hat{y}$, orthogonal projection of y on the subspace is part of the set.


We can easily compute $\hat{y}$. All we need to know is an orthogonal basis of W . $y = \hat{y} + z$

$$\hat{y} = \frac{v_1 (y \cdot v_1)}{v_1 \cdot v_1}$$

if $B = \{v_1, \dots, v_m\}$ is an orthogonal basis of W
 W is a line in $\mathbb{R}^n$

If you're given 2 vectors:

$$\hat{y} = \hat{y}_1 + \hat{y}_2 = \frac{v_1 (y \cdot v_1)}{v_1 \cdot v_1} + \frac{v_2 (y \cdot v_2)}{v_2 \cdot v_2}$$



$\|y - \hat{y}\|$ shortest distance between y and W and so $\hat{y}$ is best approx. $\hat{y}$ to y by elmts of W .

$\hat{y}_1$ is orthogonal projection of y onto $\text{span}\{v_1\}$
 $\hat{y}_2$ is orthogonal projection of y onto $\text{span}\{v_2\}$

If $B = \{v_1, v_2\}$ is an orthonormal basis of W

$$\text{then } v_1 \cdot v_1 + v_2 \cdot v_2 = 1$$

$$\hat{y}_1 = v_1 (y \cdot v_1)$$

$$\hat{y}_2 = v_2 (y \cdot v_2)$$

$$\hat{y} = \hat{y}_1 + \hat{y}_2 = v_1 (y \cdot v_1) + v_2 (y \cdot v_2)$$

$$= [v_1 \ v_2] [y \cdot v_1 \ y \cdot v_2]^T$$

$$[v_1 \ v_2] [v_1 \ v_2]^T [y] = B B^T y$$

$$\begin{bmatrix} | & | & \dots & | \\ v_1 & v_2 & \dots & v_p \\ | & | & \dots & | \end{bmatrix} \begin{bmatrix} -v_1- \\ -v_2- \\ -v_p- \end{bmatrix} \begin{bmatrix} y \end{bmatrix}$$

$U \qquad U^T$

$\text{proj}_W y = U \cdot U^T \cdot y$ where $\{v_1, v_2, \dots, v_p\}$ is an orthonormal basis of W



When something is orthonormal, v_1 & v_2 are = to

eh... i guess
 I y listening?

$$\vec{v}_1 = (1, 1, 0) \quad \vec{v}_2 = (-1, 1, 0), \quad y = (-1, 4, 3)$$

Find orthogonal proj of y onto $\text{span}\{v_1, v_2\}$

$\text{Span}\{v_1, v_2\}$ is a plane

$$v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

They are linearly independent, b/c one is not a multiple of the other. or

If v_1 & v_2 are dependent, it's a line. If they are

not a multiple of the other. or
 we could do $c_1 \vec{u}_1 + c_2 \vec{u}_2 = 0$ where
 the only soln for c_1, c_2 is the
 trivial one.

line. If they are
 linearly independent,
 it's a plane -

$\vec{u}_1 \cdot \vec{u}_2 = 0$ and so this is sufficient b/c if one was
 a multiple of the other you wouldn't get 0.

And they are also $\perp$, dot product being zero.

To find the projection, first normalize the vectors.

$$\vec{u}_1 = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right)$$

$$\vec{u}_2 = \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \right)$$

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 4 \\ 3 \end{bmatrix} \quad (\text{Proj})$$

$\vec{u}_1 = (1, 1, 0)$ $\vec{u}_2 = (-1, 1, 0)$ $\vec{y} = (-1, 4, 3)$ write
 as the sum of a vector in $W = \text{Span}\{\vec{u}_1, \vec{u}_2\}$ & a vector
 in $W^\perp$.

$$\vec{y} = \hat{\vec{y}} + \vec{z} \quad \vec{z} \text{ is a vector in } W^\perp$$

$$\vec{y} - \hat{\vec{y}} = \vec{z}$$

$$(-1, 4, 0) - (-1, 4, 3) = (0, 0, -3)$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 4 \\ 3 \end{bmatrix} = \begin{bmatrix} -1 \\ 4 \\ 0 \end{bmatrix}$$

$\|\vec{z}\|$ = closest distance between $\vec{y}$ & $W^\perp$

$$\text{sq}(0)^2 + \text{sq}(0)^2 + \text{sq}(-3)^2 = 9$$

$\vec{u}_1 = (1, 1, 2)$ $\vec{u}_2 = (-1, 1, 0)$ $\vec{y} = (-1, 4, 3)$ Find orthogonal
 projection of $\vec{y}$ onto $W = \text{Span}\{\vec{u}_1, \vec{u}_2\}$

$\vec{y}$ is in W therefore its orthogonal projection will
 be $\vec{y}$ itself

$\vec{z}$ will be zero vector.

$\vec{u}_1 = (1, 1, -2)$ $\vec{u}_2 = (5, -1, 2)$, $\vec{y} = (0, 0, 1)$

Verify that $\vec{y}$ is not in the span of $W = \text{Span}\{\vec{u}_1, \vec{u}_2\}$


$\vec{y}$ is not a lin. comb. of $\vec{u}_1, \vec{u}_2$: reduce augmented
 matrix of $[\vec{u}_1, \vec{u}_2, \vec{y}]$

Use γ to find a vector of v in $\mathbb{R}^3$ that is orthog. to v_1 & v_2 .

The zero vector is always orthogonal.

Besides the zero vector, the dot products of each of them is = to zero, & the dot products set equal to each other will find you a vector.

$v = c(0, 2, 1)$ where c is a scalar (free variable)



we can use γ to find orthogonal vector on the space ($\vec{\gamma}$) & then subtract to find z .

$$\gamma = \vec{\gamma} + z$$

so find $\vec{\gamma}$ & then subtract from γ .

Matlab

Given a matrix A , show columns are orthogonal
Normalize each column of A to get U

$$D = (A') * A =$$

```
% Normalize for ii = 1 : size(A, 2)
    U(:, ii) = A(:, ii) / norm(:, ii)
end
```

closest point to γ in col U . $\gamma = (1, 1, 1, 1, 1, 1, 1)$

col U is space spanned by cols of U .

(cols of U are orthonormal set so you just have to multiply $U * U' * \gamma = \gamma$ that in Matlab


Generating Orthogonal Basis

Tuesday, November 21, 2006


8:02 AM

Generating an orthogonal basis Given $\{u_1, u_2\}$ of a subspace W , how can we come up w/ an orthogonal basis for W ?

Idea: we need to come up w/ a set $\{u_1, \text{vector orthogonal to } u_1\}$



$\{v_1, v_2\}$ where $v_1 = u_1$, $v_2 = \text{vector orthogonal to } u_1$




$$u_2 = \hat{u}_2 + z$$

$$v_1 = u_1$$

$$v_2 = u_2 - \frac{u_2 \cdot v_1}{v_1 \cdot v_1} v_1$$

orthogonal basis $\{u_1, u_2, u_3\}$?



$\{u_1, u_2, u_3\}$

We're looking for, given $u_1, u_2, \text{ and } u_3$


the basis that contains $\{u_1, \text{vector orthogonal to } u_1, \text{ vector orth. to } u_1, \text{ vector orth. to } u_1, \text{ vector orth. to } u_1\}$

b/c we want them to be an orthogonal set $\rightarrow v_1, v_2, v_3$

Subspace $\text{span}\{u_1, u_2, u_3\}$

Space spanned by $\{u_1, u_2\}$

by $\{u_1, u_2\}$



$$u_3 = \hat{u}_3 + z \quad z = v_3$$

$$\{v_1 = u_1, v_2 = u_2 - \frac{u_2 \cdot v_1}{v_1 \cdot v_1} v_1, v_3 = u_3 - \left(\frac{u_3 \cdot v_1}{v_1 \cdot v_1} v_1 + \frac{u_3 \cdot v_2}{v_2 \cdot v_2} v_2 \right)\}$$

$$v_1 = u_1$$

$$v_2 = u_2 - \text{proj}_{W_1} u_2, \quad W_1 = \text{span}\{u_1\}$$

$$v_3 = u_3 - \text{proj}_{W_2} u_3, \quad W_2 = \text{span}\{u_1, u_2\}$$

$$\begin{aligned}
 v_2 &= v_2 - \text{proj}_{w_1} v_2, \quad w_1 = \text{span}\{v_1\} \\
 v_3 &= v_3 - \text{proj}_{w_2} v_3, \quad w_2 = \text{span}\{v_1, v_2\} \\
 v_{k+1} &= v_{k+1} - \text{proj}_{w_k} v_{k+1} \\
 w_k &= \text{span}\{v_1, v_2, \dots, v_k\}
 \end{aligned}$$

This is the Gram-Schmidt Process for finding an orthogonal basis

A: $l \times n$ ind. columns

$A = QR \rightarrow$ upper triangular invertible matrix

$\uparrow$
matrix whose columns form orthogonal basis of $\text{col } A$

$$\begin{array}{cccc}
 a_{11} & a_{12} & \dots & a_{1n} \\
 a_{21} & a_{22} & \dots & a_{2n} \\
 \vdots & \vdots & & \vdots \\
 a_{m1} & a_{m2} & \dots & a_{mn} \\
 \uparrow & \uparrow & & \uparrow \\
 v_1 & v_2 & & v_n
 \end{array}$$

A

$$\begin{array}{cccc}
 q_{11} & q_{12} & \dots & q_{1n} \\
 q_{21} & q_{22} & \dots & q_{2n} \\
 \vdots & \vdots & & \vdots \\
 q_{m1} & q_{m2} & \dots & q_{mn} \\
 \uparrow & \uparrow & & \uparrow \\
 v_1 & v_2 & & v_n
 \end{array}$$

$Q \rightarrow$ orthonormal basis of A .

Want... exact
... need...
Something...
not... being

which we
got like we
did above.

$$A = [v_1 \ v_2 \ \dots \ v_n]$$

vector
in $\mathbb{R}^m$

$$Q = \left[\frac{v_1}{\|v_1\|} \ \frac{v_2}{\|v_2\|} \ \dots \ \frac{v_n}{\|v_n\|} \right] \begin{bmatrix} r_{11} & & & \\ 0 & r_{22} & & \\ & 0 & r_{33} & \\ & & 0 & r_{nn} \end{bmatrix}$$

$$A = QR$$

$$v_1 = r_{11} v_1 + 0 v_2 + \dots + 0 v_n$$

$$v_2 = r_{12} v_1 + r_{22} v_2$$


$$v_1 = v_1$$

$$v_2 = v_2 - \text{proj}_{w_1} v_2$$

v_2 is a lin.
comb. of v_1 &
 $v_1 = v_1$

v_1 belongs in
 w_1 & in w_2
because it's
just a line
intersecting w/
the plane created
w/ v_3 .

Last night,
Mulder was:
almost abducted
kidnapped
brainwashed
returned



fooled into believing he
was onto something
rescued by scully.

$$Q^T A = (Q^T Q) R$$

I

Today's
lecture

$$v_3 \in W_3,$$

just like

$$v_1 \in W_2.$$

$$v_3 \in W_3$$

$$\text{b/c } v_3 \in W_3$$

$$\text{or } \text{proj}_{W_2} v_3 \in W_2$$

$$* \text{ not } W_2$$

$$A = \begin{bmatrix} 3 & 5 & 1 \\ -1 & 5 & 2 \\ 3 & -7 & 8 \end{bmatrix}$$

$$v_1 = (3, 1, -1, 3)$$

$$v_2 = (-5, 1, 5, -7)$$

$$v_3 = (1, 1, -2, 8)$$

$$v_1, v_2, v_3$$

$$v_1 = (3, 1, -1, 3)$$

$$v_2 = v_2 - \frac{v_2 \cdot v_1}{v_1 \cdot v_1} v_1 = v_2 - (-2)v_1 = (1, 3, 3, 1)$$

okay... more than 1
episode.

$$v_3 = v_3 - \left(\frac{v_3 \cdot v_1}{v_1 \cdot v_1} v_1 + \frac{v_3 \cdot v_2}{v_2 \cdot v_2} v_2 \right) = v_3 - \left(\frac{3}{2} v_1 + \frac{1}{2} v_2 \right) = (-3, 1, 1, 3)$$

If you wanted to decompose
A into Q + R, normalize it.

$$v_1' = \left(\frac{3}{\sqrt{20}}, \frac{1}{\sqrt{20}}, \frac{1}{\sqrt{20}}, \frac{3}{\sqrt{20}} \right)$$

$$v_2' = \left(\frac{1}{\sqrt{20}}, \frac{3}{\sqrt{20}}, \frac{3}{\sqrt{20}}, -\frac{1}{\sqrt{20}} \right)$$

$$v_3' = \left(-\frac{3}{\sqrt{20}}, \frac{1}{\sqrt{20}}, \frac{1}{\sqrt{20}}, \frac{3}{\sqrt{20}} \right)$$

$$Q = [v_1' \ v_2' \ v_3']$$

$$R = Q^T A$$

$$v_1 \cdot v_1 = 20$$

$$Q^T A = \begin{bmatrix} \frac{3}{\sqrt{20}} & \frac{1}{\sqrt{20}} & -\frac{3}{\sqrt{20}} \\ \frac{1}{\sqrt{20}} & \frac{3}{\sqrt{20}} & \frac{1}{\sqrt{20}} \\ \frac{1}{\sqrt{20}} & \frac{3}{\sqrt{20}} & \frac{1}{\sqrt{20}} \end{bmatrix}$$

One episode,
Mulder was:

- Attacked by a genetic mutant
- Fell in a hole
- Lost his weapon
- Broke his arm
- Rescued by scully.

Another episode,
Mulder was:

- Tricked by a psychic psycho.
- Shot
- Rescued by scully.

Least Squares Problems

Overdetermined systems of equations.

Dimensions of A are $m \times n$ where $m > n$ A is a full rank matrix. Rank of A is n

$$\text{Rank}(A) = n$$

So it has a soln b/c



there's a pivot in every column.

$$Ax = b$$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$


$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$b = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$$

Col A span W if b is not a lin. comb. of columns of A (no x that will make that true), then the relationship w/ $b \in \text{Col } A$ is b is not on the plane. Ax is instead $= b + r$, where r is error.the closest vector in W

$$A\hat{x} = \hat{b}$$

So we're looking for the solution for this system

 $\hat{z}$ represents error that we have.

you have to minimize these distances.

$$y_1 = ax_1 + b$$

$$y_2 = ax_2 + b$$

$$\vdots$$

$$y_p = ax_p + b$$

Data fitting

F. t the given data to a function.

Polynomial interpolation.

$$p(x) = c_0 + c_1x + c_2x^2 + \dots + c_{n-1}x^{n-1}$$

to data $(x_1, y_1), (x_2, y_2), \dots, (x_m, y_m)$

$$p(1980) = 1025$$

$$1025 = 1980 + c_1(1980) + c_2(1980)^2 + \dots + c_{n-1}(1980)^{n-1}$$

$$[1 \ 1980 \ 1980^2 \ \dots \ 1980^{n-1}] [c_0 \ c_1 \ c_2 \ \dots \ c_{n-1}] = [1025]$$


1996 was

4th season

$$\begin{bmatrix} 1 & 1984 & \dots & \dots \end{bmatrix} - \int \begin{bmatrix} c_1 \\ \vdots \\ c_{n-1} \end{bmatrix} = \begin{bmatrix} 999 \\ \vdots \end{bmatrix}$$

There will probably not be a soln b/c $m > n$, so find approx.

Navigation - given the distances between a pt. & known position, find location of the pt.



You have distance between $B_1, B_2, B_3, \& B_4$, & you're trying to find x .
3 equations, 3 unknowns, & you approximate.

The set of least squares soln to $Ax = b$ is the same as the non-empty set of solns of the normal equations $A^T A x = A^T b$. $A^T A$ is not necessarily invertible.

The columns of A are linearly independent the soln is unique & is given by $x_{LS} = (A^T A)^{-1} A^T b$

Prove

x_{LS} is a soln to the system $A^T A x = A^T b$

Given A & b , by the Best Approx Theorem, the orthogonal projection b_p of b onto $\text{Col } A$ is the closest point in $\text{Col } A$ to b .

$$\|b - b_p\| < \|b - y\| \text{ for any } y \neq b_p \text{ in } \text{Col } A.$$

Since b_p is in $\text{Col } A$, the sys: $Ax = b_p$ has a soln x_{LS} .
The above becomes $\|b - Ax_{LS}\| < \|b - Ax\|$. x_{LS} is a least squares soln.

By the Orthogonal Decomp. Theorem, the vector $b - Ax_{LS}$ is orthogonal to $\text{Col } A$, therefore $A^T(b - Ax_{LS}) = 0$

$$\begin{aligned} \Rightarrow A^T b - A^T A x_{LS} &= 0 \\ \Rightarrow A^T A x_{LS} &= A^T b \end{aligned}$$



Prove

$A^T A$ is invertible

$A^T A x = 0$ has only one soln (which proves it.)

$$A^T A x = 0 \Rightarrow x^T A^T A x = 0$$

$$(Ax)^T Ax = 0$$

$$\|Ax\|^2 = 0 \text{ (definition of length)}$$

$$Ax = 0 \text{ (if } \|x\| \neq 0, \text{ then } v = 0)$$

$$x = 0 \text{ (b/c } A \text{'s columns are lin. indep.)}$$

Since A is invertible, the system has



$x=0$ (b/c A 's column

Since A is invertible, the system has a solution & it must be zero.




$$y = x\beta + c$$

Linear model

x is called the design matrix

β is called the parameter vector

y is called the observation vector



one minute later



β_0, β_1 are regression coefficients when equation is a line & is called line of regression.

We want to avoid computing the inverse

If A is lin. ind. columns, apply Gram-Schmidt process w/ normalization to columns of A will make it possible to write A as product QR where Q is an $m \times n$ matrix whose columns form an orthogonal basis for $\text{col}(A)$. $Q^T Q = I$.

It is an upper triangular matrix which is invertible.
QR factorization.

$$A^T A x = A^T b$$

$$(QR)^T QR x = (QR)^T b$$

$$R^T Q^T A x = R^T Q^T b$$

$$R^T R x = R^T Q^T b \quad (R \text{ is invertible})$$

$$R x = Q^T b$$

Since R is upper triangular, we can solve quickly & easily using back substitution.

Smallest error in set

Wednesday, November 29, 2006
8:08 AM

$$y = Xb + \epsilon \quad \text{where } \epsilon \text{ is error}$$

$$b_0 + b_1 x + b_2 x^2 + b_3 x^3 \dots + b_{n-1} x^{n-1}$$

$$X = \begin{bmatrix} 1 & x & x^2 & \dots & x^{n-1} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & x_m & x_m^2 & \dots & x_m^{n-1} \end{bmatrix} \quad Y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$

$$\begin{bmatrix} b_0 \\ b_1 \\ \vdots \\ b_{n-1} \end{bmatrix} \leftarrow \text{unknown}$$

$$y - Xb = \epsilon$$

$$\begin{matrix} \|\epsilon_1\| \\ \|\epsilon_2\| \\ \|\epsilon_3\| \end{matrix}$$

> pick which one is smallest
& choose that set of
data bc it has the smallest
error

$$y = \beta_0 f_0(x) + \beta_1 f_1(x) + \dots + \beta_n f_n(x)$$

where $f_i(x)$ are linear or non-linear functions of x .


You have y & x & you need b .

$y = Xb$ however, x is not necessarily
square. & so the system doesn't have a soln.

You need, instead, to find orthogonal projection
of y onto $\text{Col } X$. Just like orthogonal
projection of b onto $\text{Col } A$ in $Ax = b$.

$$\begin{aligned} \hat{y} &= X\hat{b} \\ \epsilon &= \hat{y} - y = \hat{y} - Xb \\ \hat{b} - b &= \hat{b} - Ax \end{aligned}$$

$Ax = b$
 $A\hat{x} = \hat{b}$



You can do it this way or use normal equations $A^T A x = A^T b$
Solve for x .

$$x = (A^T A)^{-1} A^T b$$

You can do it this way or with QR factorization

$$\begin{aligned} A &= QR \\ A^T A x &= (QR)^T (QR x) \\ &= R^T (Q^T Q) R x \\ &= R^T R x \\ A^T b &= R^T Q^T b \\ R x &= Q^T b \end{aligned}$$

$A \rightarrow$ basis $\rightarrow$ orthogonal

basis $\rightarrow$ normalize to get Q

compute R
 $R x = Q^T b$

$A = LU \rightarrow$ upper triangular matrix
 $\downarrow$
unit lower triangular


$$\begin{aligned} LUx &= b \\ Ly &= b \end{aligned}$$

$$Ux = y$$

$\underline{A} = \underline{Q} R$ what is R ?
 Q has orthonormal columns

$$\begin{aligned} Q^T A &= Q^T Q R \\ \boxed{Q^T A} &= R \end{aligned}$$

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \\ 1 & 2 \end{bmatrix}$$



If $b = [1; 0; -1]$ what can you say about least squares soln of $Ax = b$?
What is the relationship between b & columns of A ?

The dot product between columns of A & b is 0, so b is orthogonal to both columns of A . The projection of b onto A is the 0 vector.

$Ax = 0$ will give you x_{LS}

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \\ 1 & 2 \end{bmatrix}$$

$$b = \begin{bmatrix} 3 \\ 3 \\ 5 \end{bmatrix}$$

consider $u = [1; -1/2]$ & $v = [0; 1/2]$

compute Au & Av

Could u or v be a least squares soln to $Ax = b$?

$$Au = \begin{bmatrix} 0 \\ -3 \\ 0 \end{bmatrix} \rightarrow b, \quad \|b - b_1\|$$

Find the one w/

$$Av = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \longrightarrow b_2 \quad \|b - b_2\| \text{ the smallest error.}$$

$$\|b - b_1\| = 6.16$$

$$\|b - b_2\| = 5.38$$

so Av is closer to b than Au is.
 v might be a least squares soln.

If you compute Au & Av & the error is the same, then those solns are equidistant from b , & if there are 2 vectors equally close to b then they are not least squares solns $\Rightarrow$ those are unique.

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \\ 1 & 2 \end{bmatrix} \quad b = \begin{bmatrix} 3 \\ -1 \\ 5 \end{bmatrix}$$

Find the least squares soln for $Ax = b$

$$A^T A = \begin{bmatrix} 3 & 0 \\ 0 & 24 \end{bmatrix}$$

$$A^T b = \begin{bmatrix} 9 \\ 12 \end{bmatrix}$$

$$3x_1 = 9 \Rightarrow x_1 = 3$$


$$24x_2 = 12 \Rightarrow x_2 = \frac{1}{2}$$

$$x_{LS} = \begin{bmatrix} 3 \\ 1/2 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \\ 1 & 2 \end{bmatrix}$$

$$b = \begin{bmatrix} 3 \\ -1 \\ 5 \end{bmatrix}$$

Find projection of b onto $\text{Col } A$



this + this = $\text{proj}_A b$

$$\text{proj}_{A_1} b + \text{proj}_{A_2} b = \text{proj}_A b$$

$$\frac{b \cdot a_1}{a_1 \cdot a_1} a_1 + \frac{b \cdot a_2}{a_2 \cdot a_2} a_2 = \text{proj}_A b$$

$$\text{It is } b_p = A^* x_{LS} = \begin{bmatrix} \quad \end{bmatrix}$$